arXiv:1110.4868v1 [math.NT] 21 Oct 2011

MULTIPLE DIRICHLET SERIES AND SHIFTED CONVOLUTIONS

JEFF HOFFSTEIN

1. Introduction

In 1939 and 1940 in [30] and [27] Rankin and Selberg independently introduced what is now called a Rankin-Selberg convolution. Very roughly, if

$$L(s,f)=\sum_{m\geq 1}\frac{a(m)}{m^s} \text{ and } L(s,g)=\sum_{m\geq 1}\frac{b(m)}{m^s}$$

are two L-series associated to modular forms, each convergent when $\mathfrak{Re}\, s>1$, with a functional equation in s, and a meromorphic continuation to $\mathbb{C}$, the Rankin-Selberg convolution provides a functional equation and a meromorphic continuation to $\mathbb{C}$ of the new L-series

$$L(s,f\otimes g)=\sum_{m\geq 1}\frac{a(m)\bar{b}(m)}{m^s}.$$

This idea had a profound impact on the field and was responsible for what was at that time the best progress toward the Ramanujan conjecture.

In 1965, [28], in a paper summing up progress in modular forms, Selberg introduced the notion of a shifted convolution. If h is a positive integer, his shifted convolution was the series

$$\sum_{m\geq 1}\frac{a(m+h)\bar{b}(m)}{(2m+h)^s}. \tag{1.1}$$

Selberg mentioned that the meromorphic continuation of this could be obtained. (It converges absolutely for $\mathfrak{Re}\, s>1$.) He went on to remark: “We cannot make much use of this function at the moment...”.

Since then it has been generally recognized that the analytic properties of series such as (1.1) play a very important role in estimating

---

Date: October 24, 2011.

the size of certain shifted sums, such as

$$\sum_{m<x} a(m)\bar{b}(m+h),$$

and that these in turn play an important role in progress toward the Lindelöf Hypothesis, that is, the bounding from above of automorphic L-series at points inside the critical strip. See [23] for a survey of this subject.

A major factor hindering progress in this area has been a difficulty in applying the meromorphic continuation of the series (1.1) to the analysis of shifted sums. Although non-trivial results can be obtained from any continuation past the trivial line $\mathfrak{Re}\, s = 1$, technical difficulties have interfered with more refined applications.

One purpose of this paper is to suggest that the series (1.1) may not be the correct one to look at, or at any rate, might not tell the whole story. In particular, we will show that the series

$$D(s;h) = \sum_{m\geq 1} \frac{a(m+h)\bar{b}(m)}{m^s}, \tag{1.2}$$

and generalizations of it, can be meromorphically continued to all of $\mathbb{C}$. This proves to be a surprisingly difficult task and is the subject of Sections 2–5. Once done, we extend this notion to multiple shifted Dirichlet series, in which a sum is taken over the shifts h as well, and show that these as well can be meromorphically continued. One interesting aspect of these shifted multiple Dirichlet series is that just like their L-series and multiple Dirichlet series counterparts they have an analog of a critical strip where all the deep number theoretic information is buried. Once the continuations have been established, some strong results on shifted sums, in which there is significant cancellation uniformly in the length and the shift parameter are obtained(Theorem 1.4). From these estimates subconvexity bounds (Theorem 1.1) and mean values of L-series (Theorem 1.2) follow with little difficulty. The main purpose of this work is the introduction of the series (1.2) and its generalizations, and of the techniques leading to Theorem 1.4. The applications to subconvexity and mean values of L-series are included as a "proof of concept" to demonstrate the advantages of this approach.

1.1. **A subconvexity bound.**

The subconvexity problem for $GL(2)$ can be posed in several different aspects and here we will focus on the conductor of a twisting character in the context of holomorphic cusp forms. Specifically, let f be a primitive holomorphic cusp form of even weight k and character χ_1 for $\Gamma_0(N_0)$, having Fourier expansion

$$f(z) = \sum_{m\geq 1} A(m) m^{(k-1)/2} e^{2\pi i m z}.$$

For χ a Dirichlet character mod Q, we write the twisted L-series as

$$L(s, f, \chi) = \sum_{m\geq 1} \frac{A(m)\chi(m)}{m^s}.$$

This possesses a functional equation as $s \to 1-s$ and the convexity bound for the central value which follows is

$$L(1/2, f, \chi) \ll Q^{1/2+\epsilon}.$$

The Lindelöf Hypothesis in Q aspect states that the exponent can be reduced from $1/2+\epsilon$ to ϵ, and any bound that reduces $1/2$ to $1/2-\delta$, for some $\delta > 0$ is called a convexity breaking bound. The corresponding convexity bound for $GL(1)$ is $1/4$.

In a groundbreaking series of papers, Duke, Friedlander, and Iwaniec developed techniques that broke the convexity bound for various families of L-functions [8 , 9 , 10]. Another celebrated result, in the context of $GL(1)$ and a Dirichlet L-series, was obtained by Conrey and Iwaniec, [6], who showed that

$$L(1/2, \chi) \ll Q^{1/6+\epsilon},$$

where the the character χ is real with conductor Q. This is analogous to the Weyl bound for the Riemann zeta function, and it remains the best known result for a $GL(1)$ L-series.

Progress has been made in $GL(2)$ by a number of different authors, with the most recent and strongest results in the twisting parameter being obtained by Blomer and Harcos. In [4], they demonstrate that for f a primitive holomorphic or Maass cusp form of archimedean parameterμ, level N and trivial nebentypus

$$\begin{aligned} &L(1/2, f, \chi) \\ &\quad \ll_\epsilon \left(|s|^{1/4}|\mu|^{1/2}N^{1/4}q^{3/8} + |s|^{1/2}|\mu|N^{1/2}(N,q)^{1/4}q^{1/4}\right)(|s||\mu|Nq)^\epsilon \end{aligned}$$

when f is holomorphic, and

$$\begin{aligned} &L(1/2, f, \chi) \\ &\quad \ll_\epsilon \left(|s|^{1/4}(1+|\mu|)^3 N^{1/4}q^{3/8} + |s|^{1/2}(1+|\mu|)^{7/2}N^{1/2}(N,q)^{1/4}q^{1/4}\right) \\ &\qquad\qquad \times (|s|(1+|\mu|)Nq)^\epsilon \end{aligned}$$

in the Maass form case.

In [5], Blomer and Harcos, work over an arbitrary totally real field K. In this context they let π be an irreducible cuspidal representation of $GL(2,K)\backslash GL(2,\mathbb{A}_K)$, with unitary central character, and χ a Hecke character with conductor of norm Q. They also let θ, with $0 \leq \theta \leq 1/2$ be an approximation toward the Ramanujan-Petersson conjecture. (Here $\theta = 7/64$ is the best known bound. See [21] for $\mathbb{Q}$ and [3]for an arbitrary number field. The Ramanujan-Petersson conjecture predicts $\theta = 0$.) They obtain the bound

$$L(1/2, \pi \otimes \chi) \ll_{\pi,\chi_\infty,K,\epsilon} Q^{\frac{1}{2}-\frac{1}{8}(1-2\theta)+\epsilon}.$$

Here, as our point is to demonstrate a proof of concept for a new method, we work in a more restricted setting than in the work of Blower and Harcos, restricting ourselves to $\mathbb{Q}$, and the case of f holomorphic. In this setting we in some sense combine the above theorems, removing the nebentypus restriction on f as was done in [5] and the dependence on the Ramanujan-Petersson conjecture, as was done in [4]. In Section 9 we prove the following

Theorem 1.1. *Let f be a primitive holomorphic cusp form of even weight k, character χ_1 and fixed level N_0, and let χ be a Dirichlet character modulo Q. Then*

$$L(1/2, f, \chi) \ll_{N_0,\epsilon} Q^{3/8+\epsilon}.$$

1.2. A second moment application.

Let f, g be cusp forms of even weight k for $\Gamma_0(N_0)$. As discussed above, the question of the size of $L(s, f, \chi)$ for any fixed f, s and χ, is related to the Lindelöf Hypothesis and progress is slow. When an average is taken over one of the parameters the problem becomes more tractable, but answers remain more elusive than one would like.

As another application of the techniques developed here we study the second moment

$$S_{f,g}(Q) = \frac{1}{\varphi(Q)} \sum_{\chi \bmod Q} L(1/2, f, \chi)\overline{L(1/2, g, \chi)},$$

that is, the average value of the product of the two L-series over the $\varphi(Q)$ charecters modulo Q. If the Lindelöf Hypothesis is true, the values of the individual L-series should be no larger than Q^ϵ, and the average should reflect the fact that the values are close to constant.

We are not aware of previous results on this problem in the case $f \neq g$, but the case $f = g$ has attracted a great deal of attention. In [17], Heath-Brown found an asymptotic formula in the case where f was an Eisenstein series, when Q was subject to certain constraints. In [28], Soundararajan found an asymptotic formula valid without constraints on Q, and just recently,[33], Young has found such a formula with a power saving error term. The case where f is a modular form was treated in [31], by Stefanicki, who found an asymptotic for this case when the factorization of Q obeyed certain constraints and in [11], Gao, Khan, and Ricotta significantly weakened these constraints.

The ultimate goal is to prove a result for general f with a power savings in the exponent, for example

$$S_{f,f}(Q) = c_f^{(1)} \log Q + c_f^{(2)} + \mathcal{O}(Q^{-\delta}),$$

in the case $f = g$ and

$$S_{f,g}(Q) = c_{f,g} + \mathcal{O}(Q^{-\delta}),$$

in the case $f \neq g$, for some $\delta > 0$. Here $c_f^{(1)}, c_f^{(2)}, c_{f,g}$ should be constants depending on f, g, and independent of Q.

This remains out of (our) reach, but interestingly there seems to be a clear reason for this. As we will see, the values of the leading constants arise not only from the diagonal contribution of certain sums, but also from the off-diagonal terms. The contribution from the off-diagonal terms arise in the form of a special value of a multiple Dirichlet series at a point inside what corresponds to a critical strip. This multiple Dirichlet series is a function of the Q parameter, and in order to obtain the desired asymptotic expression it would be necessary to write this value as a main term plus an error term. There is, however, no reason to believe that such an expression should exist. (For example, it is not likely that the special value of the Dirichlet L-series $L(1/2, \chi_Q)$ can be expressed as a main term plus an error term.) However, it is quite reasonable to believe that an average of these special values, as Q varies over an interval, could be written as a main term plus an error term, and the question then becomes “how short can this interval be made?”

If Q is of size X, the optimal result in this setting would be an average over Q in an interval of length X^ϵ around X and this is in fact what we succeed in proving. In particular, we prove the following

Theorem 1.2. *Let*

$$S_{f,g}(Q) = \frac{1}{\varphi(Q)} \sum_{\chi \bmod Q} L(1/2, f, \chi)\overline{L(1/2, g, \chi)}.$$

The smoothed sum

$$S_{f,g}(X, y) = \sum_{Q \geq 1} S_{f,g}(Q) e^{-y^2 (\log(X/Q))^2/(4\pi)}$$

defines an average of $S_{f,g}(Q)$ *over the interval of length* X/y *centered at* X*. For any* y *in the interval*

$$1 \le y \ll X^{1-\epsilon}$$

we have the following estimates for $S_{f,g}(X,y)$*: When* $f = g$

$$S_{f,f}(X,y) = \frac{C_1(f)L(1,f,\vee^2)e^{\pi/y^2}X\log X}{y} + \frac{C_2(f)X}{y} \\ + \mathcal{O}\left(\frac{X^{1/2}}{y}\right) + \mathcal{O}(X^\epsilon) \quad (1.3)$$

When $f \neq g$

$$S_{f,g}(X,y) = \frac{C(f,g)e^{\pi/y^2}}{y} L(1, f\otimes g)X + \mathcal{O}\left(\frac{X^{1/2}}{y}\right) + \mathcal{O}(X^\epsilon).$$

Here $C_1(f), C_2(f), C(f,g)$ *are non-zero constants that can be made explicit.*

An immediate corollary of this is the following

Corollary 1.1. *Given any* $X \gg 1$ *and* $\epsilon > 0$ *there exists a character* χ *of conductor* Q*, with* $|Q - X| \ll X^\epsilon$ *such that*

$$L(1/2, f, \chi)\overline{L(1/2, g, \chi)} \neq 0.$$

That is, the L-series of f *and* g*, twisted by* χ*, are simultaneously nonzero at the center of the critical strip.*

Remark 1.3. There is a power savings in the exponent of size exactly the interval of Q we average over. In this average we sum over *all* Q in the given interval with the smallest interval length being X^ϵ. It seems likely that if there were restrictions on the divisors of Q it would be difficult to get the length of this interval down past $X^{1/2}$.

1.3. **A bit more on shifted convolutions.**

As mentioned above, the main purpose of this paper is to present a new method for approaching problems of this type. In particular, we will introduce a new class of shifted single and multiple Dirichlet series and show how their full meromorphic continuation can be obtained, and how this meromorphic continuation can be applied to problems such as the second moment problem above.

Let ℓ_1, ℓ_2 be two positive integers, and fix $h \ge 1$. Suppose that f, g are modular forms of even weight k and Fourier expansions

$$f(z) = \sum_{m\ge1} a(m)e^{2\pi imz} = \sum_{m\ge1} A(m)m^{(k-1)/2}e^{2\pi imz},$$

$$g(z) = \sum_{m\geq 1} b(m)e^{2\pi imz} = \sum_{m\geq 1} B(m)m^{(k-1)/2}e^{2\pi imz},$$

The fundamental single Dirichlet series that we will investigate is

$$D(s;h,\ell_1,\ell_2) = \sum_{m_2\geq 1, m_1\ell_1=m_2\ell_2+h} \frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}}.$$

The ℓ_1, ℓ_2 parameters are included for use in amplification arguments targeted at upper bounds for individual L-series. See, for example, [2], [4] and [26]. When $\ell_1 = \ell_2 = 1$ this specializes to

$$D(s;h) = \sum_{m\geq 1} \frac{a(m+h)\bar{b}(m)}{m^{s+k-1}}. \tag{1.4}$$

Selberg mentions that the meromorphic continuation of his series can be obtained by combining "the technique of Rankin" with a certain Poincare series:

$$\sum_{\gamma\in\Gamma_\infty\backslash\Gamma} (\mathfrak{Im}(\gamma z))^s e^{2\pi ih\gamma z}.$$

Goldfeld, [12], wrote down the series (1.4), rather than (1.1) and remarked that Selberg had briefly indicated how to obtain its meromorphic continuation. Goldfeld approaches that continuation in a different way, using instead the Poincare series

$$\sum_{\gamma\in\Gamma_\infty\backslash\Gamma} (\mathfrak{Im}\gamma z)^{1/2} I_{s-1/2}(2\pi|h|\mathfrak{Im}\,\gamma z)e^{2\pi ih\mathfrak{Re}\,\gamma z}.$$

The function $I_v(y)$ is the modified Bessel function of the first kind, and grows exponentially. Goldfeld achieves the meromorphic continuation of (1.4) to $\mathfrak{Re}\, s > 1/2$ via the analytic properties of an inner product of his Poincare series with f and g. However the Dirichlet series that comes out of this method is more closely related to (1.1) than to (1.4), and a hypergeometric series needs to be introduced to carry over the information. The presence of this hypergeometric series seriously impacts efforts to continue past $\mathfrak{Re}\, s > 1/2$.

A fundamental difficulty with the series (1.1) seems to be the tying together of the h and the m variables in the denominator. One can see, for example, that if one is interested in estimating sums of coefficients of a given length x, that this sum becomes seriously unbalanced when h greatly exceeds x. It seems to be the case that the only way to "untie" the m and h is to reverse the sign of the h in the exponent of the Poincare series obtaining, for example:

$$P_h(z;s) = \sum_{\gamma\in\Gamma_\infty\backslash\Gamma} (\mathfrak{Im}(\gamma z))^s e^{-2\pi i\, h\, \gamma z}$$

When one does this, the series $D(s;h)$ comes naturally out of a formal computation of a triple product. Unfortunately the series $P_h(z;s)$ grows exponentially in y, making it difficult to turn the formal computation into a genuine one. Goldfeld faced a similar difficulty in [12] and dealt with it by truncating the fundamental domain. In the present case the difficulties seem to be more fundamental and it appears to be necessary to work with an approximation to $P_h(z;s)$ by two extra parameters. This is done in Sections 2-5. It is shown there that the general series $D(s;h,\ell_1,\ell_2)$, which converges for $\mathfrak{Re}\, s > 1$, has a meromorphic continuation to all of $\mathbb{C}$, with poles at the points $1/2 - r \pm it_j$. Here the eigenvalues of the Laplacian are $1/2 \pm it_j$ and $r \geq 0$ runs over non-negative integers. To accomplish this a spectral expansion for $D(s;h,\ell_1,\ell_2)$ is found which is absolutely convergent for $\mathfrak{Re}\, s < 1/2 - k/2$. The gap between 1 and $1/2 - k/2$ is bridged by an approximation to $D(s;h)$ which is absolutely convergent everywhere, and whose limit approaches $D(s;h)$ as the two parameters tend to infinity and zero respectively. The consequent meromorphic continuation of $D(s;h)$ is the content of Proposition 5.1.

Most applications of shifted sums call for a sum over the shift h. Because of this it is natural to consider the collective behavior of $D(s;h,\ell_1,\ell_2)$ as h varies. This is accomplished first in the meromorphic continuation to $\mathbb{C}^2$ of the shifted double Dirichlet series:

$$Z(s,w) = \sum_{h\geq 1} \frac{D(s;h)}{h^{w+(k-1)/2}} = \sum_{\substack{h\geq 1, m_2\geq 1\\ m_1\ell_1 = m_2\ell_2 + h}} \frac{a(m_1)\bar{b}(m_2)}{(m_2)^{s+k-1} h^{w+(k-1)/2}},$$

and more generally of the series

$$Z_Q(s,w) = \sum_{\substack{h, m_2\geq 1\\ m_1\ell_1 = m_2\ell_2 + hQ}} \frac{a(m_1)\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1} (hQ)^{w+(k-1)/2}}.$$

This series has poles in s at the expected locations: $s = 1/2 - r + it_j$, for $r \geq 0$, that is, the same locations as $D(s;h)$, but its only additional polar line is at $s + w + k/2 = 3/2$. In Section 8 we show the following theorem, which is the underpinning of the applications.

Theorem 1.4. *Let*

$$S_Q(x,y) = \sum_{\substack{\ell_1 m_1 = \ell_2 m2 + hQ\\ h, m_2\geq 1}} A(m_1)\bar{B}(m_2) G_1(m_1/y) G_2(m_2/x),$$

where $G_1(x), G_2(x)$ *are smooth, with compact support in he interval* $[1,2]$. *Suppose* $y \gg x \gg 1$, $\ell_1, \ell_2 \sim L \ll Q^{1/2}$. *Then*

$$S_Q(x,y) \ll Q^{\theta - 1/2} L^{1+\epsilon} y (y/x)^{\epsilon}.$$

If, in addition, $L \gg Q^{1/4}\log(Q)$ *and* $(\ell_1\ell_2, Q) = 1$ *then*

$$S_Q(x,y) \ll Q^{-1/2}L^{1+\epsilon}y(y/x)^{\epsilon}.$$

The coefficients $A(m), B(m)$ are constant on average and the length of this sum is xy/Q. Thus this estimate is non trivial as long as

$$Q^{\theta-1/2}L^{1+\epsilon}y(y/x)^{\epsilon} \ll xy/Q,$$

i.e as long as

$$y \ll \left(\frac{x^{1+\epsilon}}{L^{1+\epsilon}Q^{\theta+1/2}}\right)^{1/\epsilon}.$$

This, in turn, is non-trivial when

$$x \gg LQ^{(\theta+1/2)/(1+\epsilon)}.$$

More generally, we investigate

$$Z_Q(s,w;\nu) = \sum_{\substack{h\equiv\nu \pmod Q \\ m_2\geq 1, m_1\ell_1=m_2\ell_2+h}} \frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}h^{w+(k-1)/2}}.$$

The case of this series when $\nu = 0$ and $\ell_1 = \ell_2 = 1$ is particularly relevant to the second moment application described above. In fact, the special value $Z_Q(1-k/2, 1/2; 0)$ appears in the main term of the estimate for the second moment. It becomes a natural question then to estimate the average of $Z_Q(s,w;0)$ as Q varies, and for this purpose we introduce the triple Dirichlet series

$$\mathcal{M}(s,w,v) = \sum_{Q\geq 1}\frac{Z_Q(s,w;0)}{Q^v} = \sum \frac{a(m+h)\bar{b}(m)\sigma_{-v}(h)}{m^{s+k-1}h^{w+(k-1)/2}}.$$

Here σ is the usual divisor function. In this case we obtain a meromorphic continuation to all $(s,w,v) \in \mathbb{C}^3$ with $\mathfrak{Re}\, v \geq 0$. As it may be of interest we list some of the polar divisors here:

- $s = 1/2 \pm it_j - r$ for $r \geq 0$,
- $w + v/2 = 1/2 \pm it_j - r$ for $r \geq 0$,
- $w + v/2 = -r$, for $r \geq 0$,
- $s + w + k/2 + v/2 - 2 = 0$

This can be translated into (for example) asymptotic formulas for double sums of coefficient products $a(m+h)\bar{b}(m)\sigma_{-v}(h)$ that display substantial cancellation in the shift and length parameters. The full description of the continuation is given in Proposition 11.4.

One interesting aspect of the one and two variable multiple Dirichlet series discussed is that each has two incarnations: a Dirichlet series that converges absolutely in one range, and a spectral expansion that converges absolutely in a disjoint region. As one might expect, all the interesting information seems to be hidden in between. The spectral expansions have another curious aspect. The terms decay polynomially in the spectral parameter t_j, rather than exponentially, and the ratio of gamma factors governing this decay seems to act in a manner analogous to the usual $m^{-\sigma}$ indexing of a Dirichlet series (polynomial decay corresponding to the size of σ.)

The author would like to thank V. Blomer, D. Hansen, A. Kontorovich, A. Reznikov and M. Young for many very helpful conversations.

2. Spectral expansions

Fix $N_0, \ell_1, \ell_2, \ell, h \in \mathbb{N}$, and set $N = N_0\ell_1\ell_2$ and $\Gamma = \Gamma_0(N)$. For $z \in \mathbb{H}$ and $s \in \mathbb{C}$, $\mathfrak{Re}(s) > 1$, consider the Poincare series:

$$P_h(z;s) := \sum_{\gamma\in\Gamma_\infty\backslash\Gamma} (\mathfrak{Im}(\gamma z))^s e^{-2\pi i\, h\, \gamma z}. \tag{2.1}$$

This is the usual Poincare series except that the sign in the exponent is negative rather than positive. For any fixed z the series converges absolutely for $\mathfrak{Re}(s) > 1$ and is invariant under action by Γ. It does, however, grow exponentially in y, which complicates efforts to determine its spectral properties. For this reason, for any fixed $Y \gg 1$ and $\delta > 0$ we define the modified series $P_{h,Y}(z, s; \delta)$ by

$$P_{h,Y}(z,s;\delta) := \sum_{\gamma\in\Gamma_\infty\backslash\Gamma} \psi_Y(\mathfrak{Im}\gamma z)(\mathfrak{Im}(\gamma z))^s e^{-2\pi i\, h\, \mathfrak{Re}\gamma z + (2\pi\, h\, \mathfrak{Im}\gamma z)(1-\delta)}, \tag{2.2}$$

where ψ_Y is the characteristic function of the interval $[Y^{-1}, Y]$.

The series $P_{h,Y}(z, s; \delta)$ consists of finitely many terms and is in $L^2(\Gamma\backslash\mathbb{H})$. It also possesses a Fourier expansion, which can be easily worked out:

$$P_{h,Y}(z,s;\delta) = \sum_m a_{m,Y}(s,y;\delta)e^{2\pi i m x}, \tag{2.3}$$

where

$$\begin{aligned} a_{m,Y}(s,y;\delta) &= \delta_{m,-h} y^s e^{2\pi h y(1-\delta)}\psi_Y(y) \\ &+ \sum_{c\neq 0} \frac{S(m,-h;cN)}{N^{2s}|c|^{2s}} \cdot y^{1-s} \\ &\times \int_{-\infty}^{\infty} \frac{e^{(2\pi i h u + 2\pi h(1-\delta))/(N^2c^2y(u^2+1))}}{(u^2+1)^s} \\ &\times e^{-2\pi i m u y}\psi_Y(\frac{1}{N^2c^2y(u^2+1)})du. \end{aligned} \tag{2.4}$$

The Petersson inner product of two forms F, G in $L^2(\Gamma\backslash\mathbb{H})$ is defined by

$$\langle F, G\rangle = \frac{1}{\mathcal{V}}\iint_{\Gamma\backslash\mathbb{H}} F(z)\overline{G(z)}\frac{dxdy}{y^2}, \tag{2.5}$$

where $\mathcal{V} = \text{volume}(\Gamma\backslash\mathbb{H}) \asymp N$. Set V to be the Γ-invariant function

$$V_{f,g}(z;\ell_1,\ell_2) := y^k\overline{f(\ell_1 z)}g(\ell_2 z), \tag{2.6}$$

which is rapidly decreasing in the cusps of Γ.

Our approach will be to compute the inner product $\langle P_{h,Y}(*, s;\delta), V\rangle$ in several different ways, and to then let $Y \to \infty$. To begin, we compute

$$\begin{aligned}
\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) &:= \langle P_{h,Y}(*, s;\delta), V\rangle \qquad (2.7)\\
&= \frac{1}{\mathcal{V}}\iint_{\Gamma\backslash\mathbb{H}} P_{h,Y}(z,s;\delta)f(\ell_1 z)\overline{g(\ell_2 z)}y^k\frac{dx\,dy}{y^2}\\
&= \frac{1}{\mathcal{V}}\sum_{m_1\geq 1}\sum_{m_2\geq 1} a(m_1)\bar{b}(m_2)\int_0^1 e^{2\pi i x(m_1\ell_1 - m_2\ell_2 - h)}dx\\
&\quad\times\int_0^\infty e^{-2\pi y(m_1\ell_1 + m_2\ell_2 - h(1-\delta))}y^{s+k-1}\psi_Y(y)\frac{dy}{y}\\
&= \frac{1}{\mathcal{V}}(4\pi)^{-(s+k-1)}\sum_{m_1\ell_1 = m_2\ell_2 + h}\frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}}\\
&\quad\times\int_{Y^{-1}4\pi\ell_2 m_2}^{Y4\pi\ell_2 m_2} e^{-y(1+h\delta/(2\ell_2 m_2))}y^{s+k-1}\frac{dy}{y}.
\end{aligned}$$

Note that for $\mathfrak{Re}s > 1$, and all $Y > 1, \delta > 0$ the integral and sum converge absolutely and satisfy the upper bound

$$\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) \ll \frac{1}{N}\Gamma(\mathfrak{Re}s + k - 1)\sum_{m_1\ell_1 = m_2\ell_2 + h}\frac{|a(m_1)\bar{b}(m_2)|}{(m_2\ell_2)^{\mathfrak{Re}s+k-1}},$$

where the implied constant is absolute.

Let $\{u_j\}_{j\geq 0}$ be an orthonormal basis (with respect to the inner product defined in (2.5)) for the discrete part of the spectrum of the Laplace operator on $L^2(\Gamma\backslash\mathbb{H})$. Suppose the u_j have eigenvalue $1/4 + t_j^2$, and Fourier expansion

$$u_j(z) = \sum_n \rho_j(n)y^{1/2}K_{it_j}(2\pi|n|y)e^{2\pi i n x}.$$

As $P_{h,Y}(z,s;\delta) \in L^2(\Gamma\backslash\mathbb{H})$ we can obtain the spectral expansion of $\mathcal{I}$:

$$\begin{aligned}\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) &= \sum_{j\geq 1} \overline{\langle V, u_j\rangle}\, \langle P_{h,Y}(*,s;\delta), u_j\rangle \\ &+ \frac{1}{4\pi}\sum_{a}\int_{-\infty}^{\infty} \overline{\langle V, E_a(*,1/2+it)\rangle}\, \langle P_{h,Y}(*,s;\delta), E_a(*,1/2+it)\rangle\, dt.\end{aligned} \tag{2.8}$$

We easily compute

$$\langle P_{h,Y}(*,s;\delta), u_j\rangle = \frac{\overline{\rho_j(-h)}}{\mathcal{V}(2\pi h)^{s-1/2}} \int_{Y^{-1}2\pi h}^{Y2\pi h} y^{s-1/2} e^{y(1-\delta)} K_{it_j}(y)\frac{dy}{y} \tag{2.9}$$

and

$$\begin{aligned}&\langle P_{h,Y}(*,s;\delta), E_a(*,1/2+it)\rangle \\ &\quad = \frac{\overline{\tilde{\rho}_a(1/2+it,-h_a)} 2\pi^{1-s-it} 2^{3/2-s} h^{1/2-s-it}\sigma_{2it}(h)}{\mathcal{V}\Gamma(1/2-it)\zeta(1-2it)} \\ &\qquad\qquad \times \int_{Y^{-1}2\pi h}^{Y2\pi h} y^{s-1/2} e^{y(1-\delta)} K_{it}(y)\frac{dy}{y}.\end{aligned} \tag{2.10}$$

Note that u_0 is constant and $\langle u_0, P_h\rangle = 0$. We have taken the Eisenstein series notation from [2], that is: $E_a(z,s)$ denotes the Eisenstein series expanded at the cusp a. Here for each cusp $a = u/w$,

$$\begin{aligned}E_a(z,s) &= \delta_{a,\infty} y^s + \frac{\sqrt{\pi}\Gamma(s-1/2)\rho_a(s,0)y^{1-s}}{\Gamma(s)} \\ &+ \frac{2\pi^s\sqrt{y}}{\Gamma(s)} \sum_{m\neq 0} |m|^{s-1/2}\rho_a(s,m) K_{s-1/2}(2\pi|m|y) e^{2\pi i m x},\end{aligned} \tag{2.11}$$

and

$$\rho_a(s,m) = \left(\frac{(w,\frac{N}{w})}{wN}\right)^s \sum_{(\gamma,N/w)=1} \frac{1}{\gamma^{2s}} \sum_{\substack{\delta \bmod \gamma w \\ (\delta,\gamma w)=1 \\ \delta\gamma\equiv u \bmod (w,N/w)}} e^{-2\pi i m\delta/(\gamma w)}. \tag{2.12}$$

The part of m that is relatively prime to N_0 factors out of $\rho_a(s,m)$ and is an ordinary divisor function.

In fact, we find it convenient to write $\rho_a(s,m)$ a bit more explicitly as

$$\rho_a(s,0) = \tilde{\rho}_a(s,0)\frac{\zeta(2s-1)}{\zeta(2s)}$$

and for $m \neq 0$

$$\rho_a(s,m) = \frac{\tilde{\rho}_a(s,m_a)\sigma_{1-2s}(m)}{\zeta(2s)},$$

where m_a is composed of powers of primes dividing N and $(m, N) = 1$. In particular, in the case of $SL(2,\mathbb{Z})$, where there is only one cusp at $a = \infty$,

$$\tilde{\rho}_\infty(s, 1) = 1.$$

The $\tilde{\rho}_a(s, m_a)$ satisfies

$$\sum_a |\tilde{\rho}_a(1/2 + it, m_a)|^2 \ll_\epsilon ((1 + |t|)N)^\epsilon. \tag{2.13}$$

This is translated from [2] Lemma 3.4.

We will ultimately need to understand the behavior of the integral appearing in (2.9) and (2.10) for fixed small δ as $Y \to \infty$. With this in mind we define

$$M(s, t, \delta) = \int_0^\infty y^{s-1/2} e^{y(1-\delta)} K_{it}(y) \frac{dy}{y}. \tag{2.14}$$

Using the well known bounds

$$K_{\gamma+it}(y) \ll \frac{e^{-y}}{\sqrt{y}} \tag{2.15}$$

for $y \gg 1$ and

$$K_{\gamma+it}(y) \ll y^{-|\gamma|} \tag{2.16}$$

for $y \ll 1$, it is clear that the integral expression for $M(s, t, \delta)$, given in (2.14), converges absolutely for any $\delta > 0$ when $\mathfrak{Re}\, s > 1/2$. (It also converges absolutely for $\delta = 0$, when $\frac{1}{2} < \mathfrak{Re} s < 1$, a fact that will be made use of later.) The following proposition makes explicit the extent to which the integral on the right hand side of (2.9) approximates $M(s, t, \delta)$ when $\mathfrak{Re}\, s > 1/2$.

Proposition 2.1. *Let $M(s, t, \delta)$ be as defined in* (2.14). *Fix $\epsilon > 0, Y \gg 1, A \gg 1, 1 > \delta > 0$ and $s \in \mathbb{C}$ with $\mathfrak{Re}\, s > 1/2 + \epsilon$. Then*

$$\left| M(s, t, \delta) - \int_{Y^{-1}2\pi h}^{Y2\pi h} y^{s-1/2} e^{y(1-\delta)} K_{it}(y) \frac{dy}{y} \right| \ll \frac{e^{-Y2\pi h\delta}(Yh)^{\mathfrak{Re}\, s+A-2}}{\delta(1 + |t|)^A} + \frac{(Y^{-1}h)^{\mathfrak{Re}\, s-1/2}}{(1 + |t|)^A},$$

with the implied constant depending only on A and ϵ.

Proof. We first need to bound $K_{it}(y)$ uniformly in t, for y both large and small. For this we have

Lemma 2.2. *For $A, |t|, y \gg 1$*

$$K_{it}(y) \ll_A \frac{e^{-y} y^A}{\sqrt{y}(1 + |t|)^A}.$$

For $A, |t| \gg 1$ *and* $0 < y < 1$

$$K_{it}(y) \ll_A \frac{1}{(1+|t|)^A}.$$

To prove Lemma 2.2 recall the definition

$$K_{it}(y) = \frac{1}{2}\int_0^\infty e^{-\frac{y}{2}(u+u^{-1})} u^{it} \frac{du}{u}.$$

The result follows after integrating by parts $[A]+1$ times and applying the bound (2.15) when y is large, (2.16) when y is small to $K_{r+it}(y)$, with $r = 0, 1, \dots, [A]$.

Applying Lemma 2.2 we find that

$$\begin{aligned}\int_{Y2\pi h}^\infty & y^{s-1/2} e^{y(1-\delta)} K_{it}(y) \frac{dy}{y} \ll \int_{Y2\pi h}^\infty y^{\mathfrak{Re}\, s-1/2} \frac{e^{y(1-\delta)} e^{-y} y^{A-1/2}}{(1+|t|)^A} \frac{dy}{y} \\ \ll & \frac{1}{(1+|t|)^A} \int_{Y2\pi h}^\infty y^{\mathfrak{Re}\, s+A-1} e^{-y\delta} \frac{dy}{y} \\ = & \frac{1}{(1+|t|)^A \delta^{\mathfrak{Re}\, s+A-1}} \Gamma[\mathfrak{Re}\, s + A - 1, Y2\pi h\delta],\end{aligned}$$

where

$$\Gamma[s,x] := \int_x^\infty e^{-y} y^s \frac{dy}{y} \ll e^{-x} x^{s-1},$$

as $x \to \infty$. Thus

$$\int_{Y2\pi h}^\infty y^{s-1/2} e^{y(1-\delta)} K_{it}(y) \frac{dy}{y} \ll \frac{e^{-Y2\pi h\delta}(Y2\pi h\delta)^{\mathfrak{Re}\, s+A-2}}{(1+|t|)^A \delta^{\mathfrak{Re} s+A-1}} \ll \frac{e^{-Y2\pi h\delta}(Yh)^{\mathfrak{Re}\, s+A-2}}{\delta(1+|t|)^A}.$$

Similarly, applying the second part of the lemma,

$$\int_0^{Y^{-1}2\pi h} y^{s-1/2} e^{y(1-\delta)} K_{it}(y) \frac{dy}{y} \ll \frac{(Y^{-1}2\pi h)^{\mathfrak{Re}\, s-1/2}}{(1+|t|)^A},$$

completing the proof of Proposition 2.1.

□

3. THE ANALYTIC CONTINUATION OF $M(s,t,\delta)$

In this section we will provide a meromorphic continuation of $M(s,t,\delta)$ to $s \in \mathbb{C}$ with $\mathfrak{Re}\, s \leq 1/2$. The following proposition locates the poles of $M(s,t,\delta)$ and gives three different estimates at points s where $M(s,t,\delta)$ is analytic.

Proposition 3.1. *Fix* $\epsilon > 0, \delta > 0$ *and* $A \gg 1$. *The function* $M(s,t,\delta)$ *has a meromorphic continuation to all* s *in the half plane* $\mathfrak{Re}\, s > 1/2 - A$. *In this region* $M(s,t,\delta)$ *is analytic except for simple poles at the* *points* $s = 1/2 \pm it - \ell$ *for* $0 \leq \ell < A$. *The residues at* $s = 1/2 + it - \ell$ *are given by*

$$\begin{aligned} \mathrm{Res}_{s=1/2\pm it-\ell}\, M(s,t,\delta) &= \frac{(-1)^{\ell}\sqrt{\pi}2^{\ell\mp it}\Gamma(1/2\mp it+\ell)\Gamma(\pm 2it-\ell)}{\ell!\Gamma(1/2+it)\Gamma(1/2-it)} \\ &+ \mathcal{O}((1+|t|)^{\ell}e^{-\frac{\pi}{2}|t|}\delta^{1/2}), \end{aligned} \tag{3.1}$$

where the implied constants are independent of t.

For s *at least a distance* ϵ *away from these points,* $M(s,t,\delta)$ *satisfies the upper bound.*

$$M(s,t,\delta) \ll (1+|t|)^{2\mathfrak{Re}\, s-2-2A}(1+|s|)^{A-\mathfrak{Re}\, s}\delta^{-A}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}. \tag{3.2}$$

Also, for $\mathfrak{Re}\, s \leq 1/2$, *at least a distance* ϵ *away from these points, and* $\delta(1+|t|)^2 \leq 1$

$$\begin{aligned} M(s,t,\delta) &= \frac{\sqrt{\pi}2^{1/2-s}\Gamma(s-1/2-it)\Gamma(s-1/2+it)\Gamma(1-s)}{\Gamma(1/2-it)\Gamma(1/2+it)} \\ &+ \mathcal{O}\left((1+|t|)^{2\mathfrak{Re}\, s-2+2\epsilon}(1+|s|)^{1/2-\mathfrak{Re}\, s}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\delta^{\epsilon}\right) \end{aligned} \tag{3.3}$$

while for $\delta(1+|t|)^2 > 1$

$$M(s,t,\delta) \ll (1+|t|)^{2\mathfrak{Re}\, s-2}(1+|s|)^{A+\mathfrak{Re}\, s-1}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}. \tag{3.4}$$

Proof. From [14] formula 6.621 # 3 we obtain, for $\mathfrak{Re} s > 1/2$,

$$\begin{aligned} M(s,t,\delta) &= \int_0^{\infty} y^{s-1/2}e^{y(1-\delta)}K_{it}(y)\frac{dy}{y} \\ &= \frac{\sqrt{\pi}2^{it}\Gamma(s-1/2+it)\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(s)} \times F(s-1/2+it, 1/2+it; s; 1-2/\delta), \end{aligned} \tag{3.5}$$

where $F(\alpha,\beta;\gamma;x)$ is the usual hypergeometric function. By [14] formula 9.113, this can be written as a contour integral

$$\begin{aligned} F(s-1/2+it, 1/2 &+ it; s; 1-2/\delta) = \frac{\Gamma(s)}{\Gamma(s-1/2+it)\Gamma(1/2+it)} \\ &\times \frac{1}{2\pi i}\int_C \frac{\Gamma(s-1/2+u+it)\Gamma(1/2+u+it)\Gamma(-u)\left(\frac{2}{\delta}-1\right)^u du}{\Gamma(s+u)} \end{aligned} \tag{3.6}$$

where C goes from $-i\infty$ to $i\infty$ and curves so that the poles of $\Gamma(-u)$ lie to the right of C and the poles of $\Gamma(s-1/2+u+it), \Gamma(1/2+u+it)$ lie to the left. For $\mathfrak{Re}\, s > 1/2$, C can be simply the vertical line $\mathfrak{Re}\, u = a$, where a is chosen so that $1/2 - \mathfrak{Re}\, s < a < 0$.

Remark 3.2. It is clear from (3.6) that $F(s-1/2+it, 1/2+it; s; 1-2/\delta)$ does *not* have a pole at $s = 1$, and so, by (3.5), neither does $M(s,t,\delta)$. This is in spite of the appearance of $\Gamma(1-s)$ in (3.3), as that expression does not converge as $\delta \to 0$ for $\mathfrak{Re}\, s > 1/2$.



Figure 1. The contour

The expression (3.6) is valid for any fixed $s \in \mathbb{C}$ and thus when combined with (3.5) provides a continuation of $M(s, t, \delta)$. For general s the curve C can be described as follows. Write $\mathfrak{Re}\, s = 1/2 - R$, with $R \geq 0$. Choose b such that $0 > b > \max(R - [R] - 1, -1/2)$. Then for $\mathfrak{Im}\, u \leq 0$, C is the line $\mathfrak{Re}\, u = b$. The poles of $\Gamma(1/2 + u + it)$ all lie to the left of $\mathfrak{Re}\, u = b$. The poles of $\Gamma(-u)$ all lie to the right of $\mathfrak{Re}\, u = b$ and the poles of $\Gamma(s - 1/2 + u + it)$ to the right of $\mathfrak{Re}\, u = b$ are at $\mathfrak{Re}\, u = 1/2 - \mathfrak{Re}\, s - \ell = R - \ell$ for $0 \leq \ell \leq [R]$. The curve C moves in a clockwise semicircle above the real line, past 0 and down to the real axis, then counter clockwise in a semicircle around $R - [R]$, then above 1 and so on until it does a final counterclockwise turn around R and continues to $i\infty$ along the line $\mathfrak{Re}\, u = c$, where $R < c < [R] + 1$. (See Figure 1.)

Fix s, with $\mathfrak{Re}\, s = 1/2 - R > 1/2 - A$, and move C to the right to the vertical line $\mathfrak{Re} u = A$. In the process, poles are passed over at

$u = \ell$, for $0 \le \ell < A$. The residue after passing the pole at $u = \ell$ is given by

$$\frac{\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(1/2+it)}R(s,t,\ell), \tag{3.7}$$

where

$$R(s,t,\ell) = \frac{\Gamma(s-1/2+it+\ell)\Gamma(1/2+it+\ell)\left(\frac{2}{\delta}-1\right)^{\ell}(-1)^{\ell+1}}{\ell!\Gamma(s+\ell)} \tag{3.8}$$

Thus

$$M(s,t,\delta) = \frac{\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(1/2+it)}N_1(s,t,\delta),$$

where

$$\begin{aligned} N_1(s,t,\delta) &= -\sum_{0\le\ell<A} R(s,t,\ell) \qquad (3.9)\\ &+ \frac{1}{2\pi i}\int_{A-i\infty}^{A+i\infty} \frac{\Gamma(s-1/2+u+it)\Gamma(1/2+u+it)\Gamma(-u)\left(\frac{2}{\delta}-1\right)^{u}du}{\Gamma(s+u)}. \end{aligned}$$

The integral in (3.9) contributes no poles in s, as $\mathfrak{Re}\, s - 1/2 + u > 1/2 - A - 1/2 + A = 0$. Inspecting lines (3.7) to (3.9) it is then clear that the only possible locations of poles of $M(s,t,\delta)$ for $\mathfrak{Re}\, s > 1/2 - A$ are at $s = 1/2 \pm it - m$, for $0 \le m < A$.

Assume now that $R \ge 0$ and move C further to the right, to the vertical line $\mathfrak{Re}\, u = A + R$. In the process, additional poles are passed over at $u = \ell$, for $A \le \ell < A + R$. Changing variables, $u \to u - (s - 1/2 + it)$, and substituting back into (3.5), we now have, after moving the contour and subtracting the residue contribution,

$$M(s,t,\delta) = \frac{\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(1/2+it)}N_2(s,t,\delta),$$

where

$$\begin{aligned} N_2(s,t,\delta) &= -\sum_{0\le\ell<A+R} R(s,t,\ell) \qquad (3.10)\\ &+ \frac{1}{2\pi i}\int_{A-i\infty}^{A+i\infty} \frac{\Gamma(u)\Gamma(1-s+u)\Gamma(s-1/2+it-u)\left(\frac{2}{\delta}-1\right)^{u-s+1/2-it}du}{\Gamma(1/2-it+u)}. \end{aligned}$$

Recall that by Stirling's formula, for fixed x, a, as $y \to \infty$,

$$|\Gamma(x+iy)| \ll |y|^{x-1/2}e^{-\frac{\pi}{2}|y|} \text{ and } \frac{\Gamma(x+a+iy)}{\Gamma(x+iy)} \ll |x+iy|^{a}. \tag{3.11}$$

Writing $s = \sigma + ir$ and $u = A + iv$ the following upper bounds are easily confirmed, using (3.11):

$$R(s,t,\ell) \ll \delta^{-\ell} e^{-\frac{\pi}{2}(|r+t|+|t|-|r|)} \frac{(1+|r+t|)^{\sigma+\ell-1}(1+|t|)^{\ell}}{(1+|r|)^{\sigma+\ell-1/2}},$$

$$\frac{\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(1/2+it)} \ll \delta^{R}(1+|r-t|)^{\sigma-1} e^{-\frac{\pi}{2}(|r-t|-|t|)}, \tag{3.12}$$

and

$$\frac{\Gamma(u)\Gamma(1-s+u)\Gamma(s-1/2+it-u)\left(\frac{2}{\delta}-1\right)^{u-s+1/2-it}}{\Gamma(1/2-it+u)} \tag{3.13}$$
$$\ll \frac{(1+|v|)^{A-1/2}(1+|v-r|)^{A+1/2-\sigma}(1+|t-v+r|)^{\sigma-1-A}}{\delta^{A+R}(1+|t-v|)^{A} e^{\frac{\pi}{2}(|v|+|v-r|+|r-v+t|-|t-v|)}}.$$

Combining the first two, one has

$$\sum_{0\le \ell < A+R} R(s,t,\ell)\frac{\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(1/2+it)} \tag{3.14}$$
$$\ll (1+|t|+|r|)^{2\sigma+2A-2}\delta^{-A} e^{-\frac{\pi}{2}\max(|t|,|r|)}.$$

Thus this term decays exponentially in $|t|$ and is on the order of δ^{-A}. Combining the second and the third, one sees that there is exponential decay in the v variable, polynomial decay in $|t|$ with exponent $2\sigma-2-2A$, and the term is on the order of δ^{-A}.

By a case analysis of the possible respective sizes and signs of v, r and t, we find that the exponent $|v|+|v-r|+|r-v+t|-|t-v|$ in (3.13), when added to the exponent $|r-t|-|t|$ of (3.12) is always $\ge |v|-|r|$ when $|v|>|r|$ and $\ge |r|$ when $0 \le |v| \le |r|$. The worst growth, i.e the smallest exponent, is when $|v| \approx |r|$. The upper bound (3.2) follows after using this analysis to bound the integral in (3.10) from above, and combining this with (3.12), (3.13) and (3.14).

To prove (3.3), (3.4) we return to the original contour integral in line (3.6). Recall that for $\mathfrak{Re}\, s \le 1/2$ we write $\mathfrak{Re}\, s = 1/2 - R$, with $R \ge 0$. Instead of deforming the contour by moving it to the right, we instead straighten the contour until it becomes the line $\mathfrak{Re}\, u = b$. In the process, poles are passed over at the points $u = 1/2 - s - \ell - it$, for $0 \le \ell \le [R]$. (If $\mathfrak{Re}\, s > 1/2$ no poles are passed over.) The residue of the integrand at $u = 1/2 - s - \ell - it$ is

$$\frac{(-1)^{\ell}\Gamma(1-s-\ell)\Gamma(-1/2+s+it+\ell)(2/\delta-1)^{1/2-s-it-\ell}}{\ell!\Gamma(1/2-it-\ell)}.$$

Thus

$$M(s,t,\delta) = C + D, \tag{3.15}$$

where

$$\begin{aligned} C &= \sum_{\ell=0}^{[R]} \frac{(-1)^{\ell}\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\ell!\Gamma(1/2+it)\Gamma(1/2-it-\ell)} \\ &\times \Gamma(1-s-\ell)\Gamma(-1/2+s+it+\ell)(2-\delta)^{1/2-s-it}(2/\delta-1)^{-\ell} \end{aligned} \tag{3.16}$$

and

$$\begin{aligned} D &= \frac{\sqrt{\pi}2^{it}\Gamma(s-1/2-it)}{\delta^{s-1/2+it}\Gamma(1/2+it)} \\ &\times \frac{1}{2\pi i}\int_{b-i\infty}^{b+i\infty} \frac{\Gamma(s-1/2+it+u)\Gamma(1/2+it+u)\Gamma(-u)\left(\frac{2}{\delta}-1\right)^{u} du}{\Gamma(s+u)}. \end{aligned} \tag{3.17}$$

It is now a simple matter to compute

$$\begin{aligned} \operatorname{Res}_{s=1/2-m+it} M(s,t,\delta) &= \frac{\sqrt{\pi}2^{it}(-1)^{m}}{m!\Gamma(1/2+it)} \\ \times \Bigg[\sum_{0\le\ell\le m} &\frac{\Gamma(1/2+m-it-\ell)\Gamma(-m+2it+\ell)(2-\delta)^{m-2it}\left(\frac{2}{\delta}-1\right)^{-\ell}(-1)^{\ell}}{\ell!\Gamma(1/2-it-\ell)} \\ + \frac{1}{2\pi i}\int_{b-i\infty}^{b+i\infty} &\frac{\Gamma(-m+2it+u)\Gamma(1/2+it+u)\Gamma(-u)\delta^{m-2it}\left(\frac{2}{\delta}-1\right)^{u} du}{\Gamma(1/2-m+it+u)}\Bigg] \end{aligned} \tag{3.18}$$

and

$$\begin{aligned} \operatorname{Res}_{s=1/2-m-it} M(s,t,\delta) &= \frac{\sqrt{\pi}2^{it}(-1)^{m}(2-\delta)^{m}\Gamma(-m-2it)}{\Gamma(1/2+it)} \\ \times \sum_{0\le\ell\le m} &\frac{\Gamma(1/2+m+it-\ell)\left(\frac{2}{\delta}-1\right)^{-\ell}}{\ell!(m-\ell)!\Gamma(1/2-it-\ell)}. \end{aligned} \tag{3.19}$$

A careful application of Stirling's formula, (3.11), shows that the main contribution to the residue is given by the $\ell = 0$ term and that

$$\begin{aligned} \operatorname{Res}_{s=1/2-m\pm it} M(s,t,\delta) &= \frac{\sqrt{\pi}2^{m\mp it}(-1)^{m}\Gamma(1/2+m\mp it)\Gamma(-m\pm 2it)}{m!\Gamma(1/2+it)\Gamma(1/2-it)} \\ &+ \mathcal{O}\left((1+|t|)^{m-b-1/2}e^{-\frac{\pi}{2}|t|}\delta^{-b}\right) \end{aligned} \tag{3.20}$$

Another application of Stirling's formula to (3.15), (3.16), (3.17) yields

$$\begin{aligned} M(s,t,\delta) &= \frac{\sqrt{\pi}2^{1/2-s}\Gamma(s-1/2+it)\Gamma(s-1/2-it)\Gamma(1-s)}{\Gamma(1/2+it)\Gamma(1/2-it)} \\ &+ \mathcal{O}\left((1+|t|)^{2\sigma-2}(1+|s|)^{1/2-\sigma}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\max_{\ell=1,[1/2-\sigma]}\left(\delta^{\ell}(1+|t|)^{2\ell}\right)\right) \\ &+ \mathcal{O}\left((1+|t|)^{-1-2b}(1+|s|)^{1/2-\sigma}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\delta^{1/2-\sigma-b}\right), \end{aligned} \tag{3.21}$$

where the main term comes from the $\ell = 0$ part of C.

Consider now the two cases: $\delta(1+|t|)^2 \leq 1$ and $\delta(1+|t|)^2 > 1$. Choose any $1 > \epsilon > 0$. In the first case

$$\max_{\ell=1,[1/2-\sigma]}\left(\delta^{\ell}(1+|t|)^{2\ell}\right) = \delta(1+|t|)^2 \leq \delta^{\epsilon}(1+|t|)^{2\epsilon}.$$

The first error term of (3.21) is then

$$\ll (1+|t|)^{2\sigma-2+2\epsilon}(1+|s|)^{1/2-\sigma}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\delta^{\epsilon}$$

and the second is

$$\ll (1+|t|)^{-1}(1+|s|)^{\sigma-3/2}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\delta^{1/2-\sigma+\epsilon-\epsilon}$$

which is

$$\ll (1+|t|)^{2\sigma-2+2\epsilon}(1+|s|)^{\sigma-3/2}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\delta^{\epsilon}.$$

Thus both pieces of the error term are

$$\mathcal{O}\left((1+|t|)^{2\sigma-2+2\epsilon}(1+|s|)^{1/2-\sigma}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}\delta^{\epsilon}\right),$$

giving us (3.3).

In the second case, $\delta(1+|t|)^2 > 1$, we return to the first estimate, (3.2). Substituting $\delta^{-A} < (1+|t|)^{2A}$ we obtain

$$M(s,t,\delta) \ll (1+|t|)^{2\sigma-2}(1+|s|)^{A+\sigma-1}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|},$$

giving us (3.4), completing the proof of Proposition 3.1. □

4. THE LIMIT AS $Y \to \infty$

We are now in a position to analyze the uniformity of convergence of the spectral decomposition of $\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h)$ given in (2.8). We assume at first that $\mathfrak{Re}\, s > 1/2$. Recall that by (2.8) we may write

$$\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) = \mathcal{I}^{\text{cusp}}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) + \mathcal{I}^{\text{cont}}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h),$$

where

$$\begin{aligned}\mathcal{I}^{\mathrm{cusp}}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) \quad &= \sum_j \frac{\overline{\rho_j(-h)}}{\mathcal{V}(2\pi h)^{s-1/2}} \\ &\times \int_{Y^{-1}2\pi h}^{Y2\pi h} y^{s-1/2} e^{y(1-\delta)} K_{it_j}(y) \frac{dy}{y} \overline{\langle V, u_j\rangle}\end{aligned}$$

and

$$\begin{aligned}\mathcal{I}^{\mathrm{cont}}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) \quad &= \sum_{\mathfrak{a}} \frac{\overline{\tilde{\rho}_{\mathfrak{a}}(1/2+it,-h_{\mathfrak{a}})} 2\pi^{1-s-it} 2^{3/2-s} h^{1/2-s-it} \sigma_{2it}(h)}{\mathcal{V}\Gamma(1/2-it)\zeta(1-2it)} \\ &\times \int_{Y^{-1}2\pi h}^{Y2\pi h} y^{s-1/2} e^{y(1-\delta)} K_{it}(y) \frac{dy}{y} \overline{\langle V, E_{\mathfrak{a}}(*, 1/2+it)\rangle}.\end{aligned}$$

Of key importance is the average size of $\overline{\rho_j(-h)}$ and of $\overline{\langle V, u_j\rangle}$. This is addressed by the following proposition, a generalization due to Reznikov of [1]:

Proposition 4.1. *With the notation as given above*

$$\sum_{|t_j|\sim T} |\langle V, u_j\rangle|^2 e^{\pi |t_j|} \ll_{N_0} (\ell_1\ell_2)^{-k} T^{2k}.$$

Remark 4.2. At this moment A. Reznikov believes that he can prove the content of Proposition 4.1, which is essentially a generalization of [1] from the compact case to the non-compact case, but the details have not been completely written down. As an alternative approach, the needed average of (4.2) should be obtainable by a relatively simple modification of the approach of [20]. V. Blomer was kind enough to make this suggestion. This will be supplied in place of Proposition 4.1 if required.

By Weyl's law, in each unit interval at height T there are on the order of $\ell_1\ell_2 T$ eigenvalues. Thus the proposition is saying that on average, for each j with $|t_j| \sim T$,

$$|\langle V, u_j\rangle|^2 \approx (\ell_1\ell_2)^{-k-1} T^{2k-2} e^{-\pi|t_j|}.$$

The following explicit analysis by Hansen, [15], of some formulas of Ichino, [18], sheds light on the nature of the triple product $\langle V, u_j\rangle$, and in particular on the dependence on ℓ_1, ℓ_2.

Proposition 4.3. *(Hansen) Let f_1, f_2 be a pair of cusp forms of even weight $k > 0$ and level N_0, normalized so their first Fourier coefficient* *equals one. Let ℓ_1, ℓ_2 be a pair of odd primes, and let φ be a Hecke-Maass cuspidal new form of level $N_0\ell_1\ell_2$. Then*

$$\left|\frac{1}{\ell_1\ell_2}\int_{\Gamma_0(N_0\ell_1\ell_2)\backslash\mathfrak{H}} f_1(\ell_1 z)\overline{f_2(\ell_2 z)}y^k\cdot\varphi(z)\frac{dxdy}{y^2}\right|^2$$
$$= \frac{1}{\ell_1\ell_2}\int_{\Gamma_0(N_0\ell_1\ell_2)\backslash\mathfrak{H}} |\varphi(z)|^2\frac{dxdy}{y^2}$$
$$\times\frac{c(N_0)L^*(1/2, f_1\otimes\bar{f}_2\otimes\varphi)}{(\ell_1\ell_2)^{k+1}L^*(1,\operatorname{Ad}\varphi)}\left(1+o(\ell_1\ell_2)\right),$$

where the global L-functions appearing are completed with their gamma factors.

Setting $f_1 = f, f_2 = g$ and $\varphi = \bar{u}_j$, Proposition 4.3 gives us

$$|\langle V, u_j\rangle|^2 \ll (\ell_1\ell_2)^{-k-1}(1+|t_j|)^{2k-2}e^{-\pi|t_j|}L(1,\operatorname{Ad}u_j)^{-1}L(1/2, f\otimes\bar{g}\otimes\bar{u}_j). \tag{4.1}$$

Here we have used Stirling's formula for the gamma factors.

Thus Proposition 4.1 is essentially stating that

$$\frac{L(1/2, f\otimes\bar{g}\otimes\bar{u}_j)}{L(1,\operatorname{Ad}u_j)} \tag{4.2}$$

is close to constant on average. By [13] we have the bounds

$$(N(1+|t_j|))^{\epsilon} \gg L(1,\operatorname{Ad}u_j) \gg \frac{1}{\log((\ell_1\ell_2(1+|t_j|))}, \tag{4.3}$$

and also, equivalently, that

$$e^{\pi|t_j|}(N(1+|t_j|))^{-\epsilon} \ll |\rho_j(1)|^2 \ll e^{\pi|t_j|}\log((\ell_1\ell_2(1+|t_j|)). \tag{4.4}$$

As we expect $L(1/2, f\otimes\bar{g}\otimes\bar{u}_j)$ to obey the Lindelöf Hypothesis this evaluation is consistent with the average result given in Proposition 4.1.

Combining (4.4) with the upper bound $\lambda_j(h) \ll h^{\theta}$ for the h^{th} Hecke eigenvalue of u_j, gives

$$|\rho_j(-h)|^2 = |\rho_j(1)\lambda_j(h)|^2 \ll e^{\pi|t_j|}h^{2\theta}\log((\ell_1\ell_2(1+|t_j|))$$

and with Proposition 4.1, Cauchy-Schwarz, and Weyl's law this gives us

$$\sum_{|t_j|\sim T} \overline{\rho_j(-h)\langle V, u_j\rangle}$$

$$\ll \left(\sum_{|t_j|\sim T} |\rho_j(h)|^2 e^{-\pi|t_j|}\right)^{1/2} \left(\sum_{|t_j|\sim T} |\langle V, u_j\rangle|^2 e^{\pi|t_j|}\right)^{1/2}$$

$$\ll \left(\ell_1\ell_2 T^2 h^{2\theta}\log((\ell_1\ell_2(1+|t_j|)))\right)^{1/2}\left((\ell_1\ell_2)^{-k}T^{2k}\right)^{1/2}$$

$$\ll (\ell_1\ell_2)^{1/2-k/2}T^{1+k}h^{\theta}(\log(\ell_1\ell_2 T))^{1/2}. \quad (4.5)$$

The presence of h^θ can be avoided for certain choices of the parameters. An application of the Kuznetsov trace formula (see, for example, [19], Theorem 16.8, line (16.56)) shows that

$$\sum_{|t_j|\sim T} |\overline{\rho_j(h)}|^2 e^{-\pi|t_j|} \ll \ell_1\ell_2T^2 + T^{7/4}\sigma_0(\ell_1\ell_2)(h,\ell_1\ell_2)^{1/2}h^{1/2}\log^2(3h). \quad (4.6)$$

Thus, for ℓ_1, ℓ_2 prime, $\ell_1, \ell_2 \gg h^{1/4}\log(h)$, with $(h, \ell_1\ell_2) = 1$,

$$\sum_{|t_j|\sim T} |\overline{\rho_j(h)}|^2 e^{-\pi|t_j|} \ll \ell_1\ell_2 T^2,$$

implying that in this context $\lambda_j(h)$ is constant on average for varying j. Summarizing the above discussion, we have proved

Proposition 4.4.

$$\sum_{|t_j|\sim T} \overline{\rho_j(-h)\langle V, u_j\rangle} \ll (\ell_1\ell_2)^{1/2-k/2}T^{1+k}h^{\theta}(\log(\ell_1\ell_2 T))^{1/2},$$

where, if ℓ_1, ℓ_2 prime, $\ell_1, \ell_2 \gg h^{1/4}\log(h)$, with $(h, \ell_1\ell_2) = 1$, we can set $\theta = 0$.

Turning to the continuous part, a simple computation shows that

$$|\langle V, E_a(z, 1/2+it)\rangle|^2 \ll (\ell_1\ell_2)^{-k-1}(1+|t|)^{2k-2}e^{-\pi|t|}L(1/2+it, f\otimes\overline{g})^2, \quad (4.7)$$

which agrees with the average cuspidal bound (up to the central value $L(1/2, f\otimes g\otimes u_j)$.

When combined with (2.13) and Stirling's formula, (4.7) implies

$$\left|\sum_{a} \frac{\overline{\tilde{\rho}_a(1/2+it, -h_a)}\sigma_{2it}(h)\,\langle V, E_a(z, 1/2+it)\rangle}{\Gamma(1/2-it)\zeta(1-2it)}\right|$$

$$\ll (\ell_1\ell_2)^{-k/2-1+\epsilon}(1+|t|)^{k-1+\epsilon}L(1/2+it, f\otimes\overline{g}). \quad (4.8)$$

Recalling (2.8), (2.9) and (2.10) we define $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ by

$$\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h) = \mathcal{I}^{\mathrm{cusp}}_{\ell_1,\ell_2,\ell,\delta}(s;h) + \mathcal{I}^{\mathrm{cont}}_{\ell_1,\ell_2,\ell,\delta}(s;h), \tag{4.9}$$

where

$$\mathcal{I}^{\mathrm{cusp}}_{\ell_1,\ell_2,\ell,\delta}(s;h) = \sum_j \frac{\overline{\rho_j(-h)}}{\mathcal{V}(2\pi h)^{s-1/2}} M(s,t_j,\delta)\overline{\langle V, u_j\rangle} \tag{4.10}$$

and

$$\begin{aligned}\mathcal{I}^{\mathrm{cont}}_{\ell_1,\ell_2,\ell,\delta}&(s;h)\\ &= \frac{1}{4\pi}\sum_a \int_{-\infty}^{\infty} \frac{\overline{\tilde{\rho}_a(1/2+it,-h_a)}2\pi^{1-s-it}2^{3/2-s}h^{1/2-s-it}\sigma_{2it}(h)}{\mathcal{V}\Gamma(1/2-it)\zeta(1-2it)}\\ &\qquad\qquad \times M(s,t,\delta)\overline{\langle V, E_a(*,1/2+it)\rangle}dt. \end{aligned} \tag{4.11}$$

By Weyl's law

$$\sum_{|t_j|<T} 1 \ll \ell_1\ell_2 T^2. \tag{4.12}$$

By the first upper bound of Proposition 3.1, (3.2), for any fixed s and δ, s at least a distance ϵ from the poles, $M(s,t,\delta)$ decays faster than any power of $|t|$ as $|t| \to \infty$. This, together with (4.5), (4.12), and the convexity bound for the L-series shows that the right hand side of (4.10) converges absolutely. Similarly, by (4.8) the integral over t in (4.11) converges absolutely. The same argument, together with Proposition 2.1, shows that choosing A sufficiently large will ensure that the expression for $\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h)$ in (2.8) converges absolutely and uniformly for any fixed Y, as long as $\mathfrak{Re}\, s > 1/2$, and that the difference, $|\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h) - \mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h)|$, converges absolutely, allowing the interchange of the limit as $Y \to \infty$ with the summation. As the limit of each term in the summation is 0 as $Y \to \infty$ this shows that for fixed $\delta > 0$ and s with $\mathfrak{Re}\, s > 1/2$, $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ is an analytic function of s and

$$\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h) = \lim_{Y\to\infty} \mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h).$$

Recall that in (12.1) we had an expression for $\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h)$ as a Dirichlet series. For $\mathfrak{Re}\, s > 1$ it is clear that the sum converges and that a limit exists as $Y \to \infty$, namely the analytic function

$$\begin{aligned}\lim_{Y\to\infty}\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h) &= \frac{1}{\mathcal{V}}(4\pi)^{-(s+k-1)}\sum_{m_1\ell_1=m_2\ell_2+h}\frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}}\\ &\quad\times\int_0^\infty e^{-y(1+\delta h/(2\ell_2 m_2))}y^{s+k-1}\frac{dy}{y}.\\ &= \frac{\Gamma(s+k-1)}{(4\pi)^{s+k-1}\mathcal{V}}D(s;h,\delta),\end{aligned}$$

where

$$D(s;h,\delta) = \sum_{m_1\ell_1=m_2\ell_2+h}\frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}}\left(1+\frac{\delta h}{2\ell_2 m_2}\right)^{-(s+k-1)} \tag{4.13}$$

converges absolutely for $\mathfrak{Re}\, s > 1$.

We have now established that the function $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ defined by (4.9) is an infinite sum that converges absolutely when $\mathfrak{Re}\, s > 1/2$ and is related to $D(s;h,\delta)$ by

$$D(s;h,\delta) = \frac{(4\pi)^{s+k-1}\mathcal{V}}{\Gamma(s+k-1)}\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h), \tag{4.14}$$

when $\mathfrak{Re}\, s > 1$.

The meromorphic continuation of $D(s;h,\delta)$ is related to that of $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ via (4.14). Using (4.9),(4.10), (4.11) we have the following spectral expansion:

$$\begin{aligned}D(s;h,\delta) &= \frac{(4\pi)^{s+k-1}}{\Gamma(s+k-1)(2\pi h)^{s-1/2}}\Bigg(\sum_j\overline{\rho_j(-h)}M(s,t_j,\delta)\overline{\langle V,u_j\rangle}\\ +&\quad\frac{1}{4\pi}\sum_a\int_{-\infty}^{\infty}\frac{\overline{\tilde{\rho}_a(1/2+it,-h_a)}2\pi^{1-s-it}2^{3/2-s}h^{1/2-s-it}\sigma_{2it}(h)}{\mathcal{V}\Gamma(1/2-it)\zeta(1-2it)}\\ \times&\quad M(s,t,\delta)\overline{\langle V,E_a(*,1/2+it)\rangle}dt\Bigg).\end{aligned} \tag{4.15}$$

By (3.2), (4.5), (4.8), (4.9), for any sufficiently large fixed $A \gg 1$, the series expressions for $\mathcal{I}^{\mathrm{cusp}}_{\ell_1,\ell_2,\ell,\delta}(s;h)$, $\mathcal{I}^{\mathrm{cont}}_{\ell_1,\ell_2,\ell,\delta}(s;h)$, $D(s;h,\delta)$ given in (4.10),(4.11), (4.15) converge absolutely for s with $\mathfrak{Re}\, s > 1/2 - A$, but the upper bound can have a factor of δ^{-A} in it. As we intend to let $\delta \to 0$ it is useful to identify an interval for $\mathfrak{Re}\, s$ in which there is absolute convergence with an upper bound that is independent of δ.

First, combining (3.3) and (3.4) we have, for $\mathfrak{Re}\, s = \sigma \le 1/2$, regardless of the relation between δ and $(1+|t|)^{-2}$,

$$M(s,t_j,\delta) \ll (1+|t|)^{2\sigma-2+2\epsilon}(1+|s|)^{A-1/2}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}. \tag{4.16}$$

Combining this with Proposition 4.4, we have, for any $T \gg 1$,

$$\left| \sum_{|t_j| \sim T} \overline{\rho_j(-h)} M(s, t_j, \delta) \overline{\langle V, u_j \rangle} \right| \ll (\ell_1 \ell_2)^{(1-k)/2} T^{2\sigma + 2\epsilon + k - 1}$$
$$\times \quad h^{\theta} \log(\ell_1 \ell_2 T) e^{-\frac{\pi}{2} |\mathfrak{Im}\, s|} (1 + |s|)^{A - 1/2}). \tag{4.17}$$

Similarly, applying (4.8) in place of the proposition, and simply using the convexity bound (notice ℓ_1, ℓ_2 do not appear in the conductor of $L(1/2, f \otimes g)$), we obtain

$$\sum_a \frac{\overline{\tilde{\rho}_a(1/2 + it, -h_a) 2\pi^{1-s-it} 2^{3/2-s} h^{1/2-s-it} \sigma_{2it}(h)}}{\mathcal{V} \Gamma(1/2 - it) \zeta(1 - 2it)} M(s, t, \delta)$$
$$\times \overline{\langle V, E_a(*, 1/2 + it) \rangle}$$
$$\ll (\ell_1 \ell_2)^{-1-k/2+\epsilon} (1 + |t|)^{2\sigma - 2 + k + \epsilon} e^{-\frac{\pi}{2} |\mathfrak{Im}\, s|} (1 + |s|)^{A - 1/2}). \tag{4.18}$$

Breaking up the interval from 1 to T diadically and letting $T \to \infty$ it is clear from (4.17) that the cuspidal part of the spectral expansion of $D(s; h, \delta)$ converges absolutely when $1/2 - A < \mathfrak{Re}\, s < 1/2 - k/2 - \epsilon$. Referring to (4.18), we see that the continuous part also converges in this region and satisfies a better upper bound. Thus (4.17), together with (4.18), (4.15) and Stirling's formula provide the upper bound for $D(s; h, \delta)$ of Proposition 4.5 stated below in (4.19).

The details of the above discussion are summarized, and some additional information is added, in the following

Proposition 4.5. *The functions $\mathcal{I}_{\ell_1, \ell_2, \ell, \delta}(s; h)$ and $D(s; h, \delta)$, defined in* (4.9), (4.13) *and* (4.14)*, have meromorphic continuations to the half plane $\mathfrak{Re}\, s > 1/2 - A$, with simple poles at $s = 1/2 \pm it_j - r$, for each t_j and each $0 \le r < A$. Letting*

$$R(s_0; h, \delta) = \mathrm{Res}_{s = s_0} D(s; h, \delta)$$

we have

$$\begin{aligned} & R(1/2 + it_j - r; h, \delta) \\ = \quad & \frac{(-1)^r 2^{2k} \pi^k h^{r - it_j} \overline{\rho_j(-h) \langle V, u_j \rangle} \Gamma(1/2 - it_j + r) \Gamma(2it_j - r)}{r! \Gamma(1/2 + it_j) \Gamma(1/2 - it_j) \Gamma(k - 1/2 + it_j - r)} \\ + \quad & \mathcal{O}\left((1 + |t_j|)^r h^r \overline{\rho_j(-h) \langle V, u_j \rangle} \delta^{1/2} \right). \end{aligned}$$

In the region $1/2 - A + \epsilon < \mathfrak{Re}\, s = \sigma < 1/2 - k/2$, for s a distance of at least ϵ from the poles the spectral expansion for $D(s; h, \delta)$ given *in* (4.15) *converges absolutely and satisfies the upper bound*

$$D(s;h,\delta) \ll (1+|s|)^{A-1/2}(\log(\ell_1\ell_2))^{1/2}(\ell_1\ell_2)^{1/2-k/2}h^{1/2-\sigma+\theta}. \tag{4.19}$$

For $2 \geq \sigma \geq 1/2-A$ *and* s *at least* ϵ *from the poles*

$$D(s;h,\delta) \ll (1+|s|)^{A-k+1/2}\delta^{-A}(\ell_1\ell_2)^{1/2-k/2}\log(\ell_1\ell_2)h^{\theta+1/2-\sigma} \tag{4.20}$$

In the preceding two inequalities the exponent θ *can be set equal to zero if* $\ell_1,\ell_2 \gg h^{1/4}$ *and* $(\ell_1\ell_2,h)=1$.

For $\sigma > 1+\epsilon$, $D(s;h,\delta)$ *satisfies the bound*

$$D(s;h,\delta) \ll \frac{\left(1+\frac{h}{\ell_2}\right)^{(k-1)/2}}{(\ell_1\ell_2)^{(k-1)/2}\,\ell_2^{1+\epsilon}}. \tag{4.21}$$

Proof. Proposition 3.1, (3.2), states that for sufficiently large A, and fixed $\delta>0$ and s satisfying $\mathfrak{Re}\, s > 1/2-A$, with s at least a distance ϵ from the poles, the meromorphic continuation of $M(s,t,\delta)$ decays faster than any power of $|t|$. Thus for any fixed $\delta>0$ and such an s with $\mathfrak{Re}\, s > 1/2-A$, the series expression for $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ given in (4.9) converges absolutely, giving a meromorphic continuation of $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ back to $\mathfrak{Re}\, s > 1/2-A$, with possible simple poles at the points $s=1/2\pm it_j-r$, for each t_j and each $0\leq r<A$. The corresponding meromorphic continuation of $D(s;h,\delta)$ follows from (4.14). The residues at these points of $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$, and consequently of $D(s;h,\delta)$, follow from the corresponding residue evaluations of $M(s,t,\delta)$ given in Proposition 3.1.

The piece corresponding to the continuous part of the spectrum, $\mathcal{I}^{\mathrm{cont}}_{\ell_1,\ell_2,\ell,\delta}(s;h)$, defined in (4.11), does not contribute any poles, for the following reason. Any pole in s must arise from the term $M(s,t,\delta)$, and in fact must arise from the main term of (3.3) in Proposition 3.1. We thus take the spectral part of $D(s;h,\delta)$, namely

$$\frac{1}{4\pi}\sum_{a}\int_{-\infty}^{\infty}\frac{\overline{\tilde{\rho}_a(1/2+it,-h_a)}2\pi^{1-s-it}2^{3/2-s}h^{1/2-s-it}\sigma_{2it}(h)}{\mathcal{V}\Gamma(1/2-it)\zeta(1-2it)} \times M(s,t,\delta)\overline{\langle V,E_a(*,1/2+it)\rangle}dt.$$

and note that any polar contribution from this must come from the piece of the integrand

$$\frac{\Gamma(s-1/2+it)\Gamma(s-1/2-it)\overline{\langle V,E_a(*,1/2+it)\rangle}}{\zeta(1-2it)\Gamma(1/2-it)\Gamma(1/2+it)\Gamma(1/2-it)}.$$

This is more easily analyzed when the Eisenstein series is multiplied by the normalizing factor $\zeta^*(1+2it)$. The potential polar contribution is then

$$\frac{\Gamma(s-1/2+it)\Gamma(s-1/2-it)\overline{\langle V, E_a^*(*,1/2+it)\rangle}}{\zeta^*(1+2it)\zeta^*(1-2it)\Gamma(1/2+it)\Gamma(1/2-it)}.$$

For this to have a pole in s it must be the case that $\mathfrak{Re}\, s = 1/2 - r$, for some $r \geq 0$. To understand the behavior on this line, write $s = 1/2 - r + \epsilon + i\gamma$, for some $\epsilon > 0$, set $u = it$, and move the line of integration from $\mathfrak{Re}\, u = 0$ to $\mathfrak{Re}\, u = -2\epsilon$. In this process a pole is passed over at $u = -\epsilon - i\gamma$, and possibly at points where $\zeta^*(1+2u) = 0$. The moved integral now has no poles in s on the line $\mathfrak{Re}\, s = 1/2 - r$ and the residue at $u = -\epsilon - i\gamma$ is an analytic multiple of

$$\sum_a \frac{\Gamma(-r+2\epsilon+2i\gamma)\overline{\langle V, E_a(*,1/2-\epsilon-i\gamma)\rangle}}{\zeta(1+2\epsilon+2i\gamma)\Gamma(1/2+\epsilon+i\gamma)^2\Gamma(1/2-\epsilon-i\gamma)}.$$

For a pole to exist when $\epsilon \to 0$, it must be the case that $\gamma = 0$, and the potential pole of $\Gamma(-r+2\epsilon)$ in the numerator is cancelled by the $\zeta(1+2\epsilon)\zeta(1-2\epsilon)$ in the denominator. Any extra residues coming from poles at $\zeta^*(1+2u) = 0$ will be subtracted as $\epsilon \to 0$.

In addition to locating the possible poles, (3.2) in Proposition 3.1 allows us to establish an upper bound for $\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)$ when s is in the region $\mathfrak{Re}\, s > 1/2 - A$ and s is at least a distance of ϵ from the poles. In particular, combining this information with (4.5) and (4.9), we obtain for such s:

$$\begin{aligned}&\mathcal{I}_{\ell_1,\ell_2,\ell,\delta}(s;h)\\ &\ll (1+|s|)^{A+\mathfrak{Re}\, s-1}\delta^{-A}e^{-\frac{\pi}{2}|\mathfrak{Im}\, s|}(\ell_1\ell_2)^{-k/2-1/2}\log(\ell_1\ell_2)h^{\theta+1/2-s}, \quad (4.22)\end{aligned}$$

with the implied constant depending only on A. Combining (4.14), (4.22) and Stirling's formula, gives us (4.20) of the Proposition.

If $1/2 - A < \sigma < 1/2 - k/2$ then by the argument preceding the proposition we obtain (4.19). It is a key point here that this estimate is independent of δ.

For $\sigma > 1 + \epsilon$, as f, g are modular forms and satisfy the Ramanujan Conjecture, it is easily verified that the bound (4.21) in the proposition holds. □

5. THE LIMIT AS $\delta \to 0$

The object of the section is to prove the following

Proposition 5.1. *For $\mathfrak{Re} s > 1$, let*

$$D(s;h) = \lim_{\delta\to 0} D(s;h,\delta) = \sum_{m_1\ell_1 = m_2\ell_2 + h} \frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}}. \tag{5.1}$$

The function $D(s;h)$ converges for $\mathfrak{Re}\, s > 1$ and has an analytic continuation to all s with $\mathfrak{Re} s > 1/2 - A$, except for the points $s = 1/2 \pm it_j - r$, with $0 \le r < A$, $r \in \mathbb{Z}$. At these points there are simple poles and

$$R(1/2+it_j - r; h) = \mathrm{Res}_{s=1/2+it_j-r}\, D(s;h) = c_{r,j} h^{r-it_j}\overline{\lambda_j(h)}, \tag{5.2}$$

where

$$c_{r,j} = \frac{(-1)^r 2^{2k}\pi^k \overline{\rho_j(-1)\langle V, u_j\rangle}\Gamma(1/2 - it_j + r)\Gamma(2it_j - r)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)\Gamma(k-1/2+it_j-r)}. \tag{5.3}$$

For $T \gg 1$,the $c_{r,j}$ satisfy the average upper bounds

$$\sum_{|t_j|\sim T} |c_{r,j}|^2 \ll \log(\ell_1\ell_2 T)(\ell_1\ell_2)^{-k} T^{2r+1-k} \tag{5.4}$$

For s at least ϵ away from the polar points, and in the vertical strip $1/2 - A < \sigma < 1/2 - -k/2$, $D(s;h)$ can be expressed by the following absolutely convergent sum and integral:

$$\begin{aligned} D(s;h) = {} & \frac{2^{2k-1}\pi^k h^{1/2-s}\Gamma(1-s)}{\Gamma(s+k-1)} \\ & \times \Bigg(\sum_j \overline{\lambda_j(h)\rho_j(-1)} \frac{\Gamma(s-1/2+it_j)\Gamma(s-1/2-it_j)}{\Gamma(1/2+it_j)\Gamma(1/2-it_j)} \overline{\langle V, u_j\rangle} \\ & + \frac{1}{4\pi}\sum_{a}\int_{-\infty}^{\infty} \frac{\overline{\tilde{\rho}_a(1/2+it,-h_a)}2^{3/2-s}\pi^{1-s-it}h^{1/2-s-it}\sigma_{2it}(h)}{\mathcal{V}\zeta(1-2it)\Gamma(1/2-it)^2\Gamma(1/2+it)} \\ & \qquad \times \Gamma(s-1/2+it)\Gamma(s-1/2-it)\overline{\langle V, E_a(*,1/2+it)\rangle} dt \Bigg). \end{aligned}$$

In this region $D(s;h)$ satisfies the upper bound

$$D(s;h) \ll (1+|s|)^{A-1/2}(\log(\ell_1\ell_2))^{1/2}(\ell_1\ell_2)^{1/2-k/2}h^{1/2-\sigma+\theta}. \tag{5.5}$$

For s at least ϵ away from the polar points, and in the vertical strip $c < \sigma < 2$, for any c with $1/2 - A + \epsilon < c < 1/2 - k/2$, $D(s;h)$ is given *by*

$$\begin{aligned} D(s;h) &= \frac{1}{2\pi i}\int_{2-i\infty}^{2+i\infty}\frac{D(u;h)e^{(u-s)^2}du}{u-s} \\ &- \frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty}\frac{D(u;h)e^{(u-s)^2}du}{u-s} \\ &- \sum_{t_j,0\le r<1/2-c}\frac{R(1/2+it_j-r;h)e^{(1/2+it_j-r-s)^2}}{1/2+it_j-r-s} \end{aligned} \tag{5.6}$$

and, if $\ell_1,\ell_2\sim L$, satisfies the upper bound

$$D(s;h) \ll (1+|s|)^{-c-k+3/2}(\log(L))^{1/2}L^{1-k}h^{1/2-c+\theta}. \tag{5.7}$$

Here in the integral with $\mathfrak{Re}\, u=2$, $D(u;h)$ is given by (5.1)*, and in the integral with $\mathfrak{Re}\, u=c$, $D(u;h)$ is given by* (5.6)*. In the preceding two upper bounds the exponent θ can be set equal to zero if $\ell_1,\ell_2\gg h^{1/4}$ and $(\ell_1\ell_2,h)=1$.*

Proof. Using the information in Proposition 4.5, the Cauchy residue theorem allows us to express each $D(s;h,\delta)$, with $1/2-A+\epsilon<\mathfrak{Re}\, s<2$, as follows. Set $T_\delta=\delta^{-1}$. Then

$$\begin{aligned} D(s;h,\delta) &+ \sum_{|t_j|<T_\delta,0\le r<A}\frac{R(1/2+it_j-r;h,\delta)e^{(1/2+it_j-r-s)^2}}{1/2+it_j-r-s} \\ &= I_1-I_2-I_3+I_4, \end{aligned} \tag{5.8}$$

where

$$I_1=\frac{1}{2\pi i}\int_{2-iT_\delta}^{2+iT_\delta}\frac{D(u;h,\delta)e^{(u-s)^2}du}{u-s},$$

$$I_2=\frac{1}{2\pi i}\int_{c+iT_\delta}^{2+iT_\delta}\frac{D(u;h,\delta)e^{(u-s)^2}du}{u-s},$$

$$I_3=\frac{1}{2\pi i}\int_{c-iT_\delta}^{c+iT_\delta}\frac{D(u;h,\delta)e^{(u-s)^2}du}{u-s},$$

and

$$I_4=\frac{1}{2\pi i}\int_{c-iT_\delta}^{2-iT_\delta}\frac{D(u;h,\delta)e^{(u-s)^2}du}{u-s}.$$

By (4.20)

$$I_2\ll(1+|s|)^{A-k+1/2}\delta^{-A}(\ell_1\ell_2)^{(1-k)/2}\log(\ell_1\ell_2)h^{\theta+1/2-\sigma}e^{-T_\delta{}^2}.$$

As $T_\delta=\delta^{-1}$ the exponential decay of the $e^{-T_\delta{}^2}$ overcomes the polynomial growth of the δ^A, and $I_2\to 0$ as $\delta\to 0$. Similarly $\lim_{\delta\to 0}I_4=0$

By (4.13), we see that as $\mathfrak{Re}(s+u) > 1$ the limit as $\delta \to 0$ of the integrand of I_1 exists. By the absolute convergence of $D(\sigma+it;h)$ for $\sigma > 1$ and the exponential decay of e^{u^2} the limit as $\delta \to 0$ of I_1 exists.

Turning to I_3 and referring to (3.3) we see that for any fixed t, and $\mathfrak{Re}\, s \leq 1/2$, $\lim_{\delta\to 0} M(s,t,\delta)$ exists and

$$\begin{aligned} M(s,t) &= \lim_{\delta\to 0} M(s,t,\delta) \\ &= \frac{\sqrt{\pi}2^{1/2-s}\Gamma(s-1/2-it)\Gamma(s-1/2+it)\Gamma(1-s)}{\Gamma(1/2-it)\Gamma(1/2+it)}. \end{aligned} \tag{5.9}$$

Referring to Proposition 4.5, line (4.19), it is clear that in the given range the spectral expansion for $D(s;h)$ given in the proposition converges absolutely, and that the limit as $\delta \to 0$ of $D(s;h,\delta)$ exists, equals $D(s;h)$, and satisfies the upper bound (5.5). The at most polynomial growth of $D(s;h)$ in the imaginary part of s shown in (5.5), together with the exponential decay of e^{u^2} shows that for $1/2 - A + \epsilon < \mathfrak{Re}\, s < 1/8 - k/2$, the limit as $\delta \to 0$ of I_3 exists and equals the integral of the limit. The Dirichlet series expansion of $D(u;h,\delta)$ in integral I_1 converges absolutely as $\mathfrak{Re}\, u > 1$, and thus similarly the limit can be interchanged with the integral and sum.

By the description in Proposition 4.5, each residue term $R(1/2 + it_j - r; h, \delta)$ grows at most polynomially in $|t_j|$ for any fixed s with $1/2 - A + \epsilon < \mathfrak{Re}\, s < 2$. As $e^{(1/2+it_j-r-s)^2}$ decays exponentially in $|t_j|$, the sum over the residues converges absolutely and approaches the limits given above in (5.2) and (5.6). We have thus established that for any s in the range $1/2 - A + \epsilon < \mathfrak{Re}\, s < 2$, the limit as $\delta \to 0$ of $D(s;h,\delta)$ exists, and equals the analytic function $D(s;h)$ defined in (5.6).

The average upper bounds for the $c_{r,j}$ are obtained by an application of Stirling's formula, and Propositions 4.1 and 4.4. It follows then that

$$\sum_{t_j, 0\leq r<1/2-c} \frac{R(1/2+it_j-r;h)e^{(1/2+it_j-r-s)^2}}{1/2+it_j-r-s} \ll (\log(\ell_1\ell_2))^{1/2}(\ell_1\ell_2)^{1/2-k/2}h^{1/2-\sigma+\theta}.$$

By (5.5), the integral along $\mathfrak{Re}\, s = c$ is

$$\ll (1+|s|)^{-\sigma-k+3/2}(\log(\ell_1\ell_2))^{1/2}(\ell_1\ell_2)^{1/2-k/2}h^{1/2-\sigma+\theta},$$

and by (4.21), the integral along $\mathfrak{Re}\, s = 2$ is

$$\ll \frac{\left(1+\frac{h}{\ell_2}\right)^{(k-1)/2}}{\ell_2(\ell_1\ell_2)^{(k-1)/2}} \ll \ell_2^{-1}h^{(k-1)/2}(\ell_1\ell_2)^{-(k-1)/2} \ll h^{(k-1)/2}L^{-k}.$$

Noting that $1/2-c > (k-1)/2$, it follows from the previous three lines that for any s with $c < \mathfrak{Re}\, s < 2$, $D(s;h)$ satisfies the upper bound (5.7). $\square$

6. Estimating shifted convolution sums

For $x > 0$ and $\ell_1, \ell_2 \sim L$, let

$$S(x;h) = \frac{(\ell_1\ell_2)^{(k-1)/2}}{2\pi i}\int_{2-i\infty}^{2+i\infty} D(s;h)g(s)(\ell_2 x)^s ds, \tag{6.1}$$

where

$$G(x) = \frac{1}{2\pi i}\int_{2-i\infty}^{2+i\infty} g(s)x^s ds$$

is smooth and has compact support in the interval $[1,2]$.

As $\mathfrak{Re}\, s > 1$, the Dirichlet series and integral converge absolutely, so

$$\begin{aligned} S(x;h) &= (\ell_1\ell_2)^{(k-1)/2}\sum_{m_1\ell_1=\ell_2m_2+h}\frac{a(m_1)\bar{b}(m_2)}{(\ell_2m_2)^{k-1}}G\left(\frac{x}{m_2}\right) \qquad (6.2)\\ &= \sum_{m_1\ell_1=\ell_2m_2+h} A(m_1)\bar{B}(m_2)\left(1+\frac{h}{\ell_2m_2}\right)^{(k-1)/2} G\left(\frac{x}{m_2}\right). \end{aligned}$$

where, as mentioned before,

$$a(m_1) = A(m_1)m_1^{(k-1)/2},\ b(m_1) = B(m_2)m_2^{(k-1)/2}.$$

Note that the sum is over $m_2 \sim x$ such that $\ell_2 m_2 \equiv -h \pmod{\ell_1}$ and thus the length of this sum should be $\mathcal{O}(x/\ell_1)$. The nature of the sum also changes with h. For $h \ll \ell_2 x$ each term is $\mathcal{O}(1)$, while for $h \gg \ell_2 x$ each term is $\mathcal{O}((h/(\ell_2 x))^{(k-1)/2})$.

By (5.7) the integral in (6.1) continues to converge absolutely if we move the line of integration to the left, as $g(s)$ decays in a vertical line faster than any fixed power of s. We thus move the line of integration to $\mathfrak{Re} s = 1/2-k/2-\epsilon$, picking up residues at the poles of $D(s;h)$ and obtaining

$$\begin{aligned} &S(x;h)\\ &= I+(\ell_1\ell_2)^{(k-1)/2}\sum_{t_j, 0\le r\le k/2} c_{r,j}h^{r-it_j}\overline{\lambda_j(h)}g(1/2+it_j-r)(\ell_2x)^{1/2+it_j-r} \end{aligned} \tag{6.3}$$

where

$$I = \frac{(\ell_1\ell_2)^{(k-1)/2}}{2\pi i}\int_{1/2-k/2-\epsilon-i\infty}^{1/2-k/2-\epsilon+i\infty} D(s;h)g(s)(\ell_2x)^s ds. \tag{6.4}$$

By Proposition 5.1, the integral converges absolutely and

$$I \ll \log(L) h^{\theta+1/2}(h/Lx)^{k/2-1/2+\epsilon}. \tag{6.5}$$

Thus

$$\begin{aligned} &S(x;h) \\ &= (\ell_1\ell_2)^{(k-1)/2} \sum_{t_j, 0\le r<1+k/2} c_{r,j} h^{1/2}\overline{\lambda_j(h)} g(1/2+it_j-r)(h/\ell_2 x)^{r-1/2-it_j} \\ &\qquad\qquad + \mathcal{O}\left(\log(L) h^{\theta+1/2}(h/Lx)^{k/2-1/2+\epsilon}\right), \end{aligned}$$

where the $c_{r,j}$ are described above in Proposition 5.1.

If $h \ll \ell_2 x$ then the terms in the summation part of (6.3) decrease in size as r increases and dominate the error term. Thus for $h \ll \ell_2 x$ the dominant term corresponds to the residue term at $r = 0$. By (5.4) it follows that for $h \ll Lx$

$$S(x;h) \ll (\log(L))^{1/2} h^{\theta} x^{1/2+\theta}.$$

For $h \gg \ell_2 x$ the main terms in (6.3) increase in size as r increases and the error term dominates. Thus

$$S(x;h) \ll (\log(L))^{1/2} h^{\theta+1/2}(h/Lx)^{k/2-1/2+\epsilon}. \tag{6.6}$$

The θ here denotes the best known progress toward the *non-archimedian* Ramanujan-Petersson conjecture. Recall that it can be set to zero if $\ell_1, \ell_2 \gg h^{1/4}$.

One can then ask when the upper bound in (6.6) is non-trivial. Setting $\ell_1 = \ell_2 = 1$, the length of the sum is $\mathcal{O}(x)$ and each term in the sum is of order $(h/x)^{(k-1)/2}$. This means that the trivial upper bound for the left hand side of (6.6) is $x(h/x)^{(k-1)/2}$. The result is thus non-trivial when

$$x(h/x)^{(k-1)/2} \gg h^{1/2+\theta}(h/x)^{k/2-1/2+\epsilon},$$

i.e when $h \ll x^{2/(1+2\theta)}$.

This corresponds to the result of Corollary 1.4 in [2], confirming a curious "wall" blocking progress past $h \ge x^2$. The ability to improve the result past this wall seems to be tied to the finding of a subconvexity upper bound for $D(s;h)$ in the region $-1/2-k/2 < \mathfrak{Re}\, s < 0$. However, the wall goes away when an average is taken over h as will be seen shortly.

7. A double Dirichlet series

Recall that

$$D(s;h) = \sum_{m_1\ell_1=m_2\ell_2+h} \frac{a(m_1)\bar{b}(m_2)}{(m_2\ell_2)^{s+k-1}} \tag{7.1}$$

is absolutely convergent when $\mathfrak{Re}\, s > 1$.

Fix a positive integer Q and an integer ν, with $0 \leq \nu \leq Q-1$. For s, w with real parts greater than 1 the sum

$$Z_Q(s,w;\nu) = (\ell_1\ell_2)^{(k-1)/2} \sum_{h\geq 1, h\equiv \nu \pmod{Q}} \frac{D(s;h)}{h^{w+(k-1)/2}} \tag{7.2}$$

is absolutely convergent. In future work we will study this series with general ν, but for the moment we will restrict ourselves to the case $\nu = 0$, that is, to

$$Z_Q(s,w;0) = (\ell_1\ell_2)^{(k-1)/2} \sum_{h_0\geq 1} \frac{D(s;h_0Q)}{(h_0Q)^{w+(k-1)/2}} \tag{7.3}$$

and will rewrite $Z_Q(s,w;0)$ in the more convenient form

$$Z_Q(s,w) = \sum_{\substack{h_0\geq 1\\ \ell_1m_1=\ell_2m_2+h_0Q}} \frac{A(m_1)\bar{B}(m_2)\left(1+\frac{h_0Q}{\ell_2m_2}\right)^{(k-1)/2}}{(\ell_2m_2)^s(h_0Q)^{w+(k-1)/2}}. \tag{7.4}$$

The object of this section is to obtain the meromorphic continuation of $Z_Q(s,w)$ to $\mathbb{C}^2$. The first step in this direction is

Proposition 7.1. *Choose c, with $1/2 - A < c < 1/2 - k/2$. The function $Z_Q(s,w)$ continues to the region $\mathfrak{Re}\, s > c$, $\mathfrak{Re}\, w > 1$. In this region it is analytic, with the exception of simple poles at the points $s = 1/2 - r \pm it_j$, for $0 \leq r < 1/2 - c$ and a possible simple pole at $s' = s+w+k/2-1 = 1/2$ in the case $f = g$. At the points $1/2-r+it_j$*

$$\begin{aligned}&\mathrm{Res}_{s=1/2-r+it_j} Z_Q(s,w)\\ &\qquad = (\ell_1\ell_2)^{(k-1)/2} c_{r,j} L_Q(w+(k-1)/2-r+it_j, \overline{u_j};0).\end{aligned}$$

For $\mathfrak{Re}\, s' \geq 1/2$ and $\mathfrak{Re}\, w > 1$

$$Z_Q(s,w) \ll Q^{\theta-s'} L^{\epsilon}(1+|s|)^{1-k}(1+|s'|)^{1-2s'}. \tag{7.5}$$

For $\mathfrak{Re}\, s' < 1/2$ and $\mathfrak{Re}\, w > 1$

$$Z_Q(s,w) \ll Q^{\max(\theta-s',-2s')} L^{1-2s'+\epsilon}(1+|s|)^{1-k}(1+|s'|)^{1-2s'}.$$

When $\ell_1\ell_2 \gg Q^{1/4}\log(Q)$, with $(Q,\ell_1\ell_2)=1$, and $\mathfrak{Re}\, s' \geq 0$, the θ can be removed in the previous two inequalities.

In the proposition, for $\mathfrak{Re}\, s' > 1$

$$L_Q(s', \overline{u_j}) = \sum_{h_0 \geq 1} \frac{\overline{\lambda_j(h_0 Q)}}{(h_0 Q)^{s'}}$$
$$= Q^{-s'} L(s', \overline{u_j}) \prod_{p^\gamma || Q} \left(\overline{\lambda_j(p^\gamma)} - \overline{\lambda_j(p^{\gamma-1})} p^{-s'} \right), \quad (7.6)$$

where $L(s', \overline{u_j})$ is the usual L-series associated to $\overline{u_j}$ (without gamma factors).

Proof. Fix some c, $c < 1/2 - k/2$. In the range $\mathfrak{Re}\, s < c$ the spectral decomposition of $D(s; h)$ converges. Recalling the analytic continuation of $D(s; h)$ given in Proposition 5.1 we define $\tilde{D}(s; h)$ by

$$\tilde{D}(s; h) = D(s; h) e^{s^2} - \sum_{t_j, 0 \leq r < 1/2 - c} \frac{c_{r,j} h^{r - it_j} \overline{\lambda_j(h)} e^{(1/2 - r + it_j)^2}}{s - (1/2 - r + it_j)}. \quad (7.7)$$

The function e^{s^2} has been included to insure that the sum over t_j above converges. A polynomially decaying function in the imaginary part of s would suffice, but e^{s^2} is convenient as it introduces no new poles.

By construction, $\tilde{D}(s; h)$ is analytic for all s with $\mathfrak{Re}\, s > c$. Let

$$\tilde{Z}_Q(s, w) = (\ell_1 \ell_2)^{(k-1)/2} \sum_{h_0 \geq 1} \frac{\tilde{D}(s; h_0 Q)}{(h_0 Q)^{w + (k-1)/2}}. \quad (7.8)$$

By Proposition 5.1, for $c < \mathfrak{Re}\, s < 2$, the sum over h_0 in $Z_Q(s, w)$ converges absolutely when $\mathfrak{Re}\, w + (k-1)/2 > 3/2 + \theta - c$. The sum

$$\sum_{h_0 \geq 1} \sum_{t_j, 0 \leq r < 1/2 - c} \frac{c_{r,j} (h_0 Q)^{r - it_j} \overline{\lambda_j(h_0 Q))} e^{(1/2 - r + it_j)^2}}{(h_0 Q))^{w + (k-1)/2} (s - 1/2 + r - it_j)}$$

also converges absolutely when $\mathfrak{Re}\, w + (k-1)/2 > 3/2 + \theta - c$. Thus $\tilde{Z}_Q(s, w)$ converges absolutely and is an analytic function in the region $c < \mathfrak{Re}\, s < 2$, $\mathfrak{Re}\, w + (k-1)/2 > 3/2 + \theta - c$.

By (7.7)

$$\tilde{Z}_Q(s, w) = Z_Q(s, w) e^{s^2} - P_Q(s, w), \quad (7.9)$$

where

$$
\begin{aligned}
&P_Q(s,w)\\
&= (\ell_1\ell_2)^{(k-1)/2} \sum_{t_j, 0\le r<1/2-c, h_0\ge 1} \frac{c_{r,j}(h_0Q)^{r-it_j}\overline{\lambda_j(h_0Q)}e^{(1/2-r+it_j)^2}}{(h_0Q)^{w+(k-1)/2}(s-(1/2-r+it_j))}\\
&\qquad = (\ell_1\ell_2)^{(k-1)/2}\\
&\times \sum_{t_j,0\le r<1/2-c} \frac{c_{r,j}L_Q(w+(k-1)/2-r+it_j,\overline{u_j})e^{(1/2-r+it_j)^2}}{s-(1/2-r+it_j)}. \qquad (7.10)
\end{aligned}
$$

By construction

$$
\begin{aligned}
&\mathrm{Res}_{s=1/2-r+it_j} Z_Q(s,w)e^{s^2} = \mathrm{Res}_{s=1/2-r+it_j} P_Q(s,w) \qquad (7.11)\\
= \ &(\ell_1\ell_2)^{(k-1)/2}c_{r,j}L_Q(w+(k-1)/2-r+it_j,\overline{u_j})e^{(1/2-r+it_j)^2}.
\end{aligned}
$$

The interchange in the order of summation above is allowed if the real part of w is sufficiently large. Once interchanged, it is clear that (given the analytic continuation of $L_Q(w+(k-1)/2-r+it_j,\overline{u_j})$), the summation converges absolutely for any fixed value of w.

It is now clear that the sum defining $P_Q(s,w)$ converges absolutely and is, again by construction, analytic for all points $s, w \in \mathbb{C}$ with $\mathfrak{Re}\, s > c$, except for the points $s = 1/2 - r + it_j$, where it has polar lines.

Applying Proposition 5.1 again, we are in a range where each $D(s; h_0Q)$ can be represented by an absolutely convergent sum, when s is a distance of at least ϵ from any pole. Interchanging this sum with the sum over h_0 we obtain, for $\mathfrak{Re}\, w$ sufficiently large, an expansion for $Z_Q(s,w)$. Writing $s' = w + s + k/2 - 1$ to simplify this and further equations, we have:

$$Z_Q(s,w) = (\ell_1\ell_2)^{(k-1)/2}\left(S_{\text{cusp}} + S_{\text{cont}}\right), \qquad (7.12)$$

where

$$
\begin{aligned}
S_{\text{cusp}} &= \frac{2^{2k-1}\pi^k\Gamma(1-s)}{\Gamma(s+k-1)} \qquad (7.13)\\
\times &\sum_j \frac{L_Q(s',\overline{u_j})\Gamma(s-\frac{1}{2}-it_j)\Gamma(s-\frac{1}{2}+it_j)\overline{\rho_j(-1)\langle V,u_j\rangle}}{\Gamma(\frac{1}{2}-it_j)\Gamma(\frac{1}{2}+it_j)}
\end{aligned}
$$

and

$$
\begin{aligned}
S_{\text{cont}} &= \frac{2^{2k-1}\pi^k\Gamma(1-s)}{4\pi\Gamma(s+k-1)}\sum_a\int_{-\infty}^{\infty}\zeta_{a,Q}(s',it)\\
\times \quad & \frac{\Gamma(s-\frac{1}{2}-it)\Gamma(s-\frac{1}{2}+it)\overline{\langle V,E_a(*,\frac{1}{2}+it)\rangle}}{\pi^{it}\zeta(1-2it)\Gamma(\frac{1}{2}-it)^2\Gamma(\frac{1}{2}+it)}dt.
\end{aligned}
$$

The functions $\zeta_{a,Q}(s',it)$ have the analogous definition to L_Q, with

$$
\zeta_{a,Q}(s',it) = Q^{-s'}\zeta_a(s'+it)\zeta_a(s'-it)\prod_{p^\gamma||Q}\left(\overline{\sigma_{2it}(p^\gamma)}-\overline{\sigma_{2it}(p^{\gamma-1})}p^{-s'}\right),
$$

and ζ_a differing from ζ at the factors corresponding to primes dividing N_0.

We have established so far that $Z_Q(s,w)$, as given by (7.12), is absolutely convergent (away from poles) when $\mathfrak{Re}\, s < 1/2 - k/2$ and $\mathfrak{Re}\, s' > 1$. We will now enlarge the region in which $Z_Q(s,w)$ is meromorphic to the region $\mathfrak{Re}\, w > 1$. This will be done by continuing past the line $\mathfrak{Re}\, s' = 1$ to all s'.

The objective is to find an upper bound for

$$
\begin{aligned}
&\sum_{|t_j|\sim T} L_Q(s',\overline{u_j})\overline{\rho_j(-1)\langle V,u_j\rangle}\\
&\qquad = \sum_{|t_j|\sim T} L(s',\overline{u_j})\overline{\rho_j(-1)\langle V,u_j\rangle}Q^{-s'}\prod_{p^\gamma||Q}\left(\overline{\lambda_j(p^\gamma)}-\overline{\lambda_j(p^{\gamma-1})}p^{-s'}\right)
\end{aligned}
\tag{7.14}
$$

Suppose that $\mathfrak{Re}\, s' < 0$. By the functional equation we have the bound for a single $L(s',\overline{u_j}$:

$$
L(s',\overline{u_j}) \ll (1+|s'|+|t_j|)^{1-2s'}(\ell_1\ell_2)^{1/2-s'}
$$

Also

$$
Q^{-s'}\prod_{p^\gamma||Q}\left(\overline{\lambda_j(p^\gamma)}-\overline{\lambda_j(p^{\gamma-1})}p^{-s'}\right) \ll Q^{\max(\theta-s',\mathfrak{Re}\,(-2s'))}
$$

Thus for $\mathfrak{Re}\, s' < 0$

$$
L_Q(s',\overline{u_j}) \ll Q^{\max(\theta-s',\mathfrak{Re}\,(-2s'))}(\ell_1\ell_2)^{1/2-s'}(1+|s'|+|t_j|)^{1-2s'}. \tag{7.15}
$$

For $\mathfrak{Re}\, s' \geq 1/2$ we can do better. By the approximate functional equation we may write $L(1/2+i\gamma',\overline{u_j})$ as a sum of length approximately

$$
N = \left(\ell_1\ell_2N_0(1+|t_j+\gamma'|)^2\right)^{1/2} \ll L((T+|\gamma'|),
$$

obtaining

$$\rho_j(-1)L(1/2+i\gamma',\overline{u_j}) = 2\sum_m \frac{\rho_j(m)}{|m|^{1/2+i\gamma'}} V\left(\frac{|m|}{N}\right).$$

(Recall u_j has trivial central character.) The large sieve estimate of Deshouillers and Iwaniec [7], Theorem 2 line (1.29) states that

$$\sum_{|t_j|\sim T} \left| e^{-\pi|t_j|} \sum_{m\sim N} a(m)\rho_j(m) \right|^2 << (qT^2 + N^{1+\epsilon})||a||$$

where $q = \ell_1\ell_2$ is the level of the u_j and $||a|| = \left(\sum a(m)\right)^{1/2}$. Thus taking

$$a(m) = \frac{V\left(\frac{|m|}{N}\right)}{|m|^{1/2+i\gamma}},$$

we obtain

$$\sum_{|t_j|\sim T} e^{-\pi|t_j|}|\rho_j(-1)|^2|L(1/2+i\gamma',\overline{u_j})|^2 \ll (1+|\gamma'|+T)^2L^2\log(LT(1+|\gamma'|)),$$

which is the Lindelöf bound on average.

For $\mathfrak{Re}\, s' \geq 1/2$ we have

$$\prod_{p^\gamma||Q}\left(\overline{\lambda_j(p^\gamma)} - \overline{\lambda_j(p^{\gamma-1})}p^{-s'}\right) \ll \lambda_j(Q) \ll Q^\theta.$$

This implies that for $\mathfrak{Re}\, s' \geq 1/2$

$$\sum_{|t_j|\sim T} e^{-\pi|t_j|}|\rho_j(-1)L_Q(1/2+i\gamma',\overline{u_j})|^2 \ll Q^{2\theta-1}L^2(1+|\gamma'|+T)^2L^2\log(LT(1+|\gamma'|)), \tag{7.16}$$

Alternatively, rather than bounding $\lambda_j(Q)$ individually from above, we apply Cauchy-Schwarz to the left hand side of (7.14) above. One piece of this is then

$$\sum_{|t_j|\sim T} e^{-\pi|t_j|}|\rho_j(-1)L_Q(s',\overline{u_j})|^2 \ll Q^{-2s'}\sum_{|t_j|\sim T}|\rho_j(-1)L(s',\overline{u_j})|^2\sum_{|t_j|\sim T}|\lambda_j(Q)|^2. \tag{7.17}$$

To estimate $\sum_{|t_j|\sim T}|\overline{\lambda_j(Q)}|^2$ we will use an application of the Kuznetsov formula, [19], Theorem 16.8, line (16.56)) as before, but first we have to normalize as follows:

$$\sum_{|t_j|\sim T}|\overline{\lambda_j(Q)}|^2 = \sum_{|t_j|\sim T}\left|\frac{\overline{\rho_j(Q)}}{\rho_j(1)}\right|^2 \ll L^\epsilon T^\epsilon \sum_{|t_j|\sim T}|\overline{\rho_j(Q)}|^2.$$

Here we have used the lower bound for $\rho_j(1)$ given in (4.4). Switching Q for h in (4.6) gives us

$$\sum_{|t_j|\sim T} |\overline{\rho_j(Q)}|^2 e^{-\pi|t_j|} \ll \ell_1\ell_2 T^2 + T^{7/4}\sigma_0(\ell_1\ell_2)(Q,\ell_1\ell_2)^{1/2}Q^{1/2}\log^2(3Q).$$

and combining the above two expressions leads to

$$\sum_{|t_j|\sim T} |\overline{\lambda_j(Q)}|^2 \ll L^{2+\epsilon}T^{2+\epsilon} + T^{7/4+\epsilon}\sigma_0(\ell_1\ell_2)(Q,\ell_1\ell_2)^{1/2}Q^{1/2+\epsilon}\log^2(Q).$$

Thus, for $\ell_1,\ell_2 \gg Q^{1/4}\log(Q)$, with $(Q,\ell_1\ell_2)=1$ and ℓ_1,ℓ_2 prime,

$$\sum_{|t_j|\sim T} |\overline{\lambda_j(Q)}|^2 \ll L^{2+\epsilon}T^{2+\epsilon}$$

and, in this case,

$$\sum_{|t_j|\sim T} |\rho_j(-1)L_Q(1/2+i\gamma',\overline{u_j})|^2 \ll Q^{-1}L^{2+\epsilon}(1+|\gamma'|+|T|)^{2+\epsilon}. \tag{7.18}$$

From Proposition 4.1 we have

$$\sum_{|t_j|\sim T} |\overline{\langle V,u_j\rangle}|^2 \ll L^{-2k}T^{2k}. \tag{7.19}$$

Combining (7.14), (7.18), and (7.19) finally gives us

$$\begin{aligned}\sum_{|t_j|\sim T} & L_Q(s',\overline{u_j})\overline{\rho_j(-1)\langle V,u_j\rangle} \\ &\ll \left(\sum_{|t_j|\sim T} |\rho_j(-1)L_Q(1/2+i\gamma',\overline{u_j})|^2\right)^{1/2}\left(\sum_{|t_j|\sim T} |\overline{\langle V,u_j\rangle}|^2\right)^{1/2} \\ &\ll Q^{-1/2}L^{1-k+\epsilon}(1+|\gamma'|+|T|)^{1+k+\epsilon}. \end{aligned} \tag{7.20}$$

To understand the region of absolute convergence we now rewrite (7.13) as

$$S_{\text{cusp}} = \lim_{T\to\infty} S_{\text{cusp}}(T),$$

with

$$\begin{aligned} S_{\text{cusp}}(T) = \frac{2^{2k-1}\pi^k\Gamma(1-s)}{\Gamma(s+k-1)} \\ \times \sum_{|t_j|\ll T} \frac{L_Q(s',\overline{u_j})\Gamma(s-\frac12-it_j)\Gamma(s-\frac12+it_j)\overline{\rho_j(-1)\langle V,u_j\rangle}}{\Gamma(\frac12-it_j)\Gamma(\frac12+it_j)} \end{aligned} \tag{7.21}$$

The sum $S_{\mathrm{cusp}}(T)$ can in turn be rewritten, after breaking up the interval $[0,T]$ diadically, as

$$S_{\mathrm{cusp}}(T) \ll \sum_{i=0}^{\infty} S'_{\mathrm{cusp}}(T/2^i)$$

with

$$S'_{\mathrm{cusp}}(T') = \frac{2^{2k-1}\pi^k \Gamma(1-s)}{\Gamma(s+k-1)} \tag{7.22}$$
$$\times \sum_{|t_j|\sim T'} \frac{L_Q(s',\overline{u_j})\Gamma(s-\frac{1}{2}-it_j)\Gamma(s-\frac{1}{2}+it_j)\overline{\rho_j(-1)\langle V,u_j\rangle}}{\Gamma(\frac{1}{2}-it_j)\Gamma(\frac{1}{2}+it_j)}$$

By Stirling's formula we have

$$\frac{\Gamma(1-s)\Gamma(s-\frac{1}{2}-it_j)\Gamma(s-\frac{1}{2}+it_j)}{\Gamma(s+k-1)\Gamma(\frac{1}{2}-it_j)\Gamma(\frac{1}{2}+it_j)} \ll (1+|t_j|)^{2s-2}(1+|s|)^{1-k}, \tag{7.23}$$

which is valid regardless of the relative sizes of t_j and $\mathfrak{Im}\, s$.

We now substitute the upper bounds of (7.15), (7.20) and (7.23), obtaining, for $\mathfrak{Re}\, s' < 1/2$,

$$(\ell_1\ell_2)^{(k-1)/2} S'_{\mathrm{cusp}}(T) \tag{7.24}$$
$$\ll \; Q^{\max(\theta-s',-2s')} L^{1-2s'}(1+|s'|+T)^{2-2s'} T^{2s+k-2}(1+|s|)^{1-k}\log(LT).$$

In the case $\mathfrak{Re}\, s' \geq 1/2$, we similarly get

$$(\ell_1\ell_2)^{(k-1)/2} S'_{\mathrm{cusp}}(T) \tag{7.25}$$
$$\ll \; Q^{\theta-s'} L^{\epsilon}(1+|s'|+T)^{1+\epsilon} T^{2s+k-2}(1+|s|)^{1-k}\log(LT).$$

The exponent θ can be replaced by 0 when $\ell_1,\ell_2 \gg Q^{1/4}\log(Q)$, with $(Q,\ell_1\ell_2)=1$ and ℓ_1,ℓ_2 prime.

The cuspidal part contributes poles at $s = 1/2 \pm it_j - r$ coming from the gamma factors $\Gamma(s-1/2\pm it_j)$. The corresponding $\Gamma(s-1/2\pm it)$ in the continuous contribution do not contribute poles, for the same reason as they do not in the continuous contribution to $D(s;h)$. (Also, the existence of a pole in s independent of w would imply the existence of a corresponding pole in the continuous spectrum piece of $D(s;h)$, and there are no such poles.)

Unlike their counterparts in the cuspidal contribution, however, the functions $\zeta_{a,Q}$ may contribute poles. The continuous contribution is

$$S_{\mathrm{cont}}(s,s') = \frac{2^{2k-1}\pi^k\Gamma(1-s)}{4\pi\Gamma(s+k-1)} \sum_a \int_{-\infty}^{\infty} Q^{-s'}\zeta_a(s'+it)\zeta_a(s'-it)$$
$$\times \prod_{p^\gamma||Q} \left(\overline{\sigma_{2it}(p^\gamma)} - \overline{\sigma_{2it}(p^{\gamma-1})}p^{-s'}\right)$$
$$\times \frac{\Gamma(s-\frac{1}{2}-it)\Gamma(s-\frac{1}{2}+it)\overline{\langle V, E(*,\frac{1}{2}+it)\rangle}}{\pi^{it}\zeta(1-2it)\Gamma(\frac{1}{2}-it)^2\Gamma(\frac{1}{2}+it)} dt.$$

Multiplying top and bottom by the normalizing factor $\zeta^*(1+2it)$, and replacing it by u,

$$S_{\mathrm{cont}}(s,s') = \frac{2^{2k-2}\pi^{k-1/2}\Gamma(1-s)}{2\pi i\Gamma(s+k-1)} \sum_a \int_{(0)} Q^{-s'}\zeta_a(s'+u)\zeta_a(s'-u)$$
$$\times \prod_{p^\gamma||Q} \left(\overline{\sigma_u(p^\gamma)} - \overline{\sigma_u(p^{\gamma-1})}p^{-s'}\right)$$
$$\times \frac{\Gamma(s-\frac{1}{2}-u)\Gamma(s-\frac{1}{2}+u)\overline{\langle V, E^*(*,\frac{1}{2}+u)\rangle}}{\zeta^*(1+2u)\zeta^*(1-2u)\Gamma(\frac{1}{2}-u)\Gamma(\frac{1}{2}+u)} du. \tag{7.26}$$

The function $S_{\mathrm{cont}}(s,s')$ is initially defined for $\mathfrak{Re}\, s' > 1$, and in this region $S_{\mathrm{cont}}(s,s')$ has no poles in s' or s. To meromorphically continue to the region $\mathfrak{Re}\, s' \leq 1$, choose c such that $\zeta(\beta+i\gamma) \neq 0$ for $0 \leq \beta \leq c/\log(1+|\gamma|)$. Fix $s' = \sigma + i\gamma$ to be in the region $0 < \sigma < c/\log(1+|\gamma|)$ and shift u to the contour $u(t) = c/\log(1+|t|)+it$. A pole of $\zeta(s'-u)$ is passed over at $u = s'-1$. The residue at this point is

$$\frac{2^{2k-2}\pi^{k-1/2}\Gamma(1-s)}{2\pi i\Gamma(s+k-1)}\kappa_a(s')Q^{-s'} \prod_{p^\gamma||Q} \left(\overline{\sigma_{s'-1}(p^\gamma)} - \overline{\sigma_{s'-1}(p^{\gamma-1})}p^{-s'}\right)$$
$$\times \frac{\Gamma(s+\frac{1}{2}-s')\Gamma(s-\frac{3}{2}+s')\overline{\langle V, E^*(*,s'-\frac{1}{2})\rangle}}{\pi^{1/2-s'}\zeta^*(3-2s')\Gamma(s'-\frac{1}{2})^2\Gamma(\frac{3}{2}-s')}, \tag{7.27}$$

where $\kappa_a(s')$ is a Dirichlet polynomial. The singularities of the integrand in s' are at $u = s'-1, 1-s'$ and after the shift both singularities are to the left of the contour. The function of s' can thus be continued to the region $1 - c/\log(1+|\gamma|) < \sigma \leq 1$ as both singularities remain to the left of the contour. Note that neither the residue nor the integral have a pole in s' on the line $\mathfrak{Re}\, s' = 1$ (and in fact the residue vanishes at $s' = 1$).

Now shift u back to $\mathfrak{Re}\, u = 0$, passing over the pole of $\zeta_a(s'+u)$ at $1-s'$. The residue here equals the residue in (7.27), and is added with the same sign, as the integral is moved in the opposite direction and $\zeta(s'-u)$ is replaced by $\zeta(s'+u)$. Thus $S_{\text{cont}}(s,s')$ has no poles in s' when $\mathfrak{Re} s' \geq 1$, and, when $\mathfrak{Re} s' < 1$, is given by

$$\begin{aligned} S_{\text{cont}}(s,s') = {} & \frac{2^{2k-2}\pi^{k-1/2}\Gamma(1-s)}{2\pi i\Gamma(s+k-1)} \sum_a \int_{(0)} Q^{-s'}\zeta_a(s'+u)\zeta_a(s'-u) \\ & \times \prod_{p^\gamma || Q} \left(\overline{\sigma_u(p^\gamma)} - \overline{\sigma_u(p^{\gamma-1})}p^{-s'}\right) \\ & \times \frac{\Gamma(s-\frac{1}{2}-u)\Gamma(s-\frac{1}{2}+u)\overline{\langle V, E^*(*,\frac{1}{2}+u)\rangle}}{\zeta^*(1+2u)\zeta^*(1-2u)\Gamma(\frac{1}{2}-u)\Gamma(\frac{1}{2}+u)} du \\ & + \frac{2^{2k-1}\pi^{k-1/2}\Gamma(1-s)}{2\pi i\Gamma(s+k-1)} \kappa_a(s') Q^{-s'} \prod_{p^\gamma || Q} \left(\overline{\sigma_{s'-1}(p^\gamma)} - \overline{\sigma_{s'-1}(p^{\gamma-1})}p^{-s'}\right) \\ & \times \frac{\Gamma(s+\frac{1}{2}-s')\Gamma(s-\frac{3}{2}+s')\overline{\langle V, E^*(*, s'-\frac{1}{2})\rangle}}{\pi^{1/2-s'}\zeta^*(3-2s')\Gamma(s'-\frac{1}{2})^2\Gamma(\frac{3}{2}-s')}. \end{aligned} \tag{7.28}$$

Recall that $s' = s + w + k/2 - 1$. The integral above is convergent for any s' when $\mathfrak{Re}\, w > 1/2$ and $\mathfrak{Re}\, s < 1 - k/2$, and has no poles in s, as remarked before. It thus has no poles in s' when $\mathfrak{Re}\, s' < 1$ and $\mathfrak{Re}\, w > 1/2$. The residual term has a potential pole when $s' = 1/2$ and $f = g$ and, for $\mathfrak{Re}\, s' < 1$, has possible poles contributed by $\Gamma(s+\frac{1}{2}-s')\Gamma(s-\frac{3}{2}+s')$. A polar line of $\Gamma(s+\frac{1}{2}-s')$ is of the form $s+1/2-s-w-k/2+1 = 3/2-w-k/2 = --r$, for some $r \geq 0$ and thus when $\mathfrak{Re}\, w > 1$ would extend into the region $\mathfrak{Re}\, s' > 1, \mathfrak{Re}\, s' < 1-k/2$ where $S_{\text{cont}}(s,s')$ is analytic. A polar line of $G(s-\frac{3}{2}+s')$ is of the form $s-3/2+s+w+k/2-1 = 2s+w-5/2+k/2 = -r$ and this also, for $\mathfrak{Re}\, s$ sufficiently negative, extends into the region of analyticity. This leaves $s' = 1/2$ when $f = g$ as the only possible polar line. It follows that $(s'-1/2)S_{\text{cont}}(s,s')$ has no poles in the region

$$\{(s,w) | \mathfrak{Re}\, w > 1/2, \mathfrak{Re}\, s < 1-k/2\}.$$

It is easily verified that, except at the polar points, the continuous spectrum contribution, defined in (7.14) to (7.12), satisfies the same upper bound as the cuspidal contribution. Details are omitted.

Absolute convergence of the expansion, given in (7.12), of $Z_Q(s,w)$ as $T \to \infty$, (away from the potential polar line at $s' = 1/2$), will thus occur in (7.24), (7.25) when the exponent of T is negative. This occurs for

$$\mathfrak{Re}\, s < \begin{cases} \mathfrak{Re}\, s' - k/2 & \text{if } \mathfrak{Re}\, s' < 1/2, \\ 1/2 - k/2 & \text{if } \mathfrak{Re}\, s' \geq 1/2, \end{cases} \tag{7.29}$$

It follows that $Z_Q(s,w)$ is absolutely convergent when $\mathfrak{Re}\, s < 1/2 - k/2$ and $\mathfrak{Re}\, w > 1$.

As $s' = w + s + k/2 - 1$, the bounds above can be translated into ones involving w. In particular,for s at least a distance ϵ from the potential polar lines $s' = 1/2$, and$s = 1/2 - r + it_j$, for any $r < 1/2 - c$ and any $j \geq 1$, the following bounds hold: For $\mathfrak{Re}\, s' < 1/2$ and $\mathfrak{Re}\, w > 1$

$$Z_Q(s,w) \ll Q^{\max(\theta - s', -2s')} L^{1-2s'} \log(L)(1+|s|)^{1-k}(1+|s'|)^{1-2s'}. \tag{7.30}$$

For $\mathfrak{Re}\, s' \geq 1/2$ and $\mathfrak{Re}\, w > 1$

$$Z_Q(s,w) \ll Q^{\theta - s'} L^{\epsilon}(1+|s|)^{1-k}(1+|s'|)^{1-2s'}. \tag{7.31}$$

The function $P_Q(s,w)$ converges for any $w \in \mathbb{C}$ when s is in the region $c < \mathfrak{Re}\, s < 2$ and is at least a distance ϵ from the points $s = 1/2 - r + it_j$. Using (7.10) and Cauchy-Schwarz to estimate $P_Q(s,w)$ from above we obtain, in this region,

$$\begin{aligned} P_Q(s,w) \quad &\ll \quad L^{k-1} \left(\sum_{|t_j| \sim 1, 0 \leq r < 1/2 - c} |c_{r,j}|^2 \right)^{1/2} \\ &\times \quad \left(\sum_{|t_j| \sim 1, 0 \leq r < 1/2 - c} |L_Q(w + (k-1)/2 - r + it_j, \overline{u_j})|^2 \right)^{1/2} \end{aligned}$$

As $r < 1/2 - c$,

$$\mathfrak{Re}\, (w + (k-1)/2 - r + it_j) \geq \mathfrak{Re}\, w + k/2 - 1 + c.$$

Combining the previous two lines with with (5.4) and (7.15), (7.16) it follows that in this region

$$P_Q(s,w) \ll B(w) L^{-1} L^{\epsilon}, \tag{7.32}$$

for any $B(w)$ satisfying

$$B(w) \gg \left(\sum_{|t_j| \sim 1} |L_Q(w + k/2 - 1 + c + it_j, \overline{u_j})|^2 \right)^{1/2}.$$

Thus, for example, for $\mathfrak{Re}\, w + k/2 - 1 + c \geq 1/2$

$$B(w) \ll Q^{\theta - 1/2} L^{1+\epsilon}. \tag{7.33}$$

This means that $P_Q(s,w)$ also satisfies the bounds of (7.30),(7.31), from which it follows that the difference $\tilde{Z}_Q(s,w)$ also satisfies these bounds, and $(s'-1/2)\tilde{Z}_Q(s,w)$ is analytic in the region $c < \mathfrak{Re}\, s < 2$, $\mathfrak{Re}\, w > 1$.

For $\mathfrak{Re}\, s, w > 1$, the function $Z_Q(s,w)$ converges absolutely. By (7.4) it satisfies

$$Z_Q(s,w) \ll (QL)^{-1-\epsilon} L^{-(k-1)/2}$$

in the region $\mathfrak{Re}\, s, w > 1$.

We have shown that $(s'-1/2)\tilde{Z}_Q(s,w)$ is analytic in the tube domain $\mathfrak{Re}\, w > 1$, $c-\epsilon < \mathfrak{Re}\, s < 2$ and the tube domain $\mathfrak{Re}\, w > 1$, $\mathfrak{Re}\, s > 1$. Consequently it is analytic in the convex closure of the union, namely the tube domain $\mathfrak{Re}\, w > 1$, $c-\epsilon < \mathfrak{Re}\, s$. This completes the proof of the proposition. □

Referring to (7.1) and (7.3), for $\mathfrak{Re}\, s, \mathfrak{Re}\, w > 1$

$$Z_Q(s,w) = \sum_{\substack{h_0, m_2 \geq 1 \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1}(h_0 Q)^{w+(k-1)/2}}, \tag{7.34}$$

We will now analytically continue $Z_Q(s,w)$ into the region $\mathfrak{Re}\, w \leq 1$ and $\mathfrak{Re}\,(w+(k-1)/2+s-1/2) > 3/2$ and obtain an upper bound for $Z_Q(s,w)$ in this region. The bound is given in the following

Proposition 7.2. *Let $s' = w+k/2+s-1$ and suppose that $\mathfrak{Re}\, s' > 3/2$, $\mathfrak{Re}\, s > 1$ and $\mathfrak{Re}\, w \leq 1$. Choose K such that $1+\epsilon > K + \mathfrak{Re}\, w > 1$. Suppose also that $\ell_1, \ell_2 \sim L \ll Q$. Then $Z_Q(s,w)$ has an analytic continuation into this region and can be expressed by*

$$Z_Q(s,w) = \frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell_2^{s+(k-1)/2}} \sum_{0 \leq j \leq K} \binom{-(w+(k-1)/2)}{j}$$
$$\times \left(\frac{-\ell_2}{\ell_1}\right)^j \left(\sum_{d|Q} S_5'(s,w,j;Q/d) - S_8(s,w,j;Q)\right) + \mathcal{O}\left(\frac{1}{LQ}\right).$$

Here

$$S_5'(s,w,j;Q)$$
$$= \frac{1}{\varphi(Q)} \sum_{\chi \pmod Q} L(s+(k-1)/2-j, f, \chi) L(w+j, g, \overline{\chi}) \chi(\ell_1)\overline{\chi(\ell_2)}$$

and

$$S_8(s,w,j) \quad = \quad \sum_{(m,\ell_1\ell_2)=1} \frac{A(m)\bar{B}(m)}{m^{s+w+(k-1)/2}} \sum_{(r|\ell_1\ell_2)^\infty} \frac{A(\ell r)\bar{B}(\ell r)}{r^{s+w+(k-1)/2}},$$

where

$$\ell = \frac{\ell_1\ell_2}{(\ell_1,\ell_2)^2}.$$

Remark 7.3. It's interesting that the pole at $s+w+k/2-1 = 1/2$ when $f = g$, although outside of the range we are considering, is nevertheless evident in the $S_8(s,w,j;Q)$ contribution, which is the Rankin-Selberg convolution of f with g up to factors dividing $\ell_1\ell_2$.

Proof. Write

$$Z_Q(s,w) = S_1(s,w) + S_2(s,w), \tag{7.35}$$

where

$$S_1(s,w) = (\ell_1\ell_2)^{(k-1)/2} \sum_{\substack{\ell_2 m_2 \gg h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1}(h_0 Q)^{w+(k-1)/2}}$$

and

$$S_2(s,w) = (\ell_1\ell_2)^{(k-1)/2} \sum_{\substack{\ell_2 m_2 \ll h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1}(h_0 Q)^{w+(k-1)/2}}.$$

It is easily verified that in the region $\mathfrak{Re}(s+w+k/2) > 5/2$

$$S_1(s,w) \ll \frac{1}{\ell_2 Q^{w+s+k/2-3/2}} \ll \frac{1}{\ell_2 Q}. \tag{7.36}$$

In this same region

$$S_2(s,w) = \frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell_2^{s+k-1}} \times \sum_{\substack{\ell_2 m_2 \ll h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1} m_1^{w+(k-1)/2}} \left(\frac{\ell_2 m_2 + h_0 Q}{\ell_1 h_0 Q}\right)^{w+(k-1)/2}. \tag{7.37}$$

As $\ell_2 m_2 \ll h_0 Q$ the following is absolutely convergent:

$$\begin{aligned}\left(\frac{\ell_2 m_2 + h_0 Q}{\ell_1 h_0 Q}\right)^{w+(k-1)/2} &= \left(1 - \frac{\ell_2 m_2}{\ell_2 m_2 + h_0 Q}\right)^{-(w+(k-1)/2)} \\ &= \sum_{j\ge 0} \binom{-(w+(k-1)/2)}{j} \frac{(-\ell_2 m_2)^j}{(\ell_2 m_2 + h_0 Q)^j} \\ &= \sum_{j\ge 0} \binom{-(w+(k-1)/2)}{j} \left(\frac{-\ell_2 m_2}{\ell_1 m_1}\right)^j. \end{aligned} \tag{7.38}$$

Recalling $K \approx 1 - \mathfrak{Re}\, w$, break the previous series into two parts and substitute this into (7.37). This leads to

$$S_2(s,w) = S_3(s,w) + S_4(s,w), \tag{7.39}$$

where

$$\begin{aligned} S_3(s,w) = \frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell_2^{s+(k-1)/2}} \sum_{0\le j\le K} \binom{-(w+(k-1)/2)}{j} \left(\frac{-\ell_2}{\ell_1}\right)^j \\ \times \sum_{\substack{\ell_2 m_2 \ll h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}} \end{aligned} \tag{7.40}$$

and

$$\begin{aligned} S_4(s,w) = \frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell_2^{s+(k-1)/2}} \sum_{j>K} \binom{-(w+(k-1)/2)}{j} \left(\frac{-\ell_2}{\ell_1}\right)^j \\ \times \sum_{\substack{\ell_2 m_2 \ll h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}} \end{aligned} \tag{7.41}$$

As $\ell_1 \sim \ell_2 \sim L$ it is easy to check that

$$\begin{aligned} S_4(s,w) \ll \frac{1}{L^{s+(k-1)/2}} \sum_{j>K} \binom{-(w+(k-1)/2)}{j} \\ \times \sum_{\substack{\ell_2 m_2 \ll h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{A(m_1)\bar{B}(m_2)}{m_2^{s+(k-1)/2-K} m_1^{w+K}} \left(\frac{m_2}{m_1}\right)^{j-K}, \end{aligned}$$

and as $m_1 \gg m_2$ it follows that for $\mathfrak{Re}(s+w+k/2) > 5/2$

$$S_4(s,w) \ll \frac{1}{L^{K+1}Q}. \tag{7.42}$$

Write

$$S_3(s,w) = \frac{1}{\ell_2^{s+(k-1)/2}} \sum_{0\le j\le K} \binom{-(w+(k-1)/2)}{j} \left(\frac{-\ell_2}{\ell_1}\right)^j S_3(s,w,j), \tag{7.43}$$

where

$$S_3(s,w,j) = \sum_{\substack{\ell_2 m_2 \ll h_0 Q \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}}. \tag{7.44}$$

For any fixed j, $0 \le j \le K$, break $S_3(s,w,j)$ up as follows:

$$S_3(s,w,j) = S_5(s,w,j) - S_6(s,w,j) - S_7(s,w,j) - S_8(s,w,j), \tag{7.45}$$

where

$$S_5(s,w,j) = \sum_{\substack{m_2 \geq 1, h_0 \in \mathbb{Z} \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}},$$

$$S_6(s,w,j) = \sum_{\substack{h_0, m_2 \geq 1, h_0 Q \ll \ell_2 m_2 \\ \ell_1 m_1 = \ell_2 m_2 + h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}},$$

$$S_7(s,w,j) = \sum_{\substack{h_0, m_2 \geq 1 \\ \ell_1 m_1 = \ell_2 m_2 - h_0 Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}},$$

and

$$S_8(s,w,j) = \sum_{\ell_1 m_1 = \ell_2 m_2} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}}.$$

It can be easily checked that

$$S_6(s,w,j), S_7(s,w,j) \ll \frac{L}{Q}.$$

The sum $S_5(s,w,j) = S_5(s,w,j);Q$ is simply

$$S_5(s,w,j;Q) = \sum_{\ell_1 m_1 \equiv \ell_2 m_2 \pmod{Q}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}}.$$

Define $S_5'(s,w,j;Q)$ to be the same sum, with the condition $(m_2,Q) = 1$ imposed. (Recall we have also assumed $(\ell_1\ell_2, Q) = 1$.) Thus

$$S_5'(s,w,j;Q) \quad = \sum_{\substack{\ell_1 m_1 \equiv \ell_2 m_2 \pmod{Q} \\ (m_2,Q)=1}} \frac{a(m_1)\bar{b}(m_2)}{m_2^{s+k-1-j} m_1^{w+(k-1)/2+j}} \tag{7.46}$$

The sum $S_5'(s,w,j;Q)$ analytically continues to

$$\begin{aligned} &S_5'(s,w,j;Q) \\ &= \frac{1}{\varphi(Q)} \sum_{\chi \pmod{Q}} L(s+(k-1)/2-j, f, \chi) L(w+j, g, \overline{\chi}) \chi(\ell_1) \overline{\chi(\ell_2)}. \end{aligned} \tag{7.47}$$

The sums $S_5'(s,w,j;Q), S_5(s,w,j;Q)$ are related by the simple sieve

$$S_5'(s,w,j;Q) = \sum_{d|Q} \mu(d) S_5(s,w,j;Q/d)$$

for the following reason: Suppose $m_2 = m_2'd$, with $d|Q$. Then the condition $\ell_1 m_1 \equiv \ell_2 m_2 \pmod{Q}$ implies that $m_1 = m_1'd$ also, and thus this is equivalent to the condition $\ell_1 m_1' \equiv \ell_2 m_2' \pmod{Q/d}$. By Moebius reciprocity we then have

$$S_5(s,w,j;Q) = \sum_{d|Q} S_5'(s,w,j;Q/d). \tag{7.48}$$

This provides an analytic continuation for $S_5(s,w,j;Q)$.

Finally, the sum $S_8(s,w,j)$ factors as

$$S_8(s,w,j) \quad = \quad \sum_{(m,\ell_1\ell_2)=1} \frac{A(m)\bar{B}(m)}{m^{s+w+(k-1)/2}} \sum_{(r|\ell_1\ell_2)^\infty} \frac{A(\ell r)\bar{B}(\ell r)}{m^{s+w+(k-1)/2}}, \tag{7.49}$$

where

$$\ell = \frac{\ell_1\ell_2}{(\ell_1,\ell_2)^2}.$$

The proposition follows after assembling the bounds and relations (7.35)-(7.49). □

It follows from the preceding two propositions that $\tilde{Z}_Q(s,w)(s+w+k/2-3/2)$ is analytic in the union of the regions $\mathfrak{Re}\, w > 1$ and $\mathfrak{Re}(s + w + k/2 > 5/2$. As the convex hull of this union is $\mathbb{C}^2$ the analytic continuation of $\tilde{Z}_Q(s,w)(s+w+k/2-3/2)$ and the consequent meromorphic continuation of $Z_Q(s,w)$ follows.

8. Double shifted sums

The analytic properties of $Z_Q(s,w)$ make it possible to derive some sharp estimates for double shifted sums. For example, take the case $Q = \ell_1 = \ell_2 = 1$ and consider

$$S(x,y) = \sum_{m_2,h\geq 1} \frac{A(m_2+h)\bar{B}(m_2)\left(1+\frac{h}{m_2}\right)^{(k-1)/2}}{h^{(k-1)/2}} G(m_2/x)G(h/y),$$

where $G(x)$ is smooth with compact support in the interval $[1,2]$. This sum will behave differently, depending on the relative sizes of y and x. Write $y = x^{1+\alpha}$. For $y \ll x$, that is, $\alpha < 0$, it essentially estimates

$$y^{-(k-1)/2} \sum_{m_2\sim x, h\sim y} A(m_2+h)\bar{B}(m_2),$$

while for $y \gg x$, that is, $\alpha > 0$ it essentially estimates

$$x^{-(k-1)/2} \sum_{m_2\sim x, h\sim y} A(m_2+h)\bar{B}(m_2).$$

A key word here is "essentially". The sum $S(x,y)$ is weighted by the factor

$$\frac{\left(1+\frac{h}{m_2}\right)^{(k-1)/2}}{h^{(k-1)/2}}$$

and it is precisely this factor which allows extremely sharp estimates to be made for $S(x,y)$. However, it is the unweighted estimates

$$\sum_{m_2\sim x, h\sim y} A(m_2+h)\bar{B}(m_2)$$

that are required for most applications.

The sum is related to $Z(s,w)=Z_1(s,w)$ by the inverse Mellin transform

$$S(x,y)=\left(\frac{1}{2\pi i}\right)^2\int_{(2)}\int_{(2)} Z(s,w)x^s y^w g(s)g(w)dsdw.$$

Here $g(s)$ is the Mellin transform of $G(x)$, as before. Referring to Proposition 7.1, move the s line of integration to the left, to $\mathfrak{Re}\, s = 1/2-k_2-\epsilon$, passing over simple poles at $s=1/2-r+it_j$ for $0\le r\le k/2$. The residue at $s=1/2-r+it_j$ is

$$\frac{c_{r,j}x^{1/2-r+it_j}g(1/2-r+it_j)}{2\pi i}\int_{(2)} L(w+(k-1)/2-r+it_j;\overline{u_j})y^w g(w)dw.$$

As the integrand has no poles, the line of integration may be moved arbitrarily far to the left, to $\mathfrak{Re} w=-C$, and, summing over j and r, the residue contribution is

$$\mathcal{O}\left(x^{1/2}y^{-C}\right).$$

Thus

$$\begin{aligned} &S(x,y)\\ &=\left(\frac{1}{2\pi i}\right)^2\int_{(2)}\int_{(1/2-k/2-\epsilon)} Z(s,w)x^s y^w g(s)g(w)dsdw+\mathcal{O}\left(x^{1/2}y^{-C}\right).\end{aligned}$$

Now move the w line of integration to the left in this double integral. When $f=g$ there is a pole at $s+w+k/2=3/2$, and when $\mathfrak{Re}\, s = 1/2-k/2-\epsilon$, this occurs on the line $\mathfrak{Re}\, w = 1+\epsilon$. Estimating the integral at his point, we get

$$S(x,y)\ll x^{1/2-k/2-\epsilon}y^{1+\epsilon}.$$

On the other hand

$$S(x,y)\approx\sum_{m_2\sim x, h\sim y}\frac{A(m_2+h)\bar{A}(m_2)\left(1+\frac{y}{x}\right)^{(k-1)/2}}{y^{(k-1)/2}}$$

and for $y \gg x$ the trivial estimate for this sum is $x^{1/2-k/2}xy$. Thus the estimate is non-trivial as long as $y^{1+\epsilon}x^{-\epsilon} \ll xy$, i.e as long as

$$y \ll x^{(1+\epsilon)/\epsilon}.$$

8.1. **Removing the weighting factor.**

The unweighted but smoothed sum we are interested in estimating is

$$S_Q(x,y) = \sum_{\substack{\ell_1 m_1 = \ell_2 m2 + hQ \\ h, m_2 \geq 1}} A(m_1)\bar{B}(m_2)G_1(m_1/y)G_2(m_2/x).$$

The sum $S_Q(x,y)$ is related to $Z_Q(s,w)$ via a triple inverse Mellin transform, as set forth in the following

Proposition 8.1. *Suppose $G_1(x), G_2(x)$ are smooth with compact support in the interval $[1,2]$, with $g_1(s), g_2(s)$ denoting the corresponding Mellin transforms as before. For $x \gg 1$,*

$$\begin{aligned} S_Q(x,y) = \left(\frac{1}{2\pi i}\right)^3 \int_{(\gamma_1)}\int_{(\gamma_2)}\int_{(\gamma_3)} \frac{Z_Q(s+w-u, u+(1-k)/2)}{\Gamma((k-1)/2+w)} \\ \times \Gamma((k-1)/2+w-u)\Gamma(u)\ell_1^w \ell_2^s x^s y^w g_1(w) g_2(s)\, du\, dw\, ds. \quad (8.1) \end{aligned}$$

Here $\gamma_1, \gamma_2, \gamma_3$ are chosen so that $\mathfrak{Re}\, s+w-u > 1$, $\mathfrak{Re}\, u+(1-k)/2) > 1$, and $\mathfrak{Re}\, (k-1)/2+w-u > 0$ and in particular we choose $\gamma_1 = (k+1)/2$, $\gamma_2 = 1+2\epsilon$, $\gamma_3 = (k+1)/2+\epsilon$, for any $\epsilon > 0$.

Proof. The following identity can be found in [14], 6.422 #3, (after changing the signs of γ and u). For $\mathfrak{Re}\, \beta > \gamma > 0$ and $\arg(t) < \pi$,

$$\frac{1}{2\pi i}\int_{(\gamma)} \Gamma(u)\Gamma(\beta-u)t^{-u}du = \Gamma(\beta)(1+t)^{-\beta}.$$

Changing the variables in (7.4), and setting $\beta = (k-1)/2 + w$,

$$\frac{1}{2\pi i}\int_{(\gamma)} Z_Q(s+w-u, u$$
$$+ (1-k)/2)\Gamma(u)\Gamma((k-1)/2+w-u)\ell_1^w\ell_2^s du$$
$$= \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} A(m_1)\bar{B}(m_2)$$
$$\times \frac{1}{2\pi i}\int_{(\gamma)} \frac{\Gamma(u)\Gamma((k-1)/2+w-u)\ell_1^w\ell_2^s\left(1+\frac{h_0 Q}{\ell_2 m_2}\right)^{(k-1)/2} du}{(\ell_2 m_2)^{s+w-u}(h_0 Q)^u}$$
$$= \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \frac{A(m_1)\bar{B}(m_2)}{(\ell_2 m_2)^{s+w}}\Gamma((k-1)/2+w)$$
$$\times \left(1+\frac{h_0 Q}{\ell_2 m_2}\right)^{-w-(k-1)/2}\left(1+\frac{h_0 Q}{\ell_2 m_2}\right)^{(k-1)/2}\ell_1^w\ell_2^s du$$
$$= \Gamma((k-1)/2+w)\sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \frac{A(m_1)\bar{B}(m_2)\ell_1^w\ell_2^s}{(\ell_2 m_2)^{s+w}\left(1+\frac{h_0 Q}{\ell_2 m_2}\right)^w}$$
$$= \Gamma((k-1)/2+w)\sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \frac{A(m_1)\bar{B}(m_2)}{m_2^s m_1^w}. \quad (8.2)$$

The proposition then follows, after setting $\gamma = \gamma_3$ and applying the inverse Mellin transform

$$\left(\frac{1}{2\pi i}\right)^2\int_{(\gamma_1)}\int_{(\gamma_2)} g_1(w)g_2(s)x^s y^w dw ds$$

to both sides. ☐

Having related $S_Q(x,y)$ to $Z_Q(s+w-u, u+(1-k)/2)$ we will now derive an estimate for $S_Q(x,y)$ from the analytic properties of $Z_Q(s+w-u, u+(1-k)/2)$. We will show the following

Proposition 8.2. *Suppose $y \gg x \gg 1$, $\ell_1, \ell_2 \sim L \ll Q^{1/2}$. Then*

$$S_Q(x,y) \ll Q^{\theta-1/2}L^{1+\epsilon}y(y/x)^{\epsilon}.$$

If, in addition, $L \gg Q^{1/4}\log(Q)$ and $(\ell_1\ell_2, Q) = 1$ then

$$S_Q(x,y) \ll Q^{-1/2}L^{1+\epsilon}y(y/x)^{\epsilon}.$$

Proof. Let I denote the triple integral on the right hand side of (8.1). Bring the s integral inside and move the s line of integration to $\mathfrak{Re}\, s =$

$-\epsilon-\delta$, where $0 \leq \delta < 1$ will be specified shortly. After this, $\mathfrak{Re}\, s + w - u = 1/2 - k/2 - \delta$. As $\mathfrak{Re}\, u + (1-k)/2 = 1+\epsilon$, Proposition 7.1 applies, in particular the bound given in (7.5). While moving the line of integration poles are passed over at the points $s+w-u = 1/2-r+it_j$, for $0 \leq r \leq k/2$ (the upper bound of $k/2$ is there as $\delta < 1$. By Proposition 7.1, the residue at $s+w-u = 1/2-r+it_j$ of the triple integral is given by

$$\left(\frac{1}{2\pi i}\right)^2 \int_{(\gamma_2)} \int_{(\gamma_3)} \frac{(\ell_1\ell_2)^{(k-1)/2} c_{r,j} L_Q(u-r+it_j, \overline{u_j})}{\Gamma((k-1)/2+w)} \times \Gamma((k-1)/2+w-u)\Gamma(u)(\ell_1 y)^w (\ell_2 x)^{u-w+1/2-r+it_j} \times g_1(w) g_2(u-w+1/2-r+it_j) du dw. \quad (8.3)$$

Now move the u line of integration of this double integral to $\mathfrak{Re}\, u = r+1/2$. No poles are passed over in the process. The argument of $\Gamma((k-1)/2+w-u)$ now satisfies

$$\mathfrak{Re}\,(k-1)/2+w-u = k/2-r+2\epsilon > 0.$$

Move the w line of integration inside, and then to $\mathfrak{Re}\,(k-1)/2+w-u = -1/2$, i.e to $\mathfrak{Re}\, w = 1/2+r-k/2$. In the process a pole is passed over at $w = u+(1-k)/2$. The residue at $w = u+(1-k)/2$ is the single integral

$$R_1(r,j) = \frac{1}{2\pi i} \int_{(0)} (\ell_1\ell_2)^{(k-1)/2} c_{r,j} L_Q(1/2+it_j+it, \overline{u_j})(\ell_1 y)^{r+1-k/2+it} \times (\ell_2 x)^{k/2-r+it_j} g_1(r+1-k/2+it) g_2(k/2-r+it_j) du, \quad (8.4)$$

where we have written $u = 1/2+it$.

After the lines have been shifted, as above, we have

$$I = \sum_{j,r,0\leq r\leq k/2} R_1(r,j) + \sum_{j,r,0\leq r\leq k/2} R_2(r,j) + \left(\frac{1}{2\pi i}\right)^3 \int_{(\gamma_2)} \int_{(\gamma_3)} \int_{(-\epsilon-\delta)} \frac{Z_Q(s+w-u, u+(1-k)/2)}{\Gamma((k-1)/2+w)} \times \Gamma((k-1)/2+w-u)\Gamma(u)\ell_1^w \ell_2^s x^s y^w g_1(w) g_2(s) ds du dw, \quad (8.5)$$

with $R_1(r,j)$ given in (8.4) and

$$R_2(r,j) = \left(\frac{1}{2\pi i}\right)^2 \int_{(r+1/2)} \int_{(1/2+r-k/2)} \frac{(\ell_1\ell_2)^{(k-1)/2} c_{r,j}}{\Gamma((k-1)/2+w)} \times L_Q(u-r+it_j, \overline{u_j})\Gamma((k-1)/2+w-u)\Gamma(u)(\ell_1 y)^w (\ell_2 x)^{u-w+1/2-r+it_j} \times g_1(w) g_2(u-w+1/2-r+it_j) dw du. \quad (8.6)$$

To estimate the sums over $R_1(r,j)$ and $R_2(r,j)$ we first recall via (5.4) and (7.18) that, for fixed r,

$$\begin{aligned}
&\sum_{|t_j|\sim T} (\ell_1\ell_2)^{(k-1)/2} c_{r,j} L_Q(1/2+it_j+it,\overline{u_j}) \\
&\qquad \ll L^{k-1}\left(\sum_{|t_j|\sim T}|c_{r,j}|^2\right)^{1/2}\left(\sum_{|t_j|\sim T}|L_Q(1/2+it_j+it,\overline{u_j})|^2\right)^{1/2} \\
&\qquad\qquad \ll L^{k-1}\left(\log(\ell_1\ell_2T)(\ell_1\ell_2)^{-k}T^{2r+1}\right)^{1/2} \\
&\qquad\qquad\quad \times \left(Q^{2\theta-1}L^{2+\epsilon}(1+|t_j+t|+|T|)^{2+\epsilon}\right)^{1/2} \\
&\qquad\qquad \ll L^{k-1-k+1+\epsilon}(1+|t_j+t|+|T|)^{1+\epsilon}Q^{\theta-1/2} \\
&\qquad\qquad\qquad \ll L^{\epsilon}(1+|t_j+t|+|T|)^{1+\epsilon}Q^{\theta-1/2}. \qquad (8.7)
\end{aligned}$$

Here $\theta = 0$ if $L \gg Q^{1/4}\log(Q)$ and $(\ell_1\ell_2, Q) = 1$.

The constraint $\ell_1 m_1 = \ell_2 m2 + hQ$, with $m_2, h \geq 1$ implies that the sum $S_Q(x,y)$ is trivial unless $y \gg x$. In the case of $R_1(r,j)$

$$(\ell_1 y)^{r+1-k/2+it}(\ell_2 x)^{k/2-r+it_j} \ll (Ly)(y/x)^{r-k/2} \ll Ly.$$

In the case of $R_2(r,j)$,

$$(\ell_1 y)^{w}(\ell_2 x)^{u-w+1/2-r+it_j} \ll (y/x)^{1/2+r-k/2}(Lx) \ll (y/x)^{1/2}(Lx) \ll Ly.$$

Substituting these bounds and (8.7) into (8.4), (8.6), and using the rapid decay of $\Gamma(u), g_1(w), g_2(s)$, we see that

$$\sum_{j,r,0\leq r\leq k/2} R_1(r,j) + \sum_{j,r,0\leq r\leq k/2} R_2(r,j) \ll L^{1+\epsilon} y Q^{\theta-1/2}. \qquad (8.8)$$

Turning to the shifted triple integral, we recall Proposition 7.1, in particular (7.5), which we restate for convenience: For $\mathfrak{Re}\, s' = \mathfrak{Re}\, s + w + k/2 - 1 \geq 1/2$ and $\mathfrak{Re}\, w > 1$

$$Z_Q(s,w) \ll Q^{\theta-s'}L^{\epsilon}(1+|s|)^{1-k}(1+|s'|)^{1-2s'}. \qquad (8.9)$$

In our integral $s' = s + w - 1/2$ and

$$\mathfrak{Re}\, s' = -\epsilon - \delta + \gamma_2 - 1/2 = 1/2 + \epsilon - \delta.$$

Setting $\delta = \epsilon$ we obtain $\mathfrak{Re}\, s' = 1/2$, which is optimal.

There is a potential pole at $s' = 1/2$ which is described in (7.28). Translating into the present parameters, the piece containing the pole is

$$\begin{aligned}&\frac{2^{2k-1}\pi^{k-1/2}\Gamma(1+u-s-w)}{2\pi i\Gamma(s+w-u+k-1)}\kappa_a(s+w-1/2)Q^{1/2-s-w}\\ &\qquad\times\prod_{p^\gamma||Q}\left(\overline{\sigma_{s+w-3/2}(p^\gamma)}-\overline{\sigma_{s+w-3/2}(p^{\gamma-1})}p^{1/2-s-w}\right)\\ &\qquad\times\frac{\Gamma(1-u)\Gamma(2s+2w-2-u)\overline{\langle V,E^*(*,s+w-1)\rangle}}{\pi^{1-s-w}\zeta^*(4-2s-2w)\Gamma(s+w-1)^2\Gamma(2-s-w)}. \quad (8.10)\end{aligned}$$

This is analytic, and in fact vanishes, at $s+w-1/2=1/2$, as $\mathfrak{Re}\, u=(k+1)/2$, which is not a positive integer. Thus in this context there is no pole on the line $\mathfrak{Re}\, s'=1/2$.

Writing $s=-\epsilon-\delta+it_s$, $w=1+2\epsilon+it_w$ and $u=(k+1)/2+\epsilon+it_u$ we have $s'=1/2+i(t_s+t_w)$. The bound of (8.9) translates to

$$Z_Q(s+w-u,u+(1-k)/2)\ll Q^{\theta-1/2}L^\epsilon(1+|t_s+t_w-t_u|)^{1-k}.$$

Also, in this region

$$(\ell_1 y)^w(\ell_2 x)^s\ll(Ly)^{1+2\epsilon}(Lx)^{-\epsilon-\epsilon}=(Ly)(y/x)^{2\epsilon}.$$

Substituting the two previous estimates into the triple integral appearing in (8.5), and using the rapid decay of $\Gamma(u)g_1(w), g_2(s)$, the integral is bounded above by

$$Q^{\theta-1/2}L^{1+\epsilon}y(y/x)^{2\epsilon}.$$

Combining this with the bound (8.8), and substituting into (8.5), we finally have

$$I\ll Q^{\theta-1/2}L^{1+\epsilon}y(y/x)^{2\epsilon},$$

with $\theta=0$ if $L\gg Q^{1/4}\log(Q)$ and $(\ell_1\ell_2,Q)=1$. Replacing ϵ by $\epsilon/2$ completes the proof of the Proposition. □

9. APPLICATION TO SUBCONVEXITY

Our amplification argument is modeled on that of Blomer in [2], which is in turn based on that of [10]. Let χ be a primitive character mod Q with $(Q,N_0)=1$, and

$$L(s,f\otimes\chi)=\sum_{m\geq 1}\frac{A(m)\chi(m)}{m^s}.$$

Write $L(1/2,f\otimes\chi)$ using the approximate functional equation:

$$L(1/2,f\otimes\chi)=\sum_m\frac{A(m)\chi(m)}{\sqrt{m}}V\left(\frac{m}{Q\sqrt{N_0}}\right)+\omega\sum_m\frac{\tilde{A}(m)\chi(m)}{\sqrt{m}}V\left(\frac{m}{Q\sqrt{N_0}}\right).$$

Let

$$B_\chi(x) = \sum_m A(m)\chi(m)\ G\left(\frac{m}{x}\right),$$

where G is smooth of compact support in $[1,2]$ and x is a parameter. By a smooth dyadic partition of unity, and summation by parts, we have

$$L(1/2, f\otimes\chi) \ll \max_{x\ll Q}(Q\sqrt{N_0})^{-1/2}B_\chi(x).$$

Suppressing the N_0 dependence, we have

$$L(1/2, f\otimes\chi) \ll \max_{x\ll Q} Q^{-1/2}B_\chi(x). \tag{9.1}$$

Set $L = Q^{1/4}\log(Q)$, and for any $x \ll Q$, consider the amplified sum

$$S = \sum_{\psi(Q)}\left|\sum_m A(m)\psi(m)G\left(\frac{m}{x}\right)\right|^2\left|\sum_{\ell\sim L, \ell\,\mathrm{prime},\,(\ell,N_0)=1}\bar{\chi}(\ell)\psi(\ell)\right|^2. \tag{9.2}$$

Notice that the term with $\psi = \chi$ contributes

$$|B_\chi(x)|^2\left|\sum_{\ell\sim L,\ell\,\mathrm{prime},\,(\ell,N_0)=1} 1\right|^2.$$

The number of $\ell \sim L$ with ℓ prime is

$$\sim L/\log(L) \sim \frac{Q^{1/4}\log(Q)}{\log(Q^{1/4}\log(Q)} \sim Q^{1/4},$$

so taking only the contribution to S from $\psi = \chi$ gives

$$B_\chi(x) \ll S^{1/2}Q^{-1/4}. \tag{9.3}$$

Combining this with (9.1) we now have

$$L(1/2, f\otimes\chi) \ll \max_{x\ll Q} Q^{-3/4}S^{1/2}. \tag{9.4}$$

Use Parseval (for the group (Z/QZ)*) on the sum (9.2) to obtain

$$S = \sum_{a \bmod Q,(a,Q)=1}\left|\sum_{m\equiv a \bmod Q} A(m)\psi(m)G\left(\frac{m}{x}\right)\right|^2$$

$$\leq \sum_{a \bmod Q}\left|\sum_{m\equiv a \bmod Q} A(m)\psi(m)G\left(\frac{m}{x}\right)\right|^2.$$

Opening this sum and using the orthogonality of characters gives

$$\begin{aligned} S \leq \sum_{m_1}\sum_{m_2}\sum_{\ell_1}\sum_{\ell_2}\sum_{\psi} & A(m_1)\bar{A}(m_2) \\ & \times \psi(m_1)\bar{\psi}(m_2) G\left(\frac{m_1}{x}\right) G\left(\frac{m_2}{x}\right) \bar{\chi}(\ell_1)\chi(\ell_2)\psi(\ell_1)\bar{\psi}(\ell_2) \\ & = \phi(Q) \sum_{\ell_1}\sum_{\ell_2} \bar{\chi}(\ell_1)\chi(\ell_2) \\ & \times \left(\sum_{m_1} \sum_{\substack{m_2 \\ \ell_1 m_1 \equiv \ell_2 m_2 \bmod Q}} A(m_1)\bar{A}(m_2) G\left(\frac{m_1}{x}\right) G\left(\frac{m_2}{x}\right) \right). \end{aligned}$$

The sum over m_1, m_2 such that $\ell_1 m_1 \equiv \ell_2 m_2 \pmod{Q}$ can be broken into three pieces:

$$S \ll \phi(Q) \sum_{\ell_1}\sum_{\ell_2} \bar{\chi}(\ell_1)\chi(\ell_2)(S_1 + S_2 + S_3), \tag{9.5}$$

where

$$S_1 = \sum_{m_2 \geq 1, \ell_1 m_1 = \ell_2 m_2} A(m_1)\bar{A}(m_2) G\left(\frac{m_1}{x}\right) G\left(\frac{m_2}{x}\right),$$

$$S_2 = \sum_{m_2, h_0 \geq 1, \ell_1 m_1 = \ell_2 m_2 + h_0 Q} A(m_1)\bar{A}(m_2) G\left(\frac{m_1}{x}\right) G\left(\frac{m_2}{x}\right),$$

and

$$S_3 = \sum_{m_2, h_0 \geq 1, \ell_2 m_1 = \ell_1 m_2 + h_0 Q} A(m_1)\bar{A}(m_2) G\left(\frac{m_1}{x}\right) G\left(\frac{m_2}{x}\right).$$

It is easily verified that for general ℓ

$$\phi(Q) \sum_{\ell_1}\sum_{\ell_2} \bar{\chi}(\ell_1)\chi(\ell_2) S_1 \ll L Q^{2+\epsilon}.$$

For prime ℓ_1, ℓ_2 the Q^ϵ can be removed as follows. The condition $\ell_1 m_1 = \ell_2 m_2$ reduces to the two cases $\ell_1 = \ell_2$ and $m_1 = m_2$, or $(\ell_1, \ell_2) = 1$, with $m_1 = \ell_2 m$, $m_2 = \ell_1 m$. The first case occurs with multiplicity $Q^{1/4}$ and the contribution to S_1 is

$$\sum_{m \geq 1} A(m)\bar{A}(m) G\left(\frac{m}{x}\right) G\left(\frac{m}{x}\right) \ll x.$$

The second case occurs with multiplicity L^2, and the contribution to S_1 is

$$\sum_{m \geq 1} A(\ell_2 m)\bar{A}(\ell_1 m) G\left(\frac{\ell_1 m}{x}\right) G\left(\frac{\ell_2 m}{x}\right).$$

This is equal to the following double inverse Mellin transform:

$$\left(\frac{1}{2\pi i}\right)^2 \int_{(2)}\int_{(2)} L(s+w;\ell_1,\ell_2)(x/\ell_1)^s(x/\ell_2)^w g(s)g(w)dsdw,$$

where

$$L(u;\ell_1,\ell_2) = \sum_{m\geq 1}\frac{A(\ell_1 m)A(\ell_2 m)}{m^u}.$$

Setting $u = s+w$, this becomes

$$\left(\frac{1}{2\pi i}\right)^2 \int_{(2)}\int_{(4)} L(u;\ell_1,\ell_2)(x/\ell_1)^s(x/\ell_2)^{u-s} g(s)g(u-s)duds. \quad (9.6)$$

Now

$$L(u;\ell_1,\ell_2) = L(u, f\otimes \bar{f})E_{\ell_1}(u)E_{\ell_1}(u)\sum_{\gamma\geq 0}\frac{A(\ell_1^{\gamma+1})\bar{A}(\ell_1^{\gamma})}{\ell_1^{\gamma u}} \times \sum_{\gamma\geq 0}\frac{A(\ell_2^{\gamma+2})\bar{A}(\ell_2^{\gamma})}{\ell_2^{\gamma u}},$$

where, for $p = \ell_1, \ell_2$,

$$E_p(u) = (1-\alpha_p^2 p^{-u})(1-\alpha_p^{-2}p^{-u})(1-p^{-u})(1+p^{-u})^{-1}$$

is the inverse of the Euler factor of $L(u, f\otimes\bar{f})$ at the prime p. In other words, the correct factors of the Rankin-Selberg convolution have been removed and replaced by altered factors.

The key point is that $L(u;\ell_1,\ell_2)$ continues to a function of u that is analytic when $\mathfrak{Re}\, u > 0$, except for a simple pole at $u = 1$ of residue κ, with $\kappa \ll 1$. The implied constant here depends, as usual, on f, (in particular on $L(1, f, \vee^2)$), but is independent of ℓ_1, ℓ_2. The Euler products and infinite sums are bounded from above by absolute constants.

Remark 9.1. If ℓ_1, ℓ_2 were allowed to vary over any integers, the length of this product would depend on the number of prime factors of $\ell_1\ell_2$. This would have introduced an extra L^ϵ. As they are restricted to primes the product is bounded absolutely.

Moving the u line of integration in (9.6) to $\mathfrak{Re}\, u = 1/2$ and passing over the pole at $u = 1$ gives us the relation

$$\sum_{m\geq 1} A(\ell_2 m)\bar{A}(\ell_1 m) G\left(\frac{\ell_1 m}{x}\right) G\left(\frac{\ell_2 m}{x}\right) = \frac{\kappa x}{\ell_2}\left(\frac{1}{2\pi i}\right)\int_{(2)} (\ell_2/\ell_1)^s (g(s)g(1-s)ds + \mathcal{O}\left(x^{1/2}\right) \ll \frac{x}{L}.$$

The total contribution of the two cases to the S_1 piece is thus

$$\phi(Q)\sum_{\ell_1}\sum_{\ell_2}\bar{\chi}(\ell_1)\chi(\ell_2)(S_1) \ll Q(Lx + L^2(x/L)) \ll QLx \ll Q^2L.$$

The sums S_2, S_3 are bounded using Proposition 8.2, with $x = y \ll Q$. As $L \sim Q^{1/4}$ and $(\ell_1\ell_2, Q) = 1$ this gives us

$$S_2, S_3 \ll Q^{-1/2}xL^{1+\epsilon} \ll Q^{1/2}L^{1+\epsilon}.$$

Substituting, we have for their contribution:

$$\phi(Q)\sum_{\ell_1}\sum_{\ell_2}\bar{\chi}(\ell_1)\chi(\ell_2)(S_2 + S_3) \ll QL^2Q^{1/2}L^{1+\epsilon} \ll Q^{3/2}L^{3+\epsilon}.$$

Referring to (9.5) we have

$$\begin{aligned} S \ll \phi(Q)\sum_{\ell_1}\sum_{\ell_2}\bar{\chi}(\ell_1)\chi(\ell_2)(S_1 + S_2 + S_3) &\ll Q^2L + Q^{3/2}L^{3+\epsilon} \\ &\ll Q^{9/4}\log(Q) + Q^{9/4+\epsilon} \ll Q^{9/4+\epsilon}. \end{aligned}$$

Finally, referring to (9.4),

$$L(1/2, f\otimes\chi) \ll \max_{x\ll Q} Q^{-3/4}S^{1/2} \ll Q^{3/8+\epsilon}.$$

This completes the proof of Theorem 1.1

10. AN APPLICATION TO MEAN VALUES OF L-SERIES

The object of this section is to use the techniques developed so far to prove the following mean value result. We will need to sharpen our notation now, and write

$$Z_Q(s, w; f, g) = Z_Q(s, w)$$

to include a reference to f and g. We will continue to suppress ℓ_1, ℓ_2 in the notation, but in this section we will take $\ell_1 = \ell_2 = 1$. Let

$$S(Q) = \frac{1}{\varphi(Q)}\sum_{\chi \bmod Q} L(1/2, f, \chi)\overline{L(1/2, g, \chi)}. \tag{10.1}$$

Also, if f, g are our usual cusp forms, with normalized Fourier coefficients $A(n), B(n)$, let $F_d(z), G_d(z)$ be defined by

$$F_d(z) = \sum_{n\geq 0} A(dn)(dn)^{(k-1)/2}e^{2\pi inz},\ G_d(z) = \sum_{n\geq 0} B(dn)(dn)^{(k-1)/2}e^{2\pi inz},$$

which are of level dN_0. Then

Proposition 10.1. *When* $f \neq g$

$$\begin{aligned} S(Q) &= L_Q(1, f \otimes g) \\ &+ \sum_{d|Q} \frac{\mu(d)}{d} \left(Z_{Q/d}(1-k/2, 1/2; F_d, G_d) + Z_{Q/d}(1-k/2, 1/2; G_d, F_d) \right) \\ &\qquad + \mathcal{O}\left(Q^{\theta - 1/2}\right), \end{aligned} \tag{10.2}$$

and when $f = g$*, for any* $x > 1$

$$\begin{aligned} S(Q) &= \operatorname{Res}_{s=0} L_Q(s+1, f \otimes f)\Gamma(s)(x/2)^s \\ &+2 \sum_{d|Q} \frac{\mu(d) \operatorname{Res}_{w=0} \Gamma(w) x^w Z_Q(1-k/2, w+1/2; F_d, F_d))}{d} + \mathcal{O}\left(Q^{\theta-1/2}\right). \end{aligned} \tag{10.3}$$

Remark 10.2. The interest in this Proposition is in the relation of the mean value to special values of the shifted multiple Dirichlet series $Z_Q(s, w; f, g)$. These special values will be shown to be constant on average in Section 12. The final mean value result will be derived by averaging the result of this Proposition over Q in short intervals.

For any $x \gg 1$, and $\mathfrak{Re}\, s > 1/2$, the following series is absolutely convergent.

$$L(s+1/2, f, \chi) = \sum_{m \geq 1} \frac{A(m)\chi(m)}{m^{s+1/2}}.$$

Multiplying by $\Gamma(s)$ and applying an inverse Mellin transform, one obtains

$$I = \int_{(2)} L(s+1/2, f, \chi) x^s \Gamma(s) ds = \sum_{m \geq 1} \frac{A(m)\chi(m)}{m^{1/2}} e^{-m/x}.$$

Fixing any $\epsilon > 0$, if $x \gg Q^{2+\epsilon}$ then by moving the line of integration to $\mathfrak{Re}\, s = -1/2 - C_1$, $C_1 > 0$, and using the functional equation to bound the L-series one obtains

$$I = L(1/2, f, \chi) + \mathcal{O}(x^{-1/2-C_1} Q^{1+2C_1}) = L(1/2, f, \chi) + \mathcal{O}(x^{-C_2})$$

for arbitrarily large C_2. Note that the character χ does *not* need to be primitive for this to be true.

Doing the same thing for g and multiplying gives us

$$\begin{aligned} &L(1/2, f, \chi)\overline{L(1/2, g, \chi)} \\ &= \sum_{m_1, m_2 \geq 1} \frac{A(m_1)\bar{B}(m_2)\chi(m_1)\bar{\chi}(m_2)}{(m_1 m_2)^{1/2}} e^{-m_1/x - m_2/x} + \mathcal{O}(x^{-C_3}), \end{aligned} \tag{10.4}$$

for arbitrarily large C_3.

Average over all characters χ modulo Q and let

$$S(Q)=\frac{1}{\varphi(Q)}\sum_{\chi \bmod Q} L(1/2,f,\chi)\overline{L(1/2,g,\chi)}. \tag{10.5}$$

Then

$$\begin{aligned} S(Q) = &\sum_{\substack{m_1\equiv m_2 \pmod Q\\ (m_2,Q)=1}} \frac{A(m_1)\bar{B}(m_2)}{(m_1m_2)^{1/2}} e^{-m_1/x-m_2/x} \\ = &\sum_{\substack{m\geq 1\\ (m,Q)=1}} \frac{A(m)\bar{B}(m)}{m} e^{-2m/x} + \sum_{\substack{m_2,h_0\geq 1\\ (m_2,Q)=1}} \frac{A(m_2+h_0Q)\bar{B}(m_2)}{((m_2+h_0Q)m_2)^{1/2}} e^{-2m_2/x-h_0Q/x} \\ &+ \sum_{\substack{m_2,h_0\geq 1\\ (m_2,Q)=1}} \frac{\bar{B}(m_2+h_0Q)A(m_2)}{((m_2+h_0Q)m_2)^{1/2}} e^{-2m_2/x-h_0Q/x} + \mathcal{O}(x^{-C_3}). \end{aligned} \tag{10.6}$$

By a similar inverse Mellin transform to that used above,

$$\sum_{\substack{m\geq 1\\ (m,Q)=1}} \frac{A(m)\bar{B}(m)e^{-2m/x}}{m} = \frac{1}{2\pi i}\int_{(2)} L_Q(s+1,f\otimes g)\Gamma(s)(x/2)^s ds.$$

Moving the line of integration to $\mathfrak{Re}\, s=-1+\epsilon$, if $f\neq g$ this becomes

$$\sum_{\substack{m\geq 1\\ (m,Q)=1}} \frac{A(m)\bar{B}(m)e^{-2m/x}}{m} = L_Q(1,f\otimes g)+\mathcal{O}(x^{-1+\epsilon}), \tag{10.7}$$

while if $f=g$ then

$$\sum_{m\geq 1} \frac{A(m)\bar{A}(m)e^{-2m/x}}{m} = \frac{L_Q(1,f,\vee^2)\log x}{\zeta_Q(2)} + c_f + \mathcal{O}(x^{-1+\epsilon}). \tag{10.8}$$

Both c_f and the implied constants depend, as usual, upon f and g. The Q subscript indicates that Euler factors dividing Q have been removed.

We have shown so far that

$$S(Q)=S_1(x)+S_2(x)+S_3(x)+S_4(x),$$

where

$$S_1(x)=\begin{cases} L_Q(1,f\otimes g) \text{ if } f\neq g\\ L_Q(1,f,\vee^2)\zeta_Q(2)^{-1}\log x + c_f \text{ if } f=g, \end{cases} \tag{10.9}$$

$$S_2(x)=\sum_{\substack{m_2,h_0\geq 1\\ (m_2,Q)=1}} \frac{A(m_2+h_0Q)\bar{B}(m_2)e^{-2m_2/x-h_0Q/x}}{((m_2+h_0Q)m_2)^{1/2}}, \tag{10.10}$$

$$S_3(x) = \sum_{\substack{m_2,h_0\geq 1\\(m_2,Q)=1}} \frac{A(m_2)\bar{B}(m_2+h_0Q)e^{-2m_2/x-h_0Q/x}}{((m_2+h_0Q)m_2)^{1/2}}, \tag{10.11}$$

and

$$S_4(x) \ll x^{-1+\epsilon}. \tag{10.12}$$

Note that although the left hand side is independent of x, the four contributions on the right hand side all depend on x. Interestingly, in the case $f = g$, as $x \to \infty$ the diagonal term, $S_1(x)$, tends to infinity, while $S(Q)$ is independent of x. In other words, $S_1(x)$ can not be the entire main term! Thus, at least in the case $f = g$, there must be a corresponding main term contribution from $S_2(x)$ and $S_3(x)$ canceling the multiple of $\log x$ in $S_1(x)$. In fact, we will see that $S_2(x)$ and $S_3(x)$ contribute to the main term even when $f \neq g$.

We would now like to transform $S_2(x)$, into a form where Proposition 12.3, to be proved in Section 12, applies. The first step is to remove the $(m_2, Q) = 1$ condition, which we do with a sieve as follows.

$$\begin{aligned}
S_2(x) &= \sum_{\substack{m_2,h_0\geq 1\\(m_2,Q)=1}} \frac{A(m_2+h_0Q)\bar{B}(m_2)e^{-2m_2/x-h_0Q/x}}{((m_2+h_0Q)m_2)^{1/2}}\\
&= \sum_{m_2,h_0\geq 1}\sum_{d|(m_2,Q)} \mu(d)\frac{A(m_2+h_0Q)\bar{B}(m_2)e^{-2m_2/x-h_0Q/x}}{((m_2+h_0Q)m_2)^{1/2}}\\
&= \sum_{d|Q}\mu(d)\sum_{m_2',h_0\geq 1}\frac{A(m_2'd+h_0Q)\bar{B}(m_2'd)e^{-2m_2'd/x-h_0Q/x}}{((m_2'd+h_0Q)m_2'd)^{1/2}}\\
&= \sum_{d|Q}\mu(d)\sum_{m_2',h_0\geq 1}\frac{A_d(m_2'+h_0Q/d)\bar{B}_d(m_2')e^{-2m_2'/(x/d)-h_0(Q/d)/(x/d)}}{d((m_2'+h_0(Q/d))m_2')^{1/2}}.
\end{aligned}$$

Here we have used $A_d(n) = A(dn), B_d(n) = B(dn)$ to denote the normalized Fourier coefficients of the level dN_0 forms $F_d(z), G_d(z)$, as described above.

Writing

$$S_2(x', d, Q') = \sum_{m_2,h_0\geq 1}\frac{A_d(m_2+h_0Q)\bar{B}_d(m_2)e^{-2m_2/x'-h_0Q'/x'}}{((m_2+h_0Q')m_2)^{1/2}},$$

so that $S_2(x) = S_2(x, 1, Q)$, we have shown that

$$S_2(x) = \sum_{d|Q}\frac{\mu(d)}{d}S_2(x/d, d, Q/d). \tag{10.13}$$

Now write $e^{-2m_2/x} = e^{-m_2/x}e^{-m_2/x}$ and substitute the series expansion for one of the $e^{-m_2/x}$, obtaining

$$e^{-2m_2/x-h_0Q/x} = e^{-m_2/x-h_0Q/x}\sum_{n=0}^{\infty}\frac{(-1)^n(m_2/x)^n}{n!},$$

Substituting into $S_2(x', d, Q')$ and interchanging the orders of summation leads to

$$S_2(x', d, Q') = \sum_{n=0}^{\infty}\frac{(-1)^n}{n!}\sum_{m_2,h_0\geq 1}\frac{A_d(m_2+h_0Q')\bar{B}_d(m_2)m_2^n e^{-m_2/x'-h_0Q'/x'}}{((m_2+h_0Q')m_2)^{1/2}x'^n}. \tag{10.14}$$

To apply Proposition 12.3 to this sum we take $\ell_1 = \ell_2 = 1$, replace f, g by F_d, G_d, and replace the level N_0 by dN_0. (The L in the order symbol is replaced by $Ld^{1/2}$.) This gives us

$$\begin{aligned}
&\sum_{\substack{m_2,h_0\geq 1\\ m_1=m_2+h_0Q}}\left(\frac{m_2}{x'}\right)^n\frac{A_d(m_1)\bar{B}_d(m_2)}{m_1^{1/2}m_2^{1/2}}e^{-(m_2+h_0Q')/x'}\\
&\qquad = \begin{cases} c(F_d)\log x' + c'(F_d) \text{ if } n=0 \text{ and } f=g\\ Z_{Q'}(1-k/2, 1/2; F_d, G_d) \text{ if } n=0 \text{ and } f\neq g\end{cases}\\
&\qquad\qquad + \mathcal{O}\left(Q^{\theta-1/2}\right) + \mathcal{O}\left(Q^{\theta}x^{-1/2}\right)\\
&\qquad\qquad + \mathcal{O}\left(Q'^{\theta-1/2}\log(d+1)x'^{-1/2}\right) + \mathcal{O}\left(Q'^{A}x'^{-C}\right).
\end{aligned}$$

Combining this with (10.13) and (10.14) gives us

$$\begin{aligned}
S_2(x) &= \sum_{d|Q}\frac{\mu(d)}{d}\sum_{n=0}^{\infty}\frac{(-m_2)^n}{(x/d)^n n!}\\
&\quad\times\sum_{m_2,h_0\geq 1}\frac{A_d(m_2+h_0Q/d)\bar{B}_d(m_2)e^{-m_2/x'-h_0(Q/d)/(x/d)}}{((m_2+h_0Q/d)m_2)^{1/2}}\\
&= \sum_{d|Q}\frac{\mu(d)}{d}\begin{cases} c_1(F_d)\log x + c_2(F_d) \text{ if } \ f=g\\ Z_{Q'}(1-k/2, 1/2; F_d, G_d) \text{ if } \ f\neq g\end{cases}\\
&\quad + \sum_{d|Q}\frac{\mu(d)}{d}\left(\mathcal{O}\left(Q^{\theta-1/2}\right) + \mathcal{O}\left(Q^{\theta}(x/d)^{-1/2}\right)\right.\\
&\qquad\qquad\qquad\qquad\qquad\qquad \left. + \mathcal{O}\left(Q^{A'}x^{-C'}\right)\right).
\end{aligned}$$

As

$$\sum_{d|q} \frac{\mu(d)}{d} \ll 1,$$

the above expression then simplifies to

$$S_2(x) = \sum_{d|Q} \frac{\mu(d)}{d} \begin{cases} c_1(F_d)\log x + c_2(F_d) \text{ if } \ f = g \\ Z_{Q'}(1-k/2, 1/2; F_d, G_d) \text{ if } \ f \neq g \end{cases}$$
$$+ \mathcal{O}\left(Q^{\theta-1/2}\right) + \mathcal{O}\left(Q^{\theta} x^{-1/2}\right) + \mathcal{O}\left(Q^{A'} x^{-C'}\right). \quad (10.15)$$

The same argument and bound apply to $S_3(x)$.

Collecting the above equalities and bounds, we have shown that

$$S(Q) = \begin{cases} L_Q(1, f\otimes g) \text{ if } f \neq g \\ L_Q(1, f, \vee^2)\zeta_Q(2)^{-1}\log x + c_f \text{ if } f = g, \end{cases}$$
$$+ \sum_{d|Q} \frac{\mu(d)}{d} \begin{cases} c_1(F_d)\log x + c_2(F_d) \text{ if } \ f = g \\ Z_{Q'}(1-k/2, 1/2; F_d, G_d) \text{ if } \ f \neq g \end{cases}$$
$$+ \sum_{d|Q} \frac{\mu(d)}{d} \begin{cases} c_1(F_d)\log x + c_2(F_d) \text{ if } \ f = g \\ Z_{Q'}(1-k/2, 1/2; G_d, F_d) \text{ if } \ f \neq g \end{cases}$$
$$+ \mathcal{O}\left(Q^{\theta-1/2}\right) + \mathcal{O}\left(Q^{\theta} x^{-1/2}\right) + \mathcal{O}\left(Q^{A'} x^{-C'}\right). \quad (10.16)$$

We can now let $x \to \infty$ in (10.16). When we do so it is clear that there must have an identity of constants:

$$L_Q(1, f, \vee^2)\zeta_Q(2)^{-1} + 2\sum_{d|Q} \frac{\mu(d)}{d} c_1(F_d) = 0,$$

where

$$c_1(F_d)\log x + c_2(F_d) = \mathrm{Res}_{w=0}\, \Gamma(w) x^w Z_Q(1-k/2, w+1/2).$$

The proposition then follows.

11. Additional multiple Dirichlet series

To investigate the behavior of $Z_Q(s,w)$ as Q varies, we define

$$\mathcal{M}(s,w,v) = \sum_{Q\geq 1} \frac{Z_{\ell Q}(s,w)}{Q^v}. \quad (11.1)$$

The ℓ is introduced for extra flexibility that we will need for the mean value application. The function $\mathcal{M}(s,w,v)$ would be more accurately written as $\mathcal{M}(s,w,v;\ell_1,\ell_2,\ell,f,g)$ but we will suppress the extra notation whenever it is possible to do so without confusion.

By the polynomial bounds in Q found in Propositions 7.1 and 7.2, this series will converge, for any fixed value of s,w, when the real part of v is sufficiently large. If the real parts of s,w and v are all sufficiently large then $Z_Q(s,w)$ can be expanded as a double Dirichlet series. Recalling its definition in (7.4) and bringing the sum over Q inside, we get

$$\mathcal{M}(s,w,v)=\sum_{\substack{m_2,h_0,Q\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q\ell}}\frac{A(m_1)\bar{B}(m_2)\left(1+\frac{h_0 Q\ell}{\ell_2 m_2}\right)^{(k-1)/2}}{(\ell_2 m_2)^s(h_0 Q\ell)^{w+(k-1)/2}Q^v} \tag{11.2}$$

Writing $m=h_0 Q$ this becomes

$$\begin{aligned}\mathcal{M}(s,w,v)&=\sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+\ell m}}\frac{A(m_1)\bar{B}(m_2)\left(1+\frac{\ell m}{\ell_2 m_2}\right)^{(k-1)/2}\sigma_{-v}(m)}{(\ell_2 m_2)^s(\ell m)^{w+(k-1)/2}}\\
&=(\ell_1\ell_2)^{(k-1)/2}\sum_{\substack{m_2,m\geq 1\\ \ell_1 m_1=\ell_2 m_2+\ell m}}\frac{a(m_1)\bar{b}(m_2)\sigma_{-v}(m)}{(\ell_2 m_2)^{s+k-1}(\ell m)^{w+(k-1)/2}}\\
&=(\ell_1\ell_2)^{(k-1)/2}\sum_{m\geq 1}\frac{\sigma_{-v}(m)D(s;\ell m)}{(\ell m)^{w+(k-1)/2}}\\
&=(\ell_1\ell_2)^{(k-1)/2}\sum_{m_2\geq 1}\frac{\bar{b}(m_2)D'(w,v,m_2)}{(\ell_2 m_2)^{s+k-1}},\end{aligned} \tag{11.3}$$

where

$$D'(w,v,m_2)=\sum_{\substack{m\geq 1\\ \ell_1 m_1=m\ell+\ell_2 m_2}}\frac{a(m_1)\sigma_{-v}(m)}{(\ell m)^{w+(k-1)/2}}. \tag{11.4}$$

We will obtain the analytic continuation of $D'(w,v,m_2)$ by a method very similar to the one used to analytically continue $D(s,w;h)$. The difference is some technical complications caused by the fact that the $\sigma_{-v}(m)$ are Fourier coefficients of Eisenstein series rather than cusp forms. Details are given in the following section.

11.1. **The analytic continuation of $D'(w,v,m_2)$.**

Our objective is to prove:

Proposition 11.1. *Fix $\epsilon>0$, v with $\mathfrak{Re}\, v>0$ and $B\gg 1$. For $\mathfrak{Re}\, w>1+\epsilon$ the Dirichlet series $D'(w,v,m_2)$ defined in (11.4) converges absolutely. It has an analytic continuation to all w with $\mathfrak{Re}\, w>1/2-B$,* *and v with $\mathfrak{Re}\, v \geq 0$, except for simple poles when $w+v/2 = 1/2+it_j-r$, for $r \geq 0$, and possible simple poles when $w+v/2 = -r$. The residues at the points $w+v/2 = 1/2+it_j-r$ are given by*

$$\mathrm{Res}_{w+v/2=1/2+it_j-r}\, D'(w,v,m_2)) = d_{r,j}(\ell)(\ell_2 m_2)^{r-it_j}\overline{\lambda_j(\ell_2 m_2)}, \quad (11.5)$$

where

$$d_{r,j}(\ell) = \frac{2^{-1+k/2-v/2}\pi^{-1+k/2-v/2}\Gamma((k+1+v)/2)}{\Gamma(k/2+it_j-r+v/2)\Gamma(k/2+it_j-r-v/2)} \times \frac{\Gamma(1/2+it_j-r)\Gamma(1/2-it_j+r)\Gamma(2it_j-r)(-1)^r\overline{\rho_j(-1)\left\langle U,u_j\right\rangle}}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)} \quad (11.6)$$

and $U(z) = y^{k/2}\overline{f(\ell_1 z)}E^{(k)}(\ell z,(1+v)/2)$. For $T \gg 1$ the $d_{r,j}$ satisfy the average upper bounds

$$\sum_{|t_j|\sim T} |d_{r,j}(\ell)|^2 \ll T^{2r+1}(\ell_1\ell)^{-k}.$$

When $\mathfrak{Re}\, w+v/2 < 1/2-k/2$ and $w+v/2$ is at least ϵ away from the poles, $D'(w,v,m_2)$ is given by the following absolutely convergent sum and integral:

$$\begin{aligned} D'(w,v,m_2) &= \pi^{(k-v-1)/2}2^k\ell^{(v-1)/2}(\ell_2 m_2)^{1/2-w-v/2} \\ &\quad\times \frac{\Gamma(1-w-v/2)\Gamma(w+v/2)\Gamma((k+1+v)/2)}{\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)}\left(S'_{cusp}+S'_{cont}\right) \\ &-\frac{\Gamma(1-w-v/2)\Gamma(w+v/2)}{\Gamma((1-v+k)/2)\Gamma((1+v-k)/2}\sum_{1\leq \ell m'<\ell_2 m_2}\frac{a((\ell_2 m_2-\ell m')/\ell)\sigma_{-v}(m')}{(\ell m')^{w+(k-1)/2}}. \end{aligned} \quad (11.7)$$

Here

$$S'_{cusp} = \sum_j \frac{\overline{\rho_j(-\ell_2 m_2)}\Gamma(w+v/2-1/2+it_j)\Gamma(w+v/2-1/2-it_j)\overline{\left\langle U,u_j\right\rangle}}{\Gamma(1/2+it_j)\Gamma(1/2-it_j)} \quad (11.8)$$

and

$$\begin{aligned} S'_{cont} &= \sum_a \int_{-\infty}^{\infty} \frac{\tilde{\rho}_a(1/2+it,-1)\sigma_{2it}(\ell_2 m_2)\pi^{1/2-it}(\ell_2 m_2)^{-it}}{4\pi\Gamma(1/2-it)^2\zeta(1-2it)\Gamma(1/2+it)} \\ &\times \Gamma(w+v/2-1/2+it)\Gamma(w+v/2-1/2-it)\overline{\left\langle U,E_a(*,1/2+it)\right\rangle}dt. \end{aligned} \quad (11.9)$$

In this region $D'(w,v,m_2)$ satisfies the upper bound

$$D'(w,v,m_2) \ll \ell^{(v-k)/2}\ell_1^{((1-k)/2}(\ell_2 m_2)^{1/2-w-v/2+\theta}(\log \ell\ell_1)^{1/2} \\ \times \frac{(1+|v|)^{k/2}}{(1+|w|)^{w+k/2-1}(1+|w+v|)^{w+k/2-1+v}}. \quad (11.10)$$

For $w+v/2$ at least ϵ away from the polar points, and in the vertical strip $c < \mathfrak{Re}\, w+v/2 < 2$, for any c with $1/2-B+\epsilon < c < 1/2-k/2$, $D'(w,v,m_2)$ is given by

$$\begin{aligned} D'(w,v,m_2) = {}& \frac{1}{2\pi i}\int_{2-i\infty}^{2+i\infty} \frac{D'(w,v,m_2)e^{(u-w)^2}du}{(u-w)} \\ & - \frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty} \frac{D'(w,v,m_2)e^{(u-w)^2}du}{(u-w)} \\ & - \sum_{t_j,0\le r<1/2-c-v/2} \frac{d_{r,j}(\ell)(\ell_2 m_2)^{r-it_j}\overline{\lambda_j(\ell_2 m_2)}e^{(1/2+it_j-r-v/2-w)^2}}{\Gamma(1/2+it_j-r)(1/2+it_j-r-v/2-w)}. \end{aligned} \quad (11.11)$$

In this same region it satisfies the upper bound

$$D'(w,v,m_2) \ll \ell^{(v-k)/2}\ell_1^{((1-k)/2}(\ell_2 m_2)^{1/2-c+\theta}(\log \ell\ell_1)^{1/2} \\ \times \frac{(1+|v|)^{k/2}}{(1+|w|)^{c-1}(1+|w+v|)^{c+k/2-1+v/2}}. \quad (11.12)$$

Proof. Define a partially holomorphic Eisenstein series of even weight k on $SL(2,\mathbb{Z})$ as follows:

$$E^{(k)}(z,s) = y^s + \sum_{(c,d)=1,c,d\ge 1} \frac{y^s|cz+d|^k}{|cz+d|^{2s}(cz+d)^k}.$$

The Fourier expansion of $E^{(k)}(z,s)$ is:

$$E^{(k)}(z,s) = \sum_m c^{(k)}(m,y,s)e^{2\pi i m x},$$

with

$$c^{(k)}(0,y,s) = y^s + \frac{y^{1-s}\zeta(2s-1)2^{2-2s}\pi\Gamma(1+k/2-s)\Gamma(2s-1)}{\zeta(2s)\Gamma(s)\Gamma(1-s)\Gamma(s+k/2)}, \quad (11.13)$$

and for $m \neq 0$

$$c^{(k)}(m,y,s) = \frac{\sigma_{1-2s}(|m|)\pi^s|m|^{s-1}}{\zeta(2s)\Gamma(s\pm k/2)}W_{\pm k/2,1/2-s}(4\pi|m|y). \quad (11.14)$$

Here the plus or minus sign is used according as m is positive or negative.

For reference, we record here the behavior of $W_{\lambda,\mu}(y)$ at large and small values of $y > 0$. As $y \to 0$ and $\mathfrak{Re}\,\mu \neq 0$,

$$W_{\lambda,\mu}(y) \ll y^{1/2-|\mathfrak{Re}\,\mu|}.$$

(If $\mathfrak{Re}\mu = 0$ the upper bound is $y^{1/2}|\log y|$.) For $y \gg 1$,

$$W_{\lambda,\mu}(y) \ll e^{-y/2}y^{\lambda}.$$

In analogy to the original $P_{h,Y}(z,s,\delta)$, defined in (2.2), define the following weight k Poincare series for $SL(2,\mathbb{Z})$: For $\delta > 0$, $Y \gg 1$, $m' \geq 0$, $\mathfrak{Re}\,s > 1$, set

$$\begin{aligned} &P^{(k)}_{\delta,Y}(z,s,m,m_2,\ell_1,\ell_2,\ell) \\ &\quad = \sum_{\gamma\in\Gamma_\infty\backslash SL(2,\mathbb{Z})} \frac{y^s|cz+d|^k}{|cz+d|^{2s}(cz+d)^k} e^{-2\pi i\ell m\gamma z - 2\pi\ell_2\delta m_2 \mathfrak{Im}\,\gamma z}\psi_Y(\mathfrak{Im}\,\gamma z). \end{aligned} \tag{11.15}$$

In analogy to $\mathcal{I}_{\ell_1,\ell_2,\ell,Y,\delta}(s;h)$, we now define, for v real, $v > 0$, and for w with $\mathfrak{Re}\,w + v/2 > 1$

$$\begin{aligned} &\mathcal{I}(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y) \\ &:= \frac{1}{\mathcal{V}}\iint_{\Gamma\backslash\mathbb{H}} P_{\ell_2 m_2,Y}(z,w+v/2;\delta)y^{k/2}f(\ell_1 z)\overline{E^{(k)}(\ell z,(1+v)/2)}\frac{dx\,dy}{y^2} \\ &\qquad - \frac{1}{\mathcal{V}}\iint_{SL(2,\mathbb{Z})\backslash\mathbb{H}} \sum_{\ell_2 m_2 < \ell m \leq 0} a((\ell_2 m_2 + \ell m)/\ell_1) \\ &\quad \times P^{(k)}_{\delta,Y}(z,w+v/2+k/2,m,m_2,\ell_1,\ell_2,\ell)\overline{E^{(k)}(\ell z,(1+v)/2)}\frac{dx\,dy}{y^2}. \end{aligned} \tag{11.16}$$

Here the first Poincare series $P_{\ell_2 m_2,Y}(z,w+v/2;\delta)$ is the original series of (2.2), with $w+v/2$ in place of s and $\ell_2 m_2$ in place of h, and is a sum over the group $\Gamma_0(N_0\ell_1\ell)$. Although originally defined for $\mathfrak{Re}\,w+v/2 > 1$, the meromorphic continuation of the inner product is obtainable. The subtracted term is defined and convergent for $\mathfrak{Re}\,w+v/2+k/2 > 1$, and consequently has no poles in this region. This second integration is over a fundamental domain for $SL(2,\mathbb{Z})$, (although the $\mathcal{V}$ in the denominator is the original volume of the fundamental domain for Γ).

Write

$$\mathcal{I}(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y)=\mathcal{I}_0(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y) - \sum_{0\le m'<\ell_2 m_2} d_{\delta,Y}(w,v,m',m_2), \quad (11.17)$$

where

$$\mathcal{I}_0(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y)= \Big\langle P_{\ell_2 m_2,Y}(*,w+v/2;\delta), y^{k/2}\overline{f(\ell_1 *)}E^{(k)}(\ell *,(1+v)/2)\Big\rangle \quad (11.18)$$

and

$$d_{\delta,Y}(w,v,m',m_2) = \frac{1}{\mathcal{V}}\iint_{SL(2,\mathbb{Z})\backslash\mathbb{H}} a((\ell_2 m_2-\ell m')/\ell_1) \times P^{(k)}_{\delta,Y}(z,w+v/2+k/2,-m',m_2,\ell_1,\ell_2,\ell)\overline{E^{(k)}(\ell z,(1+v)/2)}\frac{dx\,dy}{y^2}. \quad (11.19)$$

First we will interpret both integrals as a Dirichlet series, take the difference, and then take the limit as first $Y\to\infty$, and then $\delta\to 0$. Unfolding the two Poincare series, replacing m' by $-m$ and performing the x integration we obtain

$$\begin{aligned}\mathcal{I}(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y) &= \Bigg(\frac{1}{\mathcal{V}}\int_{Y^{-1}}^{Y} y^{w+k/2+v/2-1} \\ &\times \sum_{\ell m>-\ell_2 m_2} a((\ell_2 m_2+\ell m)/\ell_1)e^{-2\pi y(\ell m+\delta\ell_2 m_2)}\overline{c^{(k)}(m,\ell y,(1+v)/2)}\frac{dy}{y}\Bigg) \\ &- \frac{1}{\mathcal{V}}\int_{Y^{-1}}^{Y} y^{w+k/2+v/2-1}\sum_{0\ge \ell m>-\ell_2 m_2} a((\ell_2 m_2+\ell m)/\ell_1)e^{-2\pi y(\ell m+\delta\ell_2 m_2)} \\ &\times \overline{c^{(k)}(m,\ell y,(1+v)/2)}\frac{dy}{y} \\ &= \frac{1}{\mathcal{V}}\int_{Y^{-1}}^{Y} y^{w+k/2+v/2-1}\sum_{m\ge 1} a((\ell_2 m_2+\ell m)/\ell_1)e^{-2\pi y(\ell m+\delta\ell_2 m_2)} \\ &\times \overline{c^{(k)}(m,\ell y,(1+v)/2)}\frac{dy}{y}. \quad (11.20)\end{aligned}$$

For $m \geq 1$, (11.14) becomes

$$c^{(k)}(m, y, (1+v)/2) = \frac{\sigma_{-v}(|m|)\pi^{(v+1)/2}|m|^{(v-1)/2}}{\zeta(v+1)\Gamma(k/2+(1+v)/2)} W_{k/2,-v/2}(4\pi|m|y).$$

Also, by [14], 7.621.3,

$$\int_0^\infty y^{w+k/2+v/2-1} e^{-y(1+\delta)} W_{k/2,-v/2}(2y)\frac{dy}{y}$$
$$= \frac{\Gamma(w+k/2-1/2)\Gamma(w+k/2-1/2+v)2^{-w-k/2+1-v/2}}{\Gamma(w+v/2)}\left(1+\mathcal{O}(\delta)\right).$$

Substituting these two facts into (11.20) we first take the limit as $Y \to \infty$, obtaining

$$\lim_{Y\to\infty} \mathcal{I}(w, v, m_2, \ell_1, \ell_2, \ell, \delta, Y)$$
$$= \frac{\pi^{-w-k/2+3/2}2^{-2w-v-k+2}\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)}{\mathcal{V}\zeta(1+v)\Gamma((k+1+v)/2)\Gamma(w+v/2)}$$
$$\times \sum_{m\geq 1} \frac{a((\ell_2 m_2 + \ell m)/\ell_1)\sigma_{-v}(m)m^{(v-1)/2}}{(\ell m + \delta\ell_2 m_2)^{w+v/2+k/2-1}}\left(1+\mathcal{O}(\delta)\right).$$

Then, taking the limit as $\delta \to 0$, we define

$$\mathcal{I}(w, v, m_2, \ell_1, \ell_2, \ell) = \lim_{\delta\to 0}\lim_{Y\to\infty} \mathcal{I}(w, v, m_2, \ell_1, \ell_2, \ell, \delta, Y)$$
$$= \frac{G(w, v)D'(w, v, m_2)}{\ell^{v/2-1/2}}, \quad (11.21)$$

where

$$G(w, v)$$
$$= \frac{\pi^{-w-k/2+3/2}2^{-2w-v-k+2}\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)}{\mathcal{V}\zeta(1+v)\Gamma((k+1+v)/2)\Gamma(w+v/2)} \quad (11.22)$$

This representation is valid for $\mathfrak{Re}\, w + v/2 > 1$. Our objective is to now use the representation of $\mathcal{I}(w, v, m_2, \ell_1, \ell_2, \ell, \delta, Y)$ given in (11.17) to find a spectral decomposition of $\mathcal{I}_0(w, v, m_2, \ell_1, \ell_2, \ell, \delta, Y)$. We will then take a limit again as $Y \to \infty$ and $\delta \to 0$ in the region $\mathfrak{Re}\, w + v/2 < 1/2 - k/2$, connecting this region, in the process, to the region $\mathfrak{Re}\, w + v/2 > 1$. We will thus obtain a meromorphic continuation of $\mathcal{I}(w, v, m_2, \ell_1, \ell_2)$, and hence $D'(w, v, m_2)$, to all w, v with $\mathfrak{Re}\, w + v/2 > 1/2 - B$, for a sufficiently large $B \gg 1$, as was done in the case of $D(s; h)$. Unfolding the Poincare series in (11.19)

$$d_{\delta,Y}(w,v,m',m_2) \\ = \frac{1}{\mathcal{V}}\int_{Y^{-1}}^{Y} y^{w+k/2+v/2-1} \\ \times \sum_{0\le m'<\ell_2 m_2} a((\ell_2 m_2 - \ell m')/\ell_1)e^{-2\pi y(-\ell m'+\delta\ell_2 m_2)}\overline{c^{(k)}(-m',\ell y,(1+v)/2)}\frac{dy}{y}. \tag{11.23}$$

Substituting (11.14) into (11.23) gives us, for $m' \ge 1$

$$\begin{aligned} d_{\delta,Y}(w,v,m',m_2) &= \frac{1}{\mathcal{V}}\frac{a((\ell_2 m_2-\ell m')/\ell_1)\sigma_{-v}(m')\pi^{(v+1)/2}m'^{(v-1)/2}}{\zeta(1+v)\Gamma((1+v-k)/2)} \\ &\times \int_{Y^{-1}}^{Y} y^{w+k/2+v/2-1}W_{-k/2,-v/2}(4\pi\ell m' y)e^{2\pi y(\ell m'-\delta\ell_2 m_2)}\frac{dy}{y} \\ &= \frac{2^{-w-v/2-k/2+1}\pi^{3/2-w-k/2}}{\mathcal{V}\zeta(1+v)\Gamma((1+v)/2-k/2)}\frac{a((\ell_2 m_2-\ell m')/\ell_1)\sigma_{-v}(m')}{(\ell m')^{w+(k-1)/2}} \\ &\times \int_{Y^{-1}2\pi\ell m'}^{Y2\pi\ell m'} y^{w+k/2+v/2-1}W_{-k/2,-v/2}(2y)e^{y(1-\delta\ell_2 m_2/m')}\frac{dy}{y}. \end{aligned} \tag{11.24}$$

The integral here is very similar to that studied in Section 3. It has been analyzed and its analytic continuation obtained, by Tom Hulse in [Hulse]. Using his notation, upon taking the limit as $Y \to \infty$, we have

$$\lim_{Y\to\infty}\int_{Y^{-1}2\pi m'}^{Y2\pi m'} y^{w+k/2+v/2-1}W_{-k/2,-v/2}(2y)e^{y(1-\delta\ell_2 m_2/m')}\frac{dy}{y} \\ = M_2(w+k/2+v/2, iv/2, \delta\ell_2 m_2/m'),$$

In [16] it is shown in Proposition 3.1 that for each $\delta > 0$ the function $M_2(w+k/2+v/2, iv/2, \delta\ell_2 m_2/m')$ has analytic continuation to $\mathbb{C}$, with at most polynomial growth in vertical strips in the variables w, v, δ^{-1}, except for simple poles at $w + k/2 + v/2 = 1/2 \pm v/2 - r$, for $r \ge 0$. (Note that as in Proposition 3.1, there is no pole at $w + v/2 = 1$.) It is also shown that for $\mathfrak{Re}\, w + k/2 + v/2 \le 1/2$, the limit as $\delta \to 0$ of $M_2(w + k/2 + v/2, iv/2, \delta\ell_2 m_2/m')$, exists and that we have, for $\mathfrak{Re}\, w + k/2 + v/2 \le 1/2$,

$$M_2(w+k/2+v/2, iv/2) \\ = \lim_{\delta\to 0} M_2(w+k/2+v/2, iv/2, \delta\ell_2 m_2/|m|) \\ = \frac{2^{1-w-k/2-v/2}\Gamma(w+k/2-1/2)\Gamma(w+k/2+v-1/2)\Gamma(1-w-v/2)}{\Gamma((1+k+v)/2)\Gamma((1+k-v)/2)}.$$

With this notation, after taking the limit as $Y \to \infty$ and $\delta \to 0$ we set

$$\begin{aligned} d(w,v,m',m_2) &= \lim_{\substack{\delta\to 0\\ Y\to\infty}} d_{\delta,Y}(w,v,m',m_2) \\ &= \frac{2^{-w-v/2-k/2+1}\pi^{3/2-w-k/2}}{\mathcal{V}\zeta(1+v)\Gamma((1+v)/2-k/2)} M_2(w+k/2+v/2, iv/2) \\ &\qquad \times \frac{a((\ell_2 m_2 - \ell m')/\ell_1)\sigma_{-v}(m')}{(\ell m')^{w+(k-1)/2}\ell^{v/2-1/2}} \\ &= \frac{R(w,v)F(w,v,m',m_2)}{\ell^{v/2-1/2}}, \end{aligned} \tag{11.25}$$

where

$$\begin{aligned} R(w,v) &= \frac{4^{-w-v/2-k/2+1}\pi^{3/2-w-k/2}\Gamma(w+(k-1)/2)}{\mathcal{V}\zeta(1+v)\Gamma((1+v-k)/2)\Gamma((1+v+k)/2)} \\ &\qquad \times \frac{\Gamma(w+(k-1)/2+v)\Gamma(1-w-v/2)}{\Gamma((1-v+k)/2)} \end{aligned} \tag{11.26}$$

and

$$F(w,v,m',m_2) = \frac{a((\ell_2 m_2 - \ell m')/\ell_1)\sigma_{-v}(m')}{(\ell m')^{w+(k-1)/2}}. \tag{11.27}$$

Turning to the case $m' = 0$, replace s by $(1+v)/2$ in (11.13) and write

$$c^{(k)}(0,y,(1+v)/2) = y^{(1+v)/2} + c(v)y^{(1-v)/2}, \tag{11.28}$$

where

$$c(v) = \frac{\zeta(v)2^{1-v}\pi\Gamma(k/2+(1-v)/2)\Gamma(v)}{\zeta(v+1)\Gamma((1+v)/2)\Gamma((1-v)/2)\Gamma((k+v+1)/2)} \tag{11.29}$$

Substituting the above into (11.23) gives

$$\begin{aligned} d_{\delta,Y}(w,v,0,m_2) &= \frac{1}{\mathcal{V}}\int_{Y^{-1}}^{Y} y^{w+k/2+v/2-1} \\ &\qquad \times a((\ell_2 m_2)/\ell_1)e^{-2\pi y(\delta\ell_2 m_2)}\overline{c^{(k)}(0,\ell y,(1+v)/2)}\frac{dy}{y}. \end{aligned}$$

Changing variables $y \to y/\ell$ yields

$$\begin{aligned} d_{\delta,Y}(w,v,0,m_2) &= \frac{\ell^{-(w+k/2+v/2-1)}}{\mathcal{V}}\int_{\ell Y^{-1}}^{\ell Y} y^{w+k/2+v/2-1} \\ &\qquad \times a((\ell_2 m_2)/\ell_1)e^{-2\pi y(\delta\ell_2 m_2)}\overline{c^{(k)}(0,y,(1+v)/2)}\frac{dy}{y}, \end{aligned}$$

and then expanding $c^{(k)}(0,y,(1+v)/2)$ and letting $y \to y/2\pi\delta\ell_2 m_2$ yields

$$d_{\delta,Y}(w,v,0,m_2) = \frac{\ell^{-(w+k/2+v/2-1)}a((\ell_2 m_2)/\ell_1)}{\mathcal{V}}$$
$$\times \left(\frac{1}{(2\pi\delta\ell_2 m_2)^{w+k/2+v-1/2}} \int_{2\pi\delta\ell_2 m_2 \ell Y^{-1}}^{2\pi\delta\ell_2 m_2 \ell Y} y^{w+k/2+v-1/2} e^{-y} \frac{dy}{y} \right.$$
$$\left. + \frac{c(v)}{(2\pi\delta\ell_2 m_2)^{w+k/2-1/2}} \int_{2\pi\delta\ell_2 m_2 \ell Y^{-1}}^{2\pi\delta\ell_2 m_2 \ell Y} y^{w+k/2-1/2} e^{-y} \frac{dy}{y} \right).$$

For $\mathfrak{Re}\, w + k/2 + (v \pm v)/2 > 1/2$, this is absolutely convergent as $Y \to \infty$, and converges to

$$d_\delta(w,v,0,m_2) := \lim_{Y\to\infty} d_{\delta,Y}(w,v,0,m_2) =$$
$$\frac{a((\ell_2 m_2)/\ell_1)}{\mathcal{V}\ell^{w+k/2+v/2-1}} \left(\frac{\Gamma(w+k/2+v-1/2)}{(2\pi\delta\ell_2 m_2)^{w+k/2+v-1/2}} + \frac{c(v)\Gamma(w+k/2-1/2)}{(2\pi\delta\ell_2 m_2)^{w+k/2-1/2}} \right) \tag{11.30}$$

which has a continuation to all $w, v \in \mathbb{C}$ with poles at $w + k/2 + (v \pm v)/2 - 1/2 = -r$ for $r \geq 0$. Furthermore, we note that for $\mathfrak{Re}\, w + k/2 + (v \pm v)/2 < 1/2$, the limit of $d_\delta(w,v,0,m_2)$ as $\delta \to 0$ is 0.

Recalling (11.17), the next step in the meromorphic continuation of $\mathcal{I}(w,v,H,\ell_1,\ell_2,\ell)$, and hence also of $D(w,v,m_2)$ will be achieved once the analytic continuation of

$$\mathcal{I}_0(w,v,m_2,\ell_1,\ell_2,\ell) = \lim_{\substack{\delta\to 0 \\ Y\to\infty}} \mathcal{I}_0(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y) \tag{11.31}$$

is obtained. To accomplish this, the same analysis as in Section 2, applies, as we are using the same Poincare series (with ℓ in place of ℓ_2) and only changing the automorphic function V to

$$U = y^{k/2}\overline{f(\ell_1 z)} E_\ell^{(k)}(z,(1+v)/2),$$

where $E_\ell^{(k)}(z,(1+v)/2) = E^{(k)}(\ell z,(1+v)/2)$. The function U still vanishes at the cusps and is automorphic with respect to $\Gamma_0(\ell_1\ell)$. The resulting spectral expansion of the triple product is

$$\begin{aligned}
\mathcal{I}_0(w,v,m_2,\ell_1,\ell_2,\ell,\delta,Y) &= \langle P_{\ell_2 m_2,Y}(*,w+v/2;\delta),U\rangle \\
&= \sum_{j\geq 1} \overline{\langle U,u_j\rangle}\, \langle P_{\ell_2 m_2,Y}(*,w+v/2;\delta),u_j\rangle \\
&\quad + \frac{1}{4\pi}\sum_{a}\int_{-\infty}^{\infty} \overline{\langle U,E_a(*,1/2+it)\rangle} \\
&\quad \times \langle P_{\ell_2 m_2,Y}(*,w+v/2;\delta),E_a(*,1/2+it)\rangle\,dt \\
&= \frac{(2\pi\ell_2 m_2)^{1/2-w-v/2}}{\mathcal{V}}\Bigg(\sum_j \overline{\rho_j(-\ell_2 m_2)}M(w+v/2,t_j,\delta)\overline{\langle U,u_j\rangle} \\
&\quad + \sum_a \int_{-\infty}^{\infty} \frac{\overline{\tilde{\rho}_a(1/2+it,-\ell_2 m_2)}\sigma_{2it}(\ell_2 m_2)\pi^{1/2-it}}{2\pi\zeta(1-2it)\Gamma(1/2-it)(\ell_2 m_2)^{it}} \\
&\quad \times M(w+v/2,t,\delta)\overline{\langle U,E_a(*,1/2+it)\rangle}dt\Bigg). \qquad (11.32)
\end{aligned}$$

As was the case with $D(s;h)$, this spectral expansion has poles coming from the cuspidal contribution at $w+v/2 = 1/2+it_j-r$, with $r\geq 0$, and no poles originating in the continuous contribution. By the upper bound for $M(w+v/2,t,\delta)$ given in (3.2) this converges absolutely for any fixed δ. To determine the region of absolute convergence as $\delta\to 0$, we note first that by (3.3), (and Stirling's formula), when δ is small relative to t_j

$$M(w+v/2,t,\delta) \ll_{w,v} (1+|t_j|)^{2\mathfrak{Re}\,(w+v/2)-2}, \qquad (11.33)$$

where the implied constant depends on w,v. The inner product $\overline{\langle U,u_j\rangle}$ can be explicitly evaluated in this case as g is replaced by the Eisenstein series. For example, in the case $\ell_1=\ell=N_0=1$, this simplifies to

$$\begin{aligned}
\overline{\langle U,u_j\rangle} &= \int_{SL(2,\mathbb{Z})\backslash\mathbb{H}} u_j(z)y^{k/2}f(z)\overline{E^{(k)}(z,(1+v)/2)}\frac{dxdy}{y^2} \\
&= \int_0^\infty\int_0^1 \sum_{m_1\neq 0}\rho_j(m_1)a(m_2)e^{2\pi i m_2 z}y^{(k+\bar{v})/2}K_{it_j}(2\pi|m_1|y)e^{2\pi i m_1 x}\frac{dxdy}{y} \\
&= \frac{\rho_j(-1)\sqrt{\pi}L((1+\bar{v})/2,u_j\otimes f)\Gamma((\bar{v}+k)/2+it_j)\Gamma((\bar{v}+k)/2-it_j)}{(4\pi)^{(\bar{v}+k)/2}\Gamma((1+\bar{v}+k)/2)}.
\end{aligned} \qquad (11.34)$$

Here we have done a Rankin-Selberg unfolding, applied the identity

$$\int_0^\infty x^{\mu-1}e^{-x}K_\nu(x)dx = \frac{\sqrt{\pi}2^\nu\Gamma(\mu+\nu)\Gamma(\mu-\nu)}{2^{\mu+\nu}}$$

of [14] 6.621. 3 (with $\alpha = \beta = 1$), and set

$$L(s, u_j \otimes f) = \sum_{m \geq 1} \frac{\lambda_j(m) a(m)}{m^{s+(k-1)/2}}.$$

Replacing $u_j(z)$ by $E(z, 1/2 + it)$, we have in this special case

$$\begin{aligned}\overline{\langle U, E(*, 1/2 + it)\rangle} \\ &= \frac{\Gamma(\bar{v}/2 + k/2 + it)\Gamma(\bar{v}/2 + k/2 - it)}{2^{\bar{v}+k-1}\pi^{\bar{v}/2+k/2-1-it}\Gamma(1/2 + it)\zeta(1 + 2it)\zeta(v + k)} \\ &\quad \times L(\bar{v}/2 + k/2 + it, f) L(\bar{v}/2 + k/2 - it, f). \quad (11.35)\end{aligned}$$

In the general case, substituting $E_\ell(z, (1 + v)/2)$ for g in the triple product estimate of Proposition 4.4, we now have instead

$$\sum_{|t_j| \sim T} \overline{\rho_j(-1)\langle U, u_j\rangle} \ll (\ell_1 \ell)^{(1-k)/2} T^{1+k} (\log(\ell_1 \ell T))^{1/2}, \qquad (11.36)$$

with a corresponding and improved estimate for the continuous part.

Substituting (11.33) and (11.36) into (11.32), after applying Cauchy-Schwarz and making a dyadic subdivision of the interval $[0, T]$ we see that the exponent of T will be bounded above by $2\mathfrak{Re}\, w + v/2 + k - 1$ and thus will be negative when $\mathfrak{Re}\, w + v/2 < 1/2 - k/2$. It follows that for such w, v the spectral expansion (11.32) will converge absolutely as $\delta \to 0$. The upper bound for D' claimed in the proposition follows from an examination of of the different components of $\mathcal{I}$ and the above discussion.

The full analytic continuation of $\mathcal{I}(w, v, m_2, \ell_1, \ell_2, \ell)$ is then achieved just as in the previous case, by using the analysis of $M_2(w + v/2, t, \delta)$ in [16], together with Cauchy's theorem, to fill in the gap between the regions $\mathfrak{Re}\, w + v < 1/2 - k/2$ where the spectral side converges, and the region $\mathfrak{Re}\, w + v/2 > 1$, where the Dirichlet series converges. □

11.2. The analytic continuation of $\mathcal{M}(s, w, v)$.

Consider one of the expressions for $\mathcal{M}(s, w, v)$ in (11.3):

$$\mathcal{M}(s, w, v) = (\ell_1 \ell_2)^{(k-1)/2} \sum_{m \geq 1} \frac{\sigma_{-v}(m) D(s; \ell m)}{(\ell m)^{w+(k-1)/2}}. \qquad (11.37)$$

Recall from Proposition 5.1, line (5.7), that for s a distance of at least ϵ from the polar points and in the region $c < \mathfrak{Re}\, s < 2$, for any c with $1/2 - A + \epsilon < c < 1/2 - k/2$, (with $\ell_1, \ell_2 \ll L$),

$$D(s; m) \ll (1 + |s|)^{-c-k+3/2} \log(L) L^{1-k} m^{1/2-c+\theta}.$$

It follows that when $\mathfrak{Re}\, w > A + 3/2 + \theta - k/2$ the expression above converges absolutely. Also, any poles it has must arise as poles of $D(s;m)$ and be at points $s = 1/2 + it_j - r, r \geq 0$. When $\mathfrak{Re}\, s < 1/2 - k/2$ the spectral expansion for $D(s;h)$ given in Proposition 5.1 converges absolutely. Substituting this for $D(s;h)$, replacing h by ℓm, we can, because of the absolute convergence, interchange the order of summation, obtaining

$$\mathcal{M}(s,w,v) = \frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell^{s'}}(\mathcal{M}_{\text{cusp}} + \mathcal{M}_{\text{cont}}), \tag{11.38}$$

with

$$\mathcal{M}_{\text{cusp}} = \frac{2^{2k-1}\pi^k\Gamma(1-s)}{\Gamma(s+k-1)} \sum_j L_\ell(s',\overline{u_j})L_\ell(s'+v,\overline{u_j})\mathcal{P}(s',v) \\ \frac{\Gamma(s-\frac{1}{2}-it_j)\Gamma(s-\frac{1}{2}+it_j)\overline{\rho_j(-1)\langle V,u_j\rangle}}{\zeta(2s'+v)\Gamma(\frac{1}{2}-it_j)\Gamma(\frac{1}{2}+it_j)} \tag{11.39}$$

and

$$\mathcal{M}_{\text{cont}} = \frac{2^{2k-1}\pi^{k-1/2}\Gamma(1-s)}{4\pi\Gamma(s+k-1)} \\ \times \sum_a \int_{-\infty}^{\infty} \zeta_a(s'+it)\zeta_a(s'-it)\zeta_a(s'+v+it)\zeta_a(s'+v-it) \\ \times \frac{\mathcal{P}(s',v,it)\Gamma(s-\frac{1}{2}-it)\Gamma(s-\frac{1}{2}+it)\overline{\langle V,E_a^*(*,\frac{1}{2}+it)\rangle}}{\mathcal{V}\zeta(1+it)\zeta(1-2it)\zeta(2s'+v)\Gamma(\frac{1}{2}-it)^2\Gamma(\frac{1}{2}+it)^2}dt, \tag{11.40}$$

where $s' = s + w + k/2 - 1$. Here the L-series $L_\ell(s',\overline{u_j})$ is the usual L-series with Euler factors corresponding to primes dividing ℓ removed and

$$\mathcal{P}(s',v) = \prod_{p^\gamma || \ell} \Big(\lambda_j(p^\gamma) - \lambda_j(p^{\gamma-1})p^{-s'} - \lambda_j(p)p^{-2s'-v} \\ +p^{-3s'}\lambda_j(p^{\gamma-1}) + p^{-3s'-v}\lambda_j(p^{\gamma-1})\Big). \tag{11.41}$$

The function ζ_a differs from ζ at Euler factors of primes dividing ℓN_0. Note that $\mathcal{P}(s',v) \ll \lambda_j(\ell)$ for $\mathfrak{Re}\, s' \geq 1/2$ and $\mathfrak{Re}\, v \geq 0$. We have multiplied the numerator and denominator by $\zeta^*(1+2it)$ to normalize the Eisenstein series in the inner product.

The expressions (11.38)–(11.40) allow us to obtain a meromorphic continuation of $\mathcal{M}(s,w,v)$ back to $\mathfrak{Re}\, v \geq 0$, as long as the real part of w is sufficiently large and the real part of s is sufficiently negative. Specifically, consider the cuspidal piece. Suppose that $\mathfrak{Re}\, v \geq 0$ and $\mathfrak{Re}\, s' < 0$ Then by the functional equation, for $t_j \sim T$

$$L(s', \bar{u}_j) L(s', \bar{u}_j) \ll_{\ell_1 \ell_2, \ell, s', v} T^{2-4s'}.$$

Also, recall that we had

$$\sum_{|t_j| \sim T} |\overline{\rho_j(-1)\langle V, u_j \rangle}|^2 \ll_L T^{2k},$$

Combining this with Stirling's formula, Cauchy-Schwarz and Weyl's law we see that the sum over j in (11.39) converges when $\mathfrak{Re}\, s < 1/2 - k/2$ and

$$2 - 4s' + 2s - 2 + 1 + k < 0.$$

As $s' = s + w + k/2 - 1$ this translates to the conditions $\mathfrak{Re}\, s < 1/2 - k/2$ and

$$w > -\frac{s}{2} - \frac{k}{4} + \frac{5}{4}. \tag{11.42}$$

If $\mathfrak{Re}\, v \geq 0$, and $\mathfrak{Re}\, s' \geq 1/2$ then, as

$$\sum_{|t_j| \sim T} |L(1/2, \bar{u}_j)|^2 \ll_L T^2,$$

the condition for convergence is $2s - 2 + 1 + k < 0$, which translates to the conditions $\mathfrak{Re}\, s < 1/2 - k/2$ and $\mathfrak{Re}\, w > 1$. Finally, if $\mathfrak{Re}\, v \geq 0$, and $0 \leq \mathfrak{Re}\, s' \leq 1/2$ the condition becomes again

$$w > -\frac{s}{2} - \frac{k}{4} + \frac{5}{4}.$$

Collecting this information, we have shown that the sum over j in (11.39) converges when $-k/2 < \mathfrak{Re}\, s < 1/2 - k/2$ and $\mathfrak{Re}\, w > 1$, and also when $\mathfrak{Re}\, s \leq -k/2$ and $w > -s/2 - k/4 + 5/4$.

The meromorphic continuation of the continuous piece is a little more subtle as the zeta factors only converge when $\mathfrak{Re}\, s' > 1$ and $\mathfrak{Re}\, v \geq 0$. To continue past this it is necessary to shift the line of integration to the right and then back to 0 again, as was done for the corresponding integral in the continuous part of $Z_Q(s, w)$. Specifically, suppose that $\mathfrak{Re}\, s < 1$, $\mathfrak{Re}\, s' > 1$, $\mathfrak{Re}\, v > 0$. Set $u = it$, and move the line of integration to the right, past the poles at $u = s' - 1, s' + v - 1$. After moving the line, continue to $\mathfrak{Re}\, s' < 1$, and then move the line back to $\mathfrak{Re}\, u = 0$. $\mathfrak{Re}\, u = -\mathfrak{Re}\,(1 - s' - v) - \epsilon$. Poles are passed over at the points $u = 1 - s'$, $u = 1 - s' - v$, and at $u = s - 1/2 + r_1$, $u = 1/2 - s - r_2$

for r_1, r_2 such that have real part of u lies between $1 - \mathfrak{Re}\,(s' + v)$ and 0. The contributions from the residues at these points are as follows:

$$\operatorname{Res}_{u=s'-1} = f_1(s,w)\zeta_a(1+v)\zeta_a(2s'-1+v)\zeta^*(2s'-1) \times \frac{\Gamma(s+1/2-s')\Gamma(s-3/2+s')\overline{\langle V, E_a^*(*,s'-1/2)\rangle}}{\zeta_a(3-2s')\zeta(2s'+v)\Gamma(s'-1/2)^2\Gamma(3/2-s')^2}, \quad (11.43)$$

$$\operatorname{Res}_{u=s'+v-1} = f_2(s,w)\zeta_a(1-v)\zeta_a(2s'-1+v)\zeta^*(2s'+2v-1) \times \frac{\Gamma(s+1/2-s'-v)\Gamma(s-3/2+s'+v)\overline{\langle V, E_a^*(*,s'+v-1/2)\rangle}}{\zeta_a(3-2s'-2v)\zeta(2s'+v)\Gamma(s'+v-1/2)^2\Gamma(3/2-s'-v)^2}, \quad (11.44)$$

where

$$f_1(s,w) = \frac{2^{2k-2}\pi^{k-1/2}\mathcal{P}(s',v,s'-1)\Gamma(1-s)}{\Gamma(s+k-1)\mathcal{V}},$$

and

$$f_2(s,w) = \frac{2^{2k-2}\pi^{k-1/2}\mathcal{P}(s',v,s'+v-1)\Gamma(1-s)}{\Gamma(s+k-1)\mathcal{V}},$$

Remark 11.2. It's interesting to note that the specific dependence of the residues above on f and g is contained entirely in the L-series. This will later be used to simplify some calculations of leading terms.

The residues above have potential polar lines at $2s'+v=2$, $s+1/2-s'=-r$, $s+1/2-s'-v=-r$ $s-3/2+s'=-r$, $s-3/2+s'+v=-r$, where $r \geq 0$. A line can not occur if it intersects the region $\mathfrak{Re}\,w > 1$, $1/2 - k/2 < \mathfrak{Re}\,s < 1$, or the region $\mathfrak{Re}\,s' > 1$, $\mathfrak{Re}\,s \leq 1/2 - k/2$ as we know there are no poles in this region others than those at $s = 1/2 - r + it_j$. (In other words a closer examination must show that it is cancelled.) The line $s - 3/2 + s' = -r$ does, in fact, do this, for any fixed r. This is because as $s' = 3/2 - s - r$, it must be the case that $\mathfrak{Re}\,s' > 1$ for all s with $\mathfrak{Re}\,s < 1/2 - r$. This is similarly the case with $s - 3/2 + s' + v = -r$.

The line $s + 1/2 - s' = -r$ is the same as the line $w = r + 3/2 - k/2$, (as $s' = s + w + k/2 - 1$), and thus to stay out of the region $\mathfrak{Re}\,w > 1$ we require $r \leq k/2 - 1/2$. But any such line will intersect the region $\mathfrak{Re}\,(s + w + k/2) > 5/2$, with $\mathfrak{Re}\,v > 1$, and there are no poles in that region, as the sum over Q of $Z_Q(s,w)$ is absolutely convergent. The line $s+1/2-s'-v = -r$ corresponds to the line $w+v = r+3/2-k/2$, and this has the possibility of occurring if $0 \leq r \leq k/2 - 1$. However, this lies outside of the region in which the spectral expansion is valid.

If $f = g$ there will be a simple pole introduced by the inner product, which, up to a constant, is the completed Rankin-Selberg convolution

$L^*(s'-1/2, f\otimes g)$ and $L^*(s'+v-1/2, f\otimes g)$. This will occur when $s'=1/2$ and when $s'+v=1/2$. Interestingly, this will be sometimes cancelled by a square of a gamma factor in the denominator, but not always. Thus there is another pair of potential polar lines at $s'=1/2$ and $s'+v=1/2$.

By the upper bound in Proposition 5.1, for s at least ϵ away from the polar points and $1/2-k/2-\epsilon < \mathfrak{Re}\, s < 2$, $\mathfrak{Re}\, w > 1$, the sum defining $\mathcal{M}(s,w,v)$ in (11.37) converges absolutely. Finally, the sum converges absolutely in $\mathfrak{Re}\, s > 1, \mathfrak{Re}\, w > 1$. Call the union of these connected regions R_1. Thus

$$\begin{aligned} R_1 = \{(s,w,v) | \mathfrak{Re}\, v > 0\} \cap \{\{(s,w,v)|\mathfrak{Re}\, s \geq -k/2, \mathfrak{Re}\, w > 1,\} \cup \\ \{(s,w,v)| -A < \mathfrak{Re}\, s < -k/2, \mathfrak{Re}\, w > -\mathfrak{Re}\, s/2 - k/4 + 5/4\}\}\,. \end{aligned} \tag{11.45}$$

In particular, in the part of R_1 corresponding to $\mathfrak{Re}\, s < 1/2-k/2$, $\mathfrak{Re}\, s' \geq 1/2$, $\mathfrak{Re}\, v > 0$ and at points at least ϵ away from the poles, after applying Proposition 4.4 and Sterling it follows that $\mathcal{M}(s,w,v)$ obeys the upper bound

$$\mathcal{M}(s,w,v) \ll (\log L)^{1/2} \ell^{-s'} (1+|s|)^{2-2s-k}. \tag{11.46}$$

Our goal is to now relate $\mathcal{M}(s,w,v)$ to an entire function on R_1 by subtracting off the cuspidal polar contribution and then multiplying to clear the polar lines. Specifically, as mentioned above, the poles of $\mathcal{M}(s,w,v)$ arising from the cuspidal contribution occur in the same places as the poles of $D(s;m)$, i.e at the points $s = 1/2 + it_j - r$, for $r \geq 0$. Letting

$$R^{(2)}(1/2-r+it_j, w, v) = \mathrm{Res}_{s=1/2-r+it_j} \mathcal{M}(s,w,v)$$

we compute

$$\begin{aligned} & R^{(2)}(1/2-r+it_j, w, v) \\ &= \frac{2^{2k-1}\pi^k L((k-1/)2 - r + it_j + w, \overline{u_j}) L((k-1)/2 - r + it_j + w + v, \overline{u_j})}{\Gamma(k-1/2-r+it_j)} \\ &\qquad \times \overline{\rho_j(-1))} \overline{\langle V, u_j\rangle} \frac{(-1)^r \Gamma(1/2 - it_j + r)\Gamma(2it_j - r)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)}. \end{aligned} \tag{11.47}$$

Let

$$P^{(2)}(s,w,v) = \sum_{r<A,j} \frac{R^{(2)}(1/2-r+it_j, w, v) e^{(1/2-r+it_j)^2}}{s-(1/2-r+it_j)}. \tag{11.48}$$

The absolutely convergent series $P^{(2)}(s,w,v)$ now has poles and residues at those poles in the region R_1 that are identical to the poles of

$\mathcal{M}(s,w,v)e^{(1/2-r+it_j)^2}$. Thus the function

$$\mathcal{M}(s,w,v)e^{s^2} - P^{(2)}(s,w,v)$$

has only potential polar lines arising from the continuous piece, namely at $2s'+v=2$ and at $s'=1/2$. Thus

$$(s'-1/2)(2s'+v-2)\left(\mathcal{M}(s,w,v)e^{s^2} - P^{(2)}(s,w,v)\right)$$

is holomorphic in R_1.

We will now reverse the roles of s and w in $\mathcal{M}(s,w,v)$ and perform a very similar analysis in the region R_2, defined by

$$\begin{aligned} R_2 = \{(s,w,v) | \mathfrak{Re}\, v > 0, \mathfrak{Re}\, s > 1, \mathfrak{Re}\, w > -B, \\ \mathfrak{Re}\, s + w + k/2 + v/2 > 3/2\}. \quad (11.49)\end{aligned}$$

Recall that in the region $\mathfrak{Re}\, s, w > 1$ and $\mathfrak{Re}\, v \geq 0$ we had the expression for $\mathcal{M}(s,w,v)$ given in (11.3):

$$\mathcal{M}(s,w,v) = (\ell_1\ell_2)^{(k-1)/2} \sum_{m_2 \geq 1} \frac{\bar{b}(m_2)D'(w,v,m_2)}{(\ell_2 m_2)^{s+k-1}}.$$

As for $c < \mathfrak{Re}\, w + v/2 < 2$ $D'(w,v,m_2)$ satisfies the upper bound of (11.12) it follows that $\mathcal{M}(s,w,v)$ converges for $\mathfrak{Re}\, s$ sufficiently large and $c < \mathfrak{Re}\, w + v/2 < 2$. In particular, applying the upper bound (11.10), it follows that for $\mathfrak{Re}\, w + v/2 < 1/2 - k/2$, $\mathfrak{Re}\, s' \geq 1/2$, $\mathfrak{Re}\, v > 0$,

$$\mathcal{M}(s,w,v) \ll \frac{\ell^{(v-k)/2} L^{-s'-v/2+\theta} (\log \ell L)^{1/2} (1+|v|)^{k/2}}{(1+|w|)^{w+k/2-1}(1+|w+v|)^{w+k/2-1+v}}. \quad (11.50)$$

Substituting (11.7) in the above gives us

$$\begin{aligned} \mathcal{M}(s,w,v) = (\ell_1\ell_2)^{(k-1)/2} \sum_{m_2 \geq 1} \frac{\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1}} \\ \times \pi^{(k-v-1)/2} 2^k \ell^{(v-1)/2} (\ell_2 m_2)^{1/2-w-v/2} \\ \times \frac{\Gamma(1-w-v/2)\Gamma(w+v/2)\Gamma((k+1+v)/2)}{\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)} (S_{\text{cusp}} + S_{\text{cont}}) \\ - (\ell_1\ell_2)^{(k-1)/2} \sum_{m_2 \geq 1} \frac{\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1}} \\ \times \frac{\Gamma(1-w-v/2)\Gamma(w+v/2)}{\Gamma((1-v+k)/2)\Gamma((1+v-k)/2} \sum_{1 \leq \ell m' < \ell_2 m_2} \frac{a((\ell_2 m_2 - \ell m')/\ell)\sigma_{-v}(m')}{(\ell m')^{w+(k-1)/2}}, \end{aligned}$$

with $S_{\text{cusp}}, S_{\text{cont}}$ given by (11.8), (11.9).

The sum over m_2 can be brought inside, leading to

$$\begin{aligned}\mathcal{M}(s,w,v) = {} & \ell_1^{(k-1)/2}\ell_2^{1-s-w-k/2-v/2}\ell^{(v-1)/2}2^k\pi^{k/2-1/2-v/2} \\ & \times \frac{\Gamma((k+1+v)/2)\Gamma(w+v/2)}{\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)} \\ & \times \Gamma(1-w-v/2)(\mathcal{M}_1(s,w,v)+\mathcal{M}_2(s,w,v)) \\ & - \ell_1^{(k-1)/2}\ell_2^{1/2-s-k/2}\ell^{1/2-w-k/2}\frac{\Gamma(1-w-v/2)\Gamma(w+v/2)}{\Gamma((1-v+k)/2)\Gamma((1+v-k)/2} \\ & \times \sum_{\substack{m_2\geq 1 \\ 0\leq \ell m'<\ell_2 m_2}} \frac{\bar{b}(m_2)a((\ell_2 m_2-\ell m')/\ell)\sigma_{-v}(m')}{(m')^{w+(k-1)/2}(m_2)^{s+k-1}},\end{aligned} \tag{11.51}$$

where

$$\begin{aligned}\mathcal{M}_1(s,w,v) = {} & \sum_j L_{\ell_2}(s'+v/2,\bar{g}\otimes\bar{u}_j)\overline{\rho_j(-1)\langle U,u_j\rangle} \\ & \times\frac{\Gamma(w+v/2-1/2+it_j)\Gamma(w+v/2-1/2-it_j)}{\Gamma(1/2+it_j)\Gamma(1/2-it_j)},\end{aligned} \tag{11.52}$$

(recall $s'=s+w+k/2-1$) and

$$\begin{aligned}\mathcal{M}_2(s,w,v) = {} & \sum_a\int_{-\infty}^{\infty} L_{a,\ell_2}(s'+v/2+it,\bar{g})L_{a,\ell_2}(s'+v/2-it,\bar{g})\pi^{1/2-it} \\ \times & \frac{\Gamma(w+v/2-1/2+it)\Gamma(w+v/2-1/2-it)}{4\pi\zeta(1-2it)\Gamma(1/2-it)^2\Gamma(1/2+it)}\overline{\langle U,E_a(*,1/2+it)\rangle}dt.\end{aligned} \tag{11.53}$$

In the case of $\mathcal{M}_1$ the series $L_{\ell_2}(s'+v/2,\bar{g}\otimes\bar{u}_j)$ is defined by

$$L_{\ell_2}(s'+v/2,\bar{g}\otimes\bar{u}_j) = \sum_{m_2\geq 1}\frac{\bar{B}(m_2)\bar{\lambda}_j(\ell_2 m_2)}{m_2^{s'+v/2}}$$

and equals the Rankin-Selberg convolution of $\bar{g}$ with $\bar{u}_j$, up to the Euler factors corresponding to primes dividing ℓ_2. The series L_{a,ℓ_2} appearing in $\mathcal{M}_2$ are the L-series associated to $\bar{g}$ up to Euler factors corresponding to primes dividing $\ell_1\ell_2\ell N_0$. For $Re\, v\geq 0$, $\mathfrak{Re}\, w+v/2<1/2-k/2$, and $\mathfrak{Re}\, s+w+k/2+v/2>3/2$ the series and integral $\mathcal{M}_1,\mathcal{M}_2$ converge.

The only possible poles of $\mathcal{M}_2(s,w,v)$ are at $\mathfrak{Re}\, w+v/2=1/2-\ell'$, for some $\ell'\geq 0$. To eliminate this possibility write $w+v/2=1/2-\epsilon-\ell'+i\gamma$, with $\epsilon<(2c'\log(|t|+2))^{-1}$ for some small c'. It follows from (11.35) that if the line of integration is shifted slightly (writing $u=it$ move it to $u=-(c'\log(|t|+2))^{-1}+it$), then a residue at $u=-\epsilon-i\gamma$ will have the form

$$\frac{F(s,w)\Gamma(-2\epsilon+2i\gamma-\ell)}{\zeta(1+2\epsilon-2i\gamma)\zeta(1-2\epsilon+2i\gamma)\Gamma(1/2+\epsilon-i\gamma)\Gamma(1/2-\epsilon+i\gamma)},$$

where $F(s,w)$ is analytic in the region being considered. It follows that to have a pole we must have $\gamma = 0$, and then as $\epsilon \to 0$ the poles in the numerator and denominator cancel.

The sum over m' and m_2 can be transformed and has analytic continuation to $\mathbb{C}$ as follows:

Proposition 11.3. *The Dirichlet series*

$$\begin{aligned}&\mathcal{M}_3(s,w,v)\\ &\quad= \ell_1^{(k-1)/2}\ell_2^{1/2-s-k/2}\ell^{1/2-w-k/2}\frac{\Gamma(1-w-v/2)\Gamma(w+v/2)}{\Gamma((1-v+k)/2)\Gamma((1+v-k)/2}\\ &\qquad\qquad\times \sum_{\substack{m_2\geq 1\\ 0\leq \ell m'<\ell_2 m_2}} \frac{\bar{b}(m_2)a((\ell_2 m_2-\ell m')/\ell)\sigma_{-v}(m')}{(m')^{w+(k-1)/2}(m_2)^{s+k-1}}\end{aligned}$$

is absolutely convergent for $\mathfrak{Re}\, s > 1$ *and has the following meromorphic continuation to* $\mathbb{C}$*:*

$$\begin{aligned}&\mathcal{M}_3(s,w,v)\\ &\quad= \frac{2^{2k-1-v}\pi^k(\ell_1\ell_2)^{(k-1)/2}\Gamma(w+v/2)\Gamma(1-w-v/2)\ell^{-(s+w+k/2-1)}}{\Gamma((1+v-k)/2)\Gamma((1-v+k)/2)\Gamma(s)\Gamma(s+k-1)\mathcal{V}\zeta(2s'+v)}\\ &\times\sum_{j\geq 1} L(s',\bar{u}_j)L(s'+v,\bar{u}_j)\Gamma(s-1/2+it_j)\Gamma(s-1/2-it_j)\overline{\rho_j(1)}\left\langle u_j, V\right\rangle\\ &+2^{2k-2}\pi^{k-1/2}\int_{-\infty}^{\infty}\frac{\pi^{-it}\zeta_a(s'+it)\zeta_a(s'-it)\zeta_a(s'+v+it)\zeta_a(s'+v-it)}{\zeta(2s'+v)\zeta(1-2it)\Gamma(1/2-it)\Gamma(s)\Gamma(s+k-1)}\\ &\times \ell^{-(s+w+k/2-1)}\Gamma(s-1/2+it)\Gamma(s-1/2-it)\left\langle E(*,1/2+it), V\right\rangle dt.\end{aligned}$$

The only poles in the region R_2*, defined in* (11.49)*, are at* $w+v/2 = -r$*,* $r \geq 0$*.*

Proof. The inner sum over m_2 is essentially an inner product of f and g with a standard Poincare series. In particular, if

$$P(s,z,m') = \sum_{\gamma\in\Gamma_\infty\backslash\Gamma} (\mathfrak{Im}\,\gamma z)^s e^{2\pi i\ell m'\gamma z},$$

then

$$\Gamma(s+k-1)\sum_{m_2\geq 1}\frac{a((\ell_2 m_2-\ell m')/\ell_1)\bar{b}(m_2)}{(4\pi\ell_2 m_2)^{s+k-1}} = \left\langle P(s,z,m'), \bar{f}gy^k\right\rangle.$$

The inner product can also be expressed in terms of a spectral expansion

$$\left\langle P(s,*,m'), \bar{f}gy^k \right\rangle = \sum_j \langle P(s,*,m'), u_j\rangle \left\langle u_j, \bar{f}gy^k \right\rangle$$

$$+ \frac{1}{4\pi}\sum_a \int_{-\infty}^{\infty} \langle P(s,*,m'), E_a(*,1/2+it)\rangle \left\langle E_a(*,1/2+it), \bar{f}gy^k \right\rangle .$$

A simple computation shows that

$$\begin{aligned}\langle P(s,*,m'), u_j\rangle &= \int_0^\infty \frac{y^{s-1}\overline{\rho_j(\ell m')}e^{-y}K_{it_j}(y)}{(2\pi\ell m')^{s-1/2}}\frac{dy}{y}\\ &= \frac{\overline{\rho_j(m')}\sqrt{\pi}2^{1/2-s}\Gamma(s-1/2+it_j)\Gamma(s-1/2-it_j)}{(2\pi\ell m')^{s-1/2}\Gamma(s)}\end{aligned}$$

and that

$$\begin{aligned}&\langle P(s,*,m'), E_a(*,1/2+it)\rangle\\ &= \frac{2^{2-2s}\pi^{3/2-s-it}(\ell m')^{1/2-s-it}\overline{\tilde{\rho}_a(1/2+it,\ell m'_a)}\Gamma(s-1/2+it)\Gamma(s-1/2-it)}{\Gamma(1/2-it)\zeta(1-2it)\Gamma(s)}.\end{aligned}$$

Substituting these evaluations leads to

$$\begin{aligned}&\sum_{j.m'\geq 1}\frac{\sigma_{-v}(m')}{(\ell m')^{w+(k-1)/2}}\sum_{m_2\geq 1}\frac{a((\ell_2 m_2-\ell m')/\ell_1)\bar{b}(m_2)}{(\ell_2 m_2)^{s+k-1}}\\ &= 2^{2k-1}\pi^k\sum_{m'\geq 1}\frac{\overline{\rho_j(\ell m')}\Gamma(s-1/2+it_j)\Gamma(s-1/2-it_j)\left\langle u_j,\bar{f}gy^k\right\rangle\sigma_{-v}(m')}{(\ell m')^{s+w+k/2-1}\Gamma(s)\Gamma(s+k-1)}\end{aligned}$$

plus the corresponding continuous contribution. The cuspidal part may contribute simple poles at $w+v/2=-r$, $r\geq 0$. (The factor $\Gamma(1-w-v/2)$ does not contribute a pole as $\mathfrak{Re}\, w+v/2<1$ in R_3 does not occur in R_2.) The poles of the continuous contribution to the $m'\geq 1$ piece are analyzed in an identical way to those of the continuous contribution in the region $\mathfrak{Re}\, s<1$, and there are polar lines at the same locations, none of which lie in R_2. Similarly, the poles of the cuspidal part at $s=1/2+it_j-r$ lie outside of R_2. □

We have shown that the function

$$(s'-1/2)(2s'+v-2)\left(\mathcal{M}(s,w,v)e^{s^2}-P^{(2)}(s,w,v))\right)$$

is holomorphic in the region R_1. We will now identify and remove the poles of $\mathcal{M}(s,w,v)$ in the region R_2. By the previous discussion, the expression for $\mathcal{M}$ given in (11.51) converges absolutely in the part of R_2

with $\mathfrak{Re}\, w+v/2<1$, when a distance of ϵ from the polar points. There are no other poles in the rest of R_2 as the series expansion converges absolutely there. Let κ run over the poles of $\mathcal{M}(s,w,v)$ in R_2 and set

$$R^{(3)}(s,\kappa,v)=\operatorname{Res}_{w=\kappa}\mathcal{M}(s,w,v). \tag{11.54}$$

Referring to (11.51) - (11.53) and Proposition 11.3, we see that these poles are at the following points:

- $w+v/2=1/2+it_j-r$ (from $\mathcal{M}_1$)
- The points $w+v/2=-r$, with $B>r\geq 0$. (From $\Gamma(w+v/2)$).

Writing

$$e_{j,r} \quad == \quad \operatorname{Res}_{w+v/2=1/2+it_j-r}\mathcal{M}(s,w,v),$$

we find that

$$\begin{aligned} e_{j,r} &= \frac{\Gamma((k+1+v)/2)\Gamma(1/2+it_j-r)\Gamma(1/2-it_j+r)}{\Gamma(1/2+it_j-r-v/2)\Gamma(1/2+it_j-r+v/2)} \\ &\times \frac{(-1)^r\Gamma(2it_j-r)\overline{\rho_j(-1)\langle U,u_j\rangle}L_{\ell_2}(s+k/2-1/2-r+it_j,\bar g\otimes \bar u_j)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)}. \end{aligned} \tag{11.55}$$

Let

$$P^{(3)}(s,w,v)=\sum_{\kappa}\frac{R^{(3)}(s,\kappa,v)e^{\kappa^2}}{w-\kappa}, \tag{11.56}$$

where the κ vary over the poles listed above. This series has poles and residues at poles in the region R_2 that are identical to those of $\mathcal{M}(s,w,v)$ and converges absolutely. Thus the function

$$\mathcal{M}(s,w,v)e^{w^2}-P^{(3)}(s,w,v)$$

is holomorphic in R_2.

The function $P^{(3)}(s,w,v)$ has no poles in R_1 and the function $P^{(2)}(s,w,v)$ has no poles in R_2. Thus

$$\begin{aligned} &(s'-1/2)(2s'+v-2) \\ &\times\Big(\mathcal{M}(s,w,v)e^{s^2+w^2}-P^{(2)}(s,w,v)e^{w^2}-P^{(3)}(s,w,v)e^{s^2}\Big). \end{aligned} \tag{11.57}$$

It follows that the difference in (11.57) is holomorphic in $R_1\cup R_2$. As $A,B\to\infty$ the convex hull of $R_1\cup R_2$ includes all of $\mathbb{C}^3$ such that $\mathfrak{Re}\, v>0$. Applying the theorem of Bochner, [**?**], we have completed the proof of the meromorphic continuation of $\mathcal{M}(s,w,v)$. Summarizing the above, we have proved

Proposition 11.4. *The multiple Dirichlet series*

$$\mathcal{M}(s,w,v)=\sum_{\substack{m_2,h_0,Q\geq 1\\ \ell_1 m_1=\ell_2 m_2+m}}\frac{A(m_1)\bar{B}(m_2)\left(1+\frac{m}{\ell_2 m_2}\right)^{(k-1)/2}\sigma_{-v}(m)}{(\ell_2 m_2)^s m^{w+(k-1)/2}}$$

converges absolutely when $\mathfrak{Re}\, s,w>1,\mathfrak{Re}\, v>0$ *and has a meromorphic continuation to the region* $\{(s,w,v)|\mathfrak{Re}\, v>0\}$. *It is analytic in this region, except (possibly) for the following simple polar planes:*

- $s=1/2\pm it_j-r$ *for* $r\geq 0$,
- $w+v/2=1/2\pm it_j-r$ *for* $r\geq 0$,
- $w+v/2=-r$, *for* $r\geq 0$,
- *s' = 1/2*
- $2s'+v=2$

The residues of $\mathcal{M}(s,w,v)$ *at* $s=1/2\pm it_j-r$ *and at* $w=1/2-v/2\pm it_j-r$ *are given by*

$$\begin{aligned}\operatorname{Res}_{s=1/2-r+it_j}\mathcal{M}(s,w,v)&=\frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell^{(k-1)/2-r+it_j+w}}\mathcal{P}_\ell(k-1)/2-r+it_j+w,v)\\ \times\frac{2^{2k-1}\pi^k L((k-1)/2-r+it_j+w,\overline{u_j})L((k-1)/2-r+it_j+w+v,\overline{u_j})}{\zeta(k-1-2r+2it_j+2w+v)\Gamma(k-1/2-r+it_j)}&\\ \times\overline{\rho_j(-1))}\overline{\langle V,u_j\rangle}\frac{(-1)^r\Gamma(1/2-it_j+r)\Gamma(2it_j-r)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)}&.\end{aligned}\tag{11.58}$$

$$\begin{aligned}&\operatorname{Res}_{w=1/2+it_j-r-v/2}\mathcal{M}(s,w,v)\\ &\qquad=\ell_1^{(k-1)/2}\ell_2^{1/2-s-it_j+r-k/2}\ell^{(v-1)/2}2^k\pi^{k/2-1/2-v/2}\\ &\qquad\frac{\Gamma((k+1+v)/2)\Gamma(1/2+it_j-r)\Gamma(1/2-it_j+r)}{\Gamma(k/2+1/2+it_j-r-v/2)\Gamma(k/2+1/2+it_j-r+v/2)}\\ &\times\frac{(-1)^r\Gamma(2it_j-r)\overline{\rho_j(-1)\langle U,u_j\rangle}L_{\ell_2}(s+k/2-1/2-r+it_j,\bar{g}\otimes\bar{u}_j)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)}.\end{aligned}\tag{11.59}$$

When $\mathfrak{Re}\, v\geq 0$, $-1/2-k/2<\mathfrak{Re}\, s<1/2-k/2$ *and* $\mathfrak{Re}\, w>1$, *and also when* $\mathfrak{Re}\, v\geq 0$, $\mathfrak{Re}\, s\leq -1/2-k/2$ *and* $w>-s/2-k/4+5/4$, $\mathcal{M}(s,w,v)$ *is represented by*

$$\mathcal{M}(s,w,v)=\frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell^{s+w+k/2-1}}\left(\mathcal{M}_{cusp}+\mathcal{M}_{cont}\right),$$

with

$$\mathcal{M}_{cusp} = \frac{2^{2k-1}\pi^k\Gamma(1-s)}{\Gamma(s+k-1)}\sum_j \tag{11.60}$$

$$\frac{L(s',\overline{u_j})L(s'+v,\overline{u_j})\mathcal{P}(s',v)\Gamma(s-\frac{1}{2}-it_j)\Gamma(s-\frac{1}{2}+it_j)\overline{\rho_j(-1)\langle V,u_j\rangle}}{\zeta(2s'+v)\Gamma(\frac{1}{2}-it_j)\Gamma(\frac{1}{2}+it_j)}$$

and

$$\begin{aligned}\mathcal{M}_{cont} = \frac{2^{2k-1}\pi^k\Gamma(1-s)}{4\pi\Gamma(s+k-1)}\sum_a\int_{-\infty}^{\infty}\zeta_a(s'+it)\zeta_a(s'-it)\zeta_a(s'+v+it)\\ \times\quad \frac{\zeta_a(s'+v-it)\mathcal{P}(s',v,it)\Gamma(s-\frac{1}{2}-it)\Gamma(s-\frac{1}{2}+it)\overline{\langle V,E_a^*(*,\frac{1}{2}+it)\rangle}}{\mathcal{V}\zeta^*(1+2it)\zeta(1-2it)\zeta(2s'+v)\pi^{it}\Gamma(\frac{1}{2}-it)^2\Gamma(\frac{1}{2}+it)}dt,\end{aligned} \tag{11.61}$$

When $\mathfrak{Re}\, v \geq 0$*,* $\mathfrak{Re}\, s > 1$*,* $\mathfrak{Re}\, w + v/2 < 1/2 - k/2$*, and* $\mathfrak{Re}\, s + w + k/2 + v/2 > 3/2$*,* $\mathcal{M}(s,w,v)$ *is represented by*

$$\begin{aligned}\mathcal{M}(s,w,v) = \Big(&\ell_1^{(k-1)/2}\ell_2^{1-s-w-k/2-v/2}\ell^{(v-1)/2}2^k\pi^{k/2-1/2-v/2}\\ &\times\frac{\Gamma((k+1+v)/2)\Gamma(w+v/2)}{\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)}\\ &\times\Gamma(1-w-v/2)(\mathcal{M}_1(s,w,v)+\mathcal{M}_2(s,w,v))\Big)\\ &\qquad\qquad\qquad - \mathcal{M}_3(s,w,v),\end{aligned}$$

where

$$\begin{aligned}\mathcal{M}_1(s,w,v) = \sum_j L_{\ell_2}(s+w+k/2-1+v/2,\bar{g}\otimes\bar{u}_j)\overline{\rho_j(-1)\langle U,u_j\rangle}\\ \times\frac{\Gamma(w+v/2-1/2+it_j)\Gamma(w+v/2-1/2-it_j)}{\Gamma(1/2+it_j)\Gamma(1/2-it_j)},\end{aligned}$$

$$\begin{aligned}\mathcal{M}_2(s,w,v) = \sum_a\int_{-\infty}^{\infty}L_{a,\ell_2}(s'+v/2+it,\bar{g})L_{a,\ell_2}(s'+v/2-it,\bar{g})\pi^{1/2-it}\\ \times\frac{\Gamma(w+v/2-1/2+it)\Gamma(w+v/2-1/2-it)}{4\pi\zeta(1-2it)\Gamma(1/2-it)^2\Gamma(1/2+it)}\overline{\langle U,E_a(*,1/2+it)\rangle}dt,\end{aligned}$$

and

$$\begin{aligned}
&\mathcal{M}_3(s,w,v)\\
&\quad= \frac{2^{2k-1-v}\pi^k(\ell_1\ell_2)^{(k-1)/2}\Gamma(w+v/2)\Gamma(1-w-v/2)\ell^{-(s+w+k/2-1)}}{\Gamma((1+v-k)/2)\Gamma((1-v+k)/2)\Gamma(s)\Gamma(s+k-1)\mathcal{V}\zeta(2s'+v)}\\
&\times\sum_{j\geq 1} L(s',\bar{u}_j)L(s'+v,\bar{u}_j)\Gamma(s-1/2+it_j)\Gamma(s-1/2-it_j)\overline{\rho_j(1)}\left\langle u_j,V\right\rangle\\
&+2^{2k-2}\pi^{k-1/2}\int_{-\infty}^{\infty}\frac{\pi^{-it}\zeta_a(s'+it)\zeta_a(s'-it)\zeta_a(s'+v+it)\zeta_a(s'+v-it)}{\zeta(2s'+v)\zeta(1-2it)\Gamma(1/2-it)\Gamma(s)\Gamma(s+k-1)}\\
&\times\ell^{-(s+w+k/2-1)}\Gamma(s-1/2+it)\Gamma(s-1/2-it)\left\langle E(*,1/2+it),V\right\rangle dt.
\end{aligned}$$

For points at least ϵ away from the poles, the following polynomial upper bounds are satisfied: when $\mathfrak{Re}\, s<1/2-k/2$, $\mathfrak{Re}\, s'\geq 1/2$, $\mathfrak{Re}\, v>0$,

$$\mathcal{M}(s,w,v)\ll(\log L)^{1/2}\ell^{-s'}(1+|s|)^{2-2s-k}.$$

and when $\mathfrak{Re}\, w+v/2<1/2-k/2$, $\mathfrak{Re}\, s'\geq 1/2$, $\mathfrak{Re}\, v>0$,

$$\mathcal{M}(s,w,v)\ll\frac{\ell^{(v-k)/2}L^{-s'-v/2+\theta}(\log\ell L)^{1/2}(1+|v|)^{k/2}}{(1+|w|)^{w+k/2-1}(1+|w+v|)^{w+k/2-1+v}}.$$

12. AVERAGING MEAN VALUES OF L-SERIES OVER Q

12.1. **A mean value estimate for a single Q.** For $x\gg 1$ and any $n\geq 0$ let

$$\begin{aligned}
I_Q(n)=\left(\frac{1}{2\pi i}\right)^2\int_{(2+k/2)}\int_{(1+n)}&\frac{\Gamma(s+k/2)\Gamma(s+w)\Gamma(w)}{\Gamma(s+w+k/2)}x^{s+w-n}\\
&\times Z_Q(s+1-n,w+(1-k)/2)dsdw \quad (12.1)
\end{aligned}$$

Proposition 12.1. *Then*

$$I_Q(n) \;=\; \sum_{\substack{m_2,h_0\geq 1\\ \ell_1m_1=\ell_2m_2+h_0Q}}\left(\frac{\ell_2m_2}{x}\right)^n\frac{A(m_1)\bar{B}(m_2)}{(\ell_1m_1)^{1/2}(\ell_2m_2)^{1/2}}e^{-(\ell_2m_2+h_0Q)/x}. \tag{12.2}$$

Proof. Recalling the definition of $Z_Q(s,w)$ in (7.4), write

$$\begin{aligned}
Z_Q(s+1, w+(1-k)/2) &= \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \frac{A(m_1)\bar{B}(m_2)\left(1+\frac{h_0 Q}{\ell_2 m_2}\right)^{(k-1)/2}}{(\ell_2 m_2)^{s+1}(h_0 Q)^w} \\
&= \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \frac{A(m_1)\bar{B}(m_2)\left(1+\frac{h_0 Q}{\ell_2 m_2}\right)^{k/2}}{(\ell_1 m_1)^{1/2}(\ell_2 m_2)^{1/2}(\ell_2 m_2)^{s}(h_0 Q)^w} \\
&= \sum_{R=0}^{k/2}\binom{k/2}{R} \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \frac{A(m_1)\bar{B}(m_2)}{(\ell_1 m_1)^{1/2}(\ell_2 m_2)^{1/2}(\ell_2 m_2)^{s+R}(h_0 Q)^{w-R}}
\end{aligned} \tag{12.3}$$

Let $I_Q(n)$ denote the left hand side of (12.1) above. Substituting (12.3) and interchanging the sum over R and the integrals gives

$$I_Q(n) = \sum_{R=0}^{k/2}\binom{k/2}{R}\left(\frac{1}{2\pi i}\right)^2 \int_{(2+k/2)}\int_{(2+n)} \frac{\Gamma(s+k/2)\Gamma(s+w)\Gamma(w)}{\Gamma(s+w+k/2)} x^{s+w}$$
$$\times \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \left(\frac{\ell_2 m_2}{x}\right)^n \frac{A(m_1)\bar{B}(m_2)}{(\ell_1 m_1)^{1/2}(\ell_2 m_2)^{1/2}(\ell_2 m_2)^{s+R}(h_0 Q)^{w-R}} ds dw.$$

Changing variables $s \to s-R$, $w \to w+R$ and moving the lines of integration we obtain

$$\begin{aligned}
I_Q(n) &= \sum_{R=0}^{k/2}\binom{k/2}{R}\left(\frac{1}{2\pi i}\right)^2 \int_{(2)}\int_{(2+n)} \frac{\Gamma(s+k/2-R)\Gamma(s+w)\Gamma(w+R)}{\Gamma(s+w+k/2)} x^{s+w} \\
&\times \sum_{\substack{m_2,h_0\geq 1\\ \ell_1 m_1=\ell_2 m_2+h_0 Q}} \left(\frac{\ell_2 m_2}{x}\right)^n \frac{A(m_1)\bar{B}(m_2)}{(\ell_1 m_1)^{1/2}(\ell_2 m_2)^{1/2}(\ell_2 m_2)^{s}(h_0 Q)^{w}} ds dw.
\end{aligned} \tag{12.4}$$

The sum over R can be simplified as described in the following

Lemma 12.2.

$$\Gamma(s)\Gamma(w) = \frac{\Gamma(s+w)}{\Gamma(s+w+k/2)}\sum_{R=0}^{k/2}\binom{k/2}{R}\Gamma(s+k/2-R)\Gamma(w+R)$$

Proof. Substituting the integral expressions for the gamma factors inside the summation, the right hand side above equals

$$\frac{\Gamma(s+w)}{\Gamma(s+w+k/2)}\sum_{R=0}^{k/2}\binom{k/2}{R}\int_0^\infty\int_0^\infty x_1^{s+k/2-R}x_2^{w+R}e^{-x_1-x_2}\frac{dx_1}{x_1}\frac{dx_2}{x_2}.$$

Setting $x_2=x_1x_3$ and summing over R inside the integrals we obtain

$$=\frac{\Gamma(s+w)}{\Gamma(s+w+k/2)}\int_0^\infty\int_0^\infty x_1^{s+k/2+w}x_3^w(1+x_3)^{k/2}e^{-x_1(1+x_3)}\frac{dx_1}{x_1}\frac{dx_3}{x_3}.$$

The integration over x_1 yields $\Gamma(s+w+k/2)$, canceling the corresponding term in the denominator, leaving

$$\Gamma(s+w)\int_0^\infty(1+x_3)^{-s-w}x_3^w\frac{dx_3}{x_3}.$$

Substituting the integral expressions for $\Gamma(s+w)$ we have

$$\int_0^\infty\int_0^\infty x_4^{s+w}e^{-x_4}(1+x_3)^{-s-w}x_3^w\frac{dx_3}{x_3}\frac{dx_4}{x_4},$$

and the lemma follows after writing $x_4=x_5(1+x_3)$ and integrating. ☐

Applying Lemma 12.2 to (12.4) we obtain

$$I_Q(n)=\left(\frac{1}{2\pi i}\right)^2\int_{(2)}\int_{(2)}\Gamma(s)\Gamma(w)x^{s+w}$$
$$\times\sum_{\substack{m_2,h_0\geq1\\ \ell_1m_1=\ell_2m_2+h_0Q}}\left(\frac{\ell_2m_2}{x}\right)^n\frac{A(m_1)\bar{B}(m_2)}{(\ell_1m_1)^{1/2}(\ell_2m_2)^{1/2}(\ell_2m_2)^s(h_0Q)^w}dsdw.$$

Upon interchanging the sum and the integrals and then integrating, the proposition follows. ☐

The following proposition shows how the analytic properties of $Z_Q(s,w)$ can be used to understand the sum on the right hand side of (12.1).

Proposition 12.3. *For $x\gg1$ and $n\geq0$*

$$\sum_{\substack{m_2,h_0\geq1\\ \ell_1m_1=\ell_2m_2+h_0Q}}\left(\frac{\ell_2m_2}{x}\right)^n\frac{A(m_1)\bar{B}(m_2)}{(\ell_1m_1)^{1/2}(\ell_2m_2)^{1/2}}e^{-(\ell_2m_2+h_0Q)/x}$$
$$=\begin{cases}c(f)\log x+c'(f) & \text{if } n=0 \text{ and } f=g\\ Z_Q(1-k/2,1/2) & \text{if } n=0 \text{ and } f\neq g\end{cases}$$
$$+\mathcal{O}\left(Q^{\theta-1/2}\right)+\mathcal{O}\left(Q^\theta x^{-1/2}\right)$$
$$+\mathcal{O}\left(Q^{\theta-1/2}\log(L)x^{-1/2}\right)+\mathcal{O}\left(Q^Ax^{-C}\right).$$

The constants are $c(f) = c_{1,f} + c_{3,f}, c'(f) = c_{2,f} + c_{4,f}$,

$$\begin{aligned} &c_{1,f} \log x + c_{2,f} \\ &= -\operatorname{Res}_{w=1+k/2} \frac{\Gamma(w-k/2-1)\Gamma(w)}{\Gamma(w-1)} x^{w-k/2-1} Z_Q(-k/2, w+(1-k)/2) \end{aligned}$$

$$c_{3,f} \log x + c_{4,f} = \operatorname{Res}_{w=k/2} \Gamma(w-k/2) x^{w-k/2} Z_Q(1-k/2, w+(1-k)/2) \tag{12.5}$$

Remark 12.4. Note that the main term vanishes if $n \neq 0$. (Though there is a second order main term.) When $n = 0$ the The key question left unresolved is the order of magnitude of $Z_Q(1-k/2, 1/2)$ if $f \neq g$, and of $Z_Q(s, w)$ near this point if $f = g$. This point is in the analog, in this two dimensional case, of the center of the critical strip. We will show later that the value is constant on average as Q varies.

Proof. Move the line of the s integral to $\mathfrak{Re}\, s+1-n = -1/2-k/2+\epsilon$. In the process, simple poles are encountered at $s+1-n = 1/2-r+it_j$, for $0 \leq r \leq k/2$, and also at $s = -k/2, -k/2-1$. The pole at $s = -k/2$ is only encountered if $n = 0, 1$, that at $-k/2-1$ only at $n = 0$. Recalling Proposition 7.1, the residue at $s+1-n = 1/2-r+it_j$ is

$$\begin{aligned} \frac{1}{2\pi i} \int_{(2+k/2)} & \frac{\Gamma(n-1/2-r+it_j+k/2)\Gamma(n-1/2-r+it_j+w)\Gamma(w)}{\Gamma(n-1/2-r+it_j+w+k/2)} \\ & \times x^{w-1/2-r+it_j} (\ell_1\ell_2)^{(k-1)/2} c_{r,j} L_Q(w-r+it_j, \overline{u_j}; 0)) dw \end{aligned}$$

where

$$L_Q(s', \overline{u_j}; 0) \qquad = \qquad Q^{-s'} L(s', \overline{u_j}) \prod_{p^\gamma || Q} \left(\overline{\lambda_j(p^\gamma)} - \overline{\lambda_j(p^{\gamma-1})} p^{-s'} \right)$$

Move the w line of this residual integral to $\mathfrak{Re}\, w = r$. In the case $n = 0$ a pole is passed over at $w = 1/2+r-it_j$ with residue

$$\frac{\Gamma(-1/2-r+it_j+k/2)\Gamma(1/2+r-it_j)}{\Gamma(k/2)} (\ell_1\ell_2)^{(k-1)/2} c_{r,j} L_Q(1/2, \overline{u_j}; 0))$$

Referring to (7.32), (7.33), we see that the sum over t_j of these residues is

$$\mathcal{O}\left(Q^{\theta-1/2}\right). \tag{12.6}$$

The same reference shows that the moved integral contributes at most

$$\mathcal{O}\left(LQ^{\theta} x^{-1/2}\right). \tag{12.7}$$

The residual integral at $s = -k/2$, when $n = 0, 1$, is

$$\frac{1}{2\pi i}\int_{(2+k/2)} \Gamma(w-k/2)x^{w-n-k/2} \\ \times Z_Q(1-k/2-n, w+(1-k)/2)dw \tag{12.8}$$

The residual integral at $s = -k/2-1$, when $n = 0$, is

$$\frac{-1}{2\pi i}\int_{(2+k/2)} \frac{\Gamma(w-k/2-1)\Gamma(w)}{\Gamma(w-1)}x^{w-k/2-1} \\ \times Z_Q(-k/2, w+(1-k)/2)dw \tag{12.9}$$

If $n = 1$, move the line of integration in (12.8) to $\mathfrak{Re}\, w = 1/2 + k/2$. A simple pole is passed over at $w = 1 + k/2$, (if $f = g$), and referring to the bound (7.31) we see that the expression (12.8) is

$$\delta_{f,g}\operatorname{Res}_{w=3/2} Z_Q(-k/2, w) + \mathcal{O}\left(Q^{\theta-1/2}\log(L)x^{-1/2}\right). \tag{12.10}$$

If $n = 0$, move the line of integration in (12.9) to $\mathfrak{Re}\, w = 1/2 + k/2$. A pole is passed over at $w = 1 + k/2$ because of the contribution of $\Gamma(w - k/2 - 1)$, but it is now a double pole if $f = g$. As a result a factor of $\log x$, is introduced in that case, yielding the following bound for (12.9):

$$c_{1,f}\log x + c_{2,f} + \mathcal{O}\left(Q^{\theta-1/2}\log(L)x^{-1/2}\right). \tag{12.11}$$

where

$$c_{1,f}\log x + c_{2,f} \\ = -\operatorname{Res}_{w=1+k/2}\frac{\Gamma(w-k/2-1)\Gamma(w)}{\Gamma(w-1)}x^{w-k/2-1}Z_Q(-k/2, w+(1-k)/2) \tag{12.12}$$

and $c_{1,f} = 0$ if $f \neq g$.

If $n = 0$, move the line of integration in (12.8) to $\mathfrak{Re}\, w = k/2 - 1/2$. In the process a pole is passed over at $w = k/2$, and this becomes a double pole if $f = g$. The residue at this pole is the special value $Z_Q(1 - k/2, 1/2)$, if $f \neq g$, and we set

$$c_{3,f}\log x + c_{4,f} = \operatorname{Res}_{w=k/2}\Gamma(w-k/2)x^{w-k/2}Z_Q(1-k/2, w+(1-k)/2) \tag{12.13}$$

if $f = g$. The shifted integral contributes the same error term as above.

As $Z_Q(s,w)$ has at most polynomial growth in Q, s and w away from poles, it follows that the moved integral above, and the final double moved integral are on the order of $Q^A x^{-C}$, for positive constants A, C which we need not estimate at this time. This completes the proof of the proposition. $\square$

Recall that we have shown

$$\begin{aligned} S(Q) &= L_Q(1, f\otimes g) \\ &+\sum_{d|Q}\frac{\mu(d)}{d}\left(Z_{Q/d}(1-k/2,1/2;F_d,G_d)+Z_{Q/d}(1-k/2,1/2;G_d,F_d)\right) \\ &\qquad\qquad + \mathcal{O}\left(Q^{\theta-1/2}\right) \qquad (12.14)\end{aligned}$$

when $f\neq g$ and

$$S(Q)=c_f(Q)+2\sum_{d|Q}\frac{\mu(d)c_2(F_d,Q)}{d}+\mathcal{O}\left(Q^{\theta-1/2}\right)$$

when $f=g$. We will now begin to average over Q, beginning with the case $f\neq g$.

12.2. **The case $f \neq g$.** Fix $\epsilon>0$ and $X, y\gg 1$, with $y \ll X^{1-\epsilon}$, and consider

$$S(X,y)=\sum_{Q\geq 1}S(Q)e^{-y^2(\log(X/Q))^2/(4\pi)}.$$

Applying the identity

$$ye^{-y^2(\log X)^2/(4\pi)}=-i\int_{(2)}e^{\pi v^2/y^2}X^v dv, \qquad (12.15)$$

to (12.14), we obtain

$$S(X,y)=S_1(X,y)+S_2(X,y)+S_3(X,y)+\mathcal{O}\left(\frac{X^{1/2}}{y}\right),$$

where

$$S_1(X,y)=-i\int_{(2)}\frac{e^{\pi v^2/y^2}X^v}{y}\sum_{Q\geq 1}\frac{L_Q(1,f\otimes g)}{Q^v}dv, \qquad (12.16)$$

$$\begin{aligned} &S_2(X,y) \\ &\quad=-i\int_{(2)}\frac{e^{\pi v^2/y^2}X^v}{y}\sum_{Q\geq 1}\frac{\sum_{d|Q}\frac{\mu(d)}{d}Z_{Q/d}(1-k/2,1/2;F_d,G_d)}{Q^v}dv, \end{aligned}$$

(12.17)

and

$$\begin{aligned} &S_3(X,y)\\ &\qquad = -i\int_{(2)} \frac{e^{\pi v^2/y^2}X^v}{y}\sum_{Q\geq 1}\frac{\sum_{d|Q}\frac{\mu(d)}{d}Z_{Q/d}(1-k/2,1/2;G_d,F_d)}{Q^v}dv. \end{aligned}$$

The simplest of these is $S_1(X,y)$. The sum over Q in (12.16) can be analyzed as follows.

$$\begin{aligned} &\sum_{Q\geq 1}\frac{L_Q(1,f\otimes g)}{Q^v}\\ &\qquad = L(1,f\otimes g)\sum_{Q\geq 1}\prod_{p|Q}\left(\sum_{\gamma\geq 0}\frac{A(p^\gamma)\bar{B}(p^\gamma)}{p}\right)\left(1-p^{-2}\right)^{-1}\left(1-p^{-2}\right)\\ &\qquad\qquad = \frac{\zeta(v)\zeta(2)L(1,f\otimes g)}{L(1+v,f)L(1+v,\bar{g})}E(v,f,g), \end{aligned}$$

where $E(v,f,g)$ is analytic for $\mathfrak{Re}\, v > -1$.

Substituting in (12.16) gives

$$\begin{aligned} S_1(X,y) = 2\pi &\frac{e^{\pi/y^2}X}{y}\frac{\zeta(2)L(1,f\otimes g)}{L(1,f)L(1,\bar{g})}E(0,f,g)\\ &- i\int_{(2)}\frac{e^{\pi v^2/y^2}X^v}{y}\frac{\zeta(v)\zeta(2)L(1,f\otimes g)}{L(1+v,f)L(1+v,\bar{g})}E(v,f,g)dv, \end{aligned}$$

The integral is estimated as $\mathcal{O}(X^\epsilon)$ by changing variables $v\to yv$ and applying stationary phase to save a power of $y^{1/2}$ from the ratio of gamma factors relating $\zeta(yv)$ to $\zeta(1-yv)$. Thus

$$S_1(X,y) \;=\; 2\pi\frac{e^{\pi/y^2}X}{y}\frac{\zeta(2)L(1,f\otimes g)}{L(1,f)L(1,\bar{g})}E(0,f,g)\;+\;\mathcal{O}(X^\epsilon). \quad (12.18)$$

Moving to $S_2(X,y)$, replace the Q by $Q'd$ in (12.17) and interchange the sums. This leads to

$$\begin{aligned} &S_2(X,y)\\ &\quad = -i\int_{(2)}\frac{e^{\pi v^2/y^2}X^v}{y}\sum_{Q',d\geq 1}\frac{\mu(d)}{d^{1+v}}\frac{Z_{Q'}(1-k/2,1/2;F_d,G_d)}{(Q')^v}dv, \quad (12.19) \end{aligned}$$

Let

$$I(X,y) = I(X,y;\ell_1,\ell_2,\ell) = -i\int_{(2)}\frac{e^{\pi v^2/y^2}X^v}{y}\mathcal{M}(1-k/2,1/2,v)dv. \quad (12.20)$$

Thus $S_2(X,y)$ is a sum of $I(X,y)$ over various values of the parameters ℓ_1, ℓ_2, ℓ.

There is one subtle point here. The following analysis will all be done for a fixed set of these parameters. We will do this by breaking the sum into several pieces and analyzing these separately. We will have to be certain that for each of these pieces the sum in (12.19) converges absolutely. Fortunately the sum over d is easily understandable and will converge absolutely for $\mathfrak{Re}\, v > 0$ for the following reason. Using the Hecke relations, one sees that for fixed d,

$$F_d(z) = \sum_{d=r_1r_2} A(r_1)\mu(r_2)f(r_2z).$$

Thus the inner product $< F_d\bar{G}_d y^k, u_j >$ breaks up into a linear combination of terms involving $f(r_3z)\bar{g}(r_4z)$ for all $r_3, r_4|d$. Each corresponding inner product is bounded above by $d^{-k+1/2}$. For each d there will be an extra factor of d introduced from the multiplicity of u_j of level d in a unit interval, but the $-k$ in the exponent dominates.

The function $\mathcal{M}(1-k/2, 1/2, v)$ has one pole in v at the point $s' + v/2 = 1$, i.e at $v = 1$, as $s' = (1-k/2)+(1/2)+k/2-1 = 1/2$. Moving the line of integration to $\mathfrak{Re}\, v = \epsilon$, we pass over this pole, obtaining

$$\begin{aligned} I(X,y) = \operatorname{Res}_{v=1} \mathcal{M}(1-k/2,1/2,v)\frac{e^{\pi/y^2}X}{y} \\ - i\int_{(\epsilon)} \frac{e^{\pi v^2/y^2}X^v}{y}\mathcal{M}(1-k/2,1/2,v)dv. \quad (12.21) \end{aligned}$$

To analyze the shifted integral, we will relate $\mathcal{M}(1-k/2,1/2,v)$ to an integral around a boundary as follows. Note first that most of the residues contained in our contour will be from the cuspidal part, with remaining poles at $w = -r - v/2, 1/2 - k/2, 1/2 - k/2 - v$, where $w = 3/2 - k/2 - s$. We thus write

$$\mathcal{P}(s) = (s-1)(s-1-v)\prod_{0\le r\le k/2}(s - 3/2 + k/2 - r - v/2)$$

and set

$$\tilde{\mathcal{M}}(s,w,v) = \mathcal{P}(s)\mathcal{M}(s,w,v).$$

The poles of $\tilde{\mathcal{M}}(s,w,v)$ are contributed entirely by the cuspidal part, and we obtain

$$\frac{1}{2\pi i}\int_{(1+\delta)}\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{s-1+k/2}ds=\tilde{\mathcal{M}}(1-k/2,1/2,v)$$
$$+\sum_{j,0\leq r\leq k/2}\mathrm{Res}_{s=1/2-r+it_j}\,\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{s-1+k/2}$$
$$+\sum_{j,0\leq r\leq k/2}\mathrm{Res}_{3/2-k/2-s=1/2-r+it_j-v/2}\,\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{s-1+k/2}$$
$$+\frac{1}{2\pi i}\int_{(1/2-k/2-\delta)}\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{s-1+k/2}ds.$$

Thus

$$-i\int_{(\epsilon)}\frac{e^{\pi v^2/y^2}X^v}{y}\mathcal{M}(1-k/2,1/2,v)dv$$
$$=I_1(X,y)-I_2(X,y)-I_3(X,y)-I_4(X,y), \quad (12.22)$$

where

$$I_1(X,y)$$
$$=-i\int_{(\epsilon)}\frac{e^{\pi v^2/y^2}X^v}{y}\frac{1}{2\pi i}\int_{(1+\delta)}\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{\mathcal{P}(1-k/2)(s-1+k/2)}dsdv, \quad (12.23)$$

$$I_2(X,y)=-i\int_{(\epsilon)}\frac{e^{\pi v^2/y^2}X^v}{y}$$
$$\times\sum_{j,0\leq r\leq k/2}\mathrm{Res}_{s=1/2-r+it_j}\,\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{\mathcal{P}(1-k/2)(s-1+k/2)}dv, \quad (12.24)$$

$$I_3(X,y)=-i\int_{(\epsilon)}\frac{e^{\pi v^2/y^2}X^v}{y}$$
$$\times\sum_{j,0\leq r\leq k/2}\mathrm{Res}_{3/2-k/2-s=1/2-r+it_j-v/2}\,\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{\mathcal{P}(1-k/2)(s-1+k/2)}dv, \quad (12.25)$$

and

$$I_4(X,y) \\ = -i\int_{(\epsilon)}\frac{e^{\pi v^2/y^2}X^v}{y}\frac{1}{2\pi i}\int_{(1/2-k/2-\delta)}\frac{\tilde{\mathcal{M}}(s,3/2-k/2-s,v)e^{(s-1+k/2)^2}}{\mathcal{P}(1-k/2)(s-1+k/2)}dsdv, \tag{12.26}$$

We will now estimate these four integrals from above, starting with the residual contributions. Referring to the evaluations in Proposition 11.4 and specializing the variables gives us

$$\begin{aligned}\mathrm{Res}_{s=1/2-r+it_j}\,\mathcal{M}(s,3/2-k/2-s,v) &= \frac{(\ell_1\ell_2)^{(k-1)/2}}{\ell^{1/2}} \\ &\times \mathcal{P}_\ell(1/2,v)\frac{2^{2k-1}\pi^k L(1/2,\overline{u_j})L(1/2+v,\overline{u_j})}{\zeta(1+v)\Gamma(k-1/2-r+it_j)} \\ &\times \overline{\rho_j(-1))}\overline{\langle V,u_j\rangle}\frac{(-1)^r\Gamma(1/2-it_j+r)\Gamma(2it_j-r)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)}.\end{aligned} \tag{12.27}$$

$$\begin{aligned}&\mathrm{Res}_{3/2-k/2-s=1/2-r+it_j-v/2}\,\mathcal{M}(s,3/2-k/2-s,v) \\ &\qquad = -\frac{\Gamma((k+1+v)/2)\Gamma(1/2+it_j-r)\Gamma(1/2-it_j+r)}{\Gamma(1/2+it_j-r-v/2)\Gamma(1/2+it_j-r+v/2)} \\ &\qquad\times\frac{(-1)^r\Gamma(2it_j-r)\overline{\rho_j(-1)\langle U,u_j\rangle}L_{\ell_2}(1/2+v/2,\bar{g}\otimes\bar{u}_j)}{r!\Gamma(1/2+it_j)\Gamma(1/2-it_j)}.\end{aligned} \tag{12.28}$$

To estimate $I_2(X,y)$ we will use the following

Lemma 12.5. *Let f be a function of v, analytic for $0 \le \mathfrak{Re}\, v < 3$, such that $f(it) \ll (1+|t|)^\epsilon$ for t real. Then*

$$\int_{(2)}\frac{L(1/2+i\gamma+v,\overline{u_j})e^{\pi v^2/y^2}f(v)X^v dv}{y} \ll y^\epsilon(1+|\gamma|)^{1/2+\epsilon}(1+|t_j|)^{1/2+\epsilon}.$$

Proof. We will integrate by parts. Set

$$F_1(v) = e^{\pi v^2/y^2}$$

and

$$F_2(v) = \int_0^v \frac{L(1/2+i\gamma+v',\overline{u_j})f(v')X^{v'}dv'}{y}.$$

Then

$$I(y) = \int_{2-i\infty}^{2+i\infty}F_1(v)dF_2(v)dv = F_1(v)F_2(v)|_{2-i\infty}^{2+i\infty} - \int_{2-i\infty}^{2+i\infty}F_2(v)F_1'(v)dv.$$

The boundary term vanishes as $e^{\pi v^2/y^2}$ decays exponentially in the imaginary part of v. Moving the line of integration to $\mathfrak{Re}\, v = 0$ and differentiating F_1 we now have

$$I(y) = -\int_{-i\infty}^{i\infty} \frac{2vF_2(v)e^{\pi v^2/y^2}dv}{y^2}.$$

As $v = it$ and $f(it) \ll |t|^\epsilon$, it follows that

$$f(it)X^{it} \ll |t|^\epsilon.$$

Then, by Cauchy-Schwarz,

$$\begin{aligned} |F_2(iT)| &\ll \frac{(1+|t|)^\epsilon}{y} \int_0^T |L(1/2 + i\gamma + it, \overline{u_j})|dt \\ &\ll \frac{T^{1/2+\epsilon}}{y} \left(\int_0^T |L(1/2 + i\gamma + it, \overline{u_j})|^2 dt \right)^{1/2}. \end{aligned}$$

The weak Lindelöf on average result

$$\int_0^T |L(1/2 \;+\; i\gamma \;+\; it, \overline{u_j})|^2 dt \quad \ll \quad (T(1+|\gamma|)(1+|t_j|))^{1+\epsilon}$$

is easily obtainable by an application of Montgomery's mean value theorem for Dirichlet polynomials (after replacing the L-series by a Dirichlet polynomial of length approximately $\sqrt{(\gamma + T + t_j)(\gamma + T - t_j)}$).

It follows from this that

$$\begin{aligned} I(y) &\ll (1+|\gamma|)^{1/2+\epsilon}(1+|t_j|)^{1/2+\epsilon} \int_{-\infty}^{\infty} \frac{t^{2+\epsilon}e^{-\pi t^2/y^2}}{y^3} dt \\ &\ll y^\epsilon (1+|\gamma|)^{1/2+\epsilon}(1+|t_j|)^{1/2+\epsilon}. \end{aligned}$$

□

Substituting (12.24) into (11.58) leads to

$$\begin{aligned} I_2(X,y) = -i \int_{(\epsilon)} \frac{e^{\pi v^2/y^2}X^v}{y} \\ \times \sum_{j, 0 \le r \le k/2} \frac{\mathcal{P}(1/2 - r + it_j)2^{2k-1}\pi^k L(1/2, \overline{u_j}) L(1/2 + v, \overline{u_j}) \overline{\rho_j(-1)}) \overline{\langle V, u_j \rangle}}{\mathcal{P}(1 - k/2)\zeta(1+v)\Gamma(k - 1/2 - r + it_j)} \\ \times \frac{(-1)^r \Gamma(1/2 - it_j + r)\Gamma(2it_j - r)}{r!\Gamma(1/2 + it_j)\Gamma(1/2 - it_j)} \frac{e^{(k/2 - 1/2 - r + it_j)^2}}{k/2 - 1/2 - r + it_j} dv. \end{aligned}$$

By the lemma

$$I(y) = \int_{(\epsilon)} \frac{L(1/2+v, \overline{u_j}) e^{\pi v^2/y^2} X^v dv}{\zeta(1+v)y} \ll y^{\epsilon}(1+|t_j|)^{1/2+\epsilon}. \tag{12.29}$$

The sum over t_j converges super exponentially, and so

$$I_2(X,y) \ll \frac{y^{\epsilon}}{\ell^{1/2}}. \tag{12.30}$$

Turning to (12.25), and substituting (12.28), leads to

$$\begin{aligned} I_3(X,y) = i \int_{(\epsilon)} & \frac{e^{\pi v^2/y^2} X^v}{y} \\ \times \sum_{j, 0 \le r \le k/2} & \frac{\mathcal{P}(1-k/2+r-it_j)\Gamma((k+1+v)/2)\Gamma(1/2+it_j-r)\Gamma(1/2-it_j+r)}{\mathcal{P}(1/2-k/2)\Gamma(1/2+it_j-r-v/2)\Gamma(1/2+it_j-r+v/2)} \\ \times & \frac{(-1)^r \Gamma(2it_j - r)\overline{\rho_j(-1)\langle U, u_j\rangle} L_{\ell_2}(1/2+v/2, \bar{g} \otimes \bar{u}_j)}{r! \Gamma(1/2+it_j)\Gamma(1/2-it_j)} \frac{e^{(r-it_j+v/2)^2}}{r-it_j+v/2} dv, \end{aligned}$$

Recall from (11.36) that

$$\sum_{|t_j| \sim T} \overline{\rho_j(-1)\, \langle U, u_j\rangle} \ll \ell^{(1-k)/2}(\log(1+\ell))^{1/2} \tag{12.31}$$

The super exponential decay of $e^{(r-it_j+v/2)^2}$ effectively restricts v to $|v/2 - it_j| \ll 1$ and by Stirling's formula, the resulting gamma factors decay on the order of $e^{-(\pi/2)|t_j}|$. Thus after shifting the integral to $\mathfrak{Re}\, v = 0$, we see that

$$I_3(X,y) \ll y^{-1+\epsilon} \ell^{1/2-k/2}.(\log(1+\ell))^{1/2} \tag{12.32}$$

To estimate $I_1(X,y)$, note that on the line $\mathfrak{Re}\, s = 1+\delta$, the function $\mathcal{M}(s, 3/2-k/2-s, v)$ has $\mathfrak{Re}\, w = \mathfrak{Re}\,(3/2-k/2-s) = 1/2-k/2-\delta$. We would like to use the spectral expansion valid in the region $\mathfrak{Re}\, s > 1$, $\mathfrak{Re}\, w < 1/2-k/2$ and $\mathfrak{Re}\, s+w+k/2+v/2 > 3/2$. To do this we choose any $\delta > 0$. However, the point $(1+\delta, 1/2-k/2-\delta)$ is right on the boundary, satisfying $s+w+k/2 = 3/2$. We will there do our estimation in the region $\mathfrak{Re}\, s = 1+\delta+\epsilon_1$, $\mathfrak{Re}\, w = 1/2-k/2-\delta-\epsilon_2$, for $\epsilon_1, \epsilon_2, \epsilon > 0$ satisfying $\epsilon_2/\epsilon_1 = 1-\epsilon$. This point lies in the region of convergence, and the line connecting it with $(1+\delta, 1/2-k/2-\delta)$ hits the other region of convergence, where $\mathfrak{Re}\, s < 1/2-k/2$, because the slope of the line will have absolute value less than 1 but greater than 1/2. As a result, by convexity, the upper bound at the desired point will be within epsilon of the upper bound we derive.

In this region, the function $\mathcal{M}(s, 3/2-k/2-s, v)$ breaks up into three pieces. Therefore we write

$$I_1(X,y) = \ell_1^{(k-1)/2}\ell_2^{1-s-w-k/2-v/2}2^{k-1-v}\pi^{k/2-1/2-v/2} \times (I_{1,1}(X,y)+I_{1,2}(X,y)) - I_{1,3}(X,y), \quad (12.33)$$

where for $i=1,2$

$$I_{1,i}(X,y) = i\int_{(\epsilon)} \frac{e^{\pi v^2/y^2}X^v\mathcal{P}(s)}{y\mathcal{P}(1/2-k/2)} \times \frac{1}{2\pi i}\int_{(1+\delta+\epsilon)} \frac{e^{(s-1+k/2)^2}}{s-1+k/2}\frac{\Gamma((k+1+v)/2)\Gamma(w+v/2)}{\Gamma(w+(k-1)/2)\Gamma(w+(k-1)/2+v)} \times \Gamma(1-w-v/2)\mathcal{M}_i(s,w,v)dsdv \quad (12.34)$$

and for $i=3$,

$$I_{1,3}(X,y) = i\int_{(\epsilon)} \frac{e^{\pi v^2/y^2}X^v\mathcal{P}(s)}{y\mathcal{P}(1/2-k/2)} \times \frac{1}{2\pi i}\int_{(1+\delta+\epsilon)} \frac{e^{(s-1+k/2)^2}}{s-1+k/2}\mathcal{M}_3(s,w,v)dsdv. \quad (12.35)$$

The $\mathcal{M}_i(s,w,v)$ are defined in Proposition 11.4. Substituting for $\mathcal{M}_1$ gives

$$I_{1,1}(X,y) = i\int_{(\epsilon)} \frac{e^{\pi v^2/y^2}X^v\mathcal{P}(s)}{y\mathcal{P}(1/2-k/2)} \times\frac{1}{2\pi i}\int_{(1+\delta+\epsilon)} \frac{e^{(s-1+k/2)^2}}{s-1+k/2}\frac{\Gamma((k+1+v)/2)\Gamma(3/2-k/2-s+\epsilon+v/2)}{\Gamma(1-s+\epsilon)\Gamma(1-s+\epsilon+v)} \times\Gamma(s+k/2-1/2-v/2-\epsilon)\sum_j L_{\ell_2}(1/2+\epsilon+v/2,\bar{g}\otimes\bar{u}_j)\overline{\rho_j(-1)\langle U,u_j\rangle} \times \frac{\Gamma(1-k/2-s+v/2+\epsilon+it_j)\Gamma(1-k/2-s+v/2+\epsilon-it_j)}{\Gamma(1/2+it_j)\Gamma(1/2-it_j)}dsdv \quad (12.36)$$

The sum over j converges absolutely in this range. The super exponential decay in s effectively restricts the range of s integration to $s \ll 1$ and the ratio of gamma factors decays exponentially in the imaginary part of v. Recalling the estimate in (12.31) we get

$$I_{1,1}(X,y) \ll X^\epsilon \quad (12.37)$$

Similarly, $I_{1,2}(X,y) \ll X^\epsilon$.

We now substitute for $\mathcal{M}_3$ in (12.35) as follows.

$$\begin{aligned}
I_{1,3}(X,y) = i\int_{(\epsilon)} &\frac{e^{\pi v^2/y^2} X^v \mathcal{P}(s)}{y\mathcal{P}(1/2-k/2)} \\
&\times \frac{1}{2\pi i}\int_{(1+\delta+\epsilon)} \frac{e^{(s-1+k/2)^2}}{s-1+k/2} \\
\times &\frac{2^{2k-1-v}\pi^k(\ell_1\ell_2)^{(k-1)/2}\Gamma(3/2-k/2-s+\epsilon+v/2)\Gamma(s+k/2-1/2-\epsilon-v/2)}{\ell^{s'}\Gamma((1+v-k)/2)\Gamma((1-v+k)/2)\mathcal{V}\zeta(1+2\epsilon+v)} \\
&\times \sum_{j\geq 1}\left(\frac{L(1/2+\epsilon,\bar{u}_j)L(1/2+\epsilon+v,\bar{u}_j)}{\Gamma(s)\Gamma(s+k-1)}\right. \\
&\left.\times\, \Gamma(s-1/2+it_j)\Gamma(s-1/2-it_j)\right) ds dv. \qquad (12.38)
\end{aligned}$$

By Stirling's formula

$$\frac{\Gamma(3/2-k/2-s+\epsilon+v/2)\Gamma(s+k/2-1/2-\epsilon-v/2)}{\Gamma((1+v-k)/2)\Gamma((1-v+k)/2)} \ll_s y^{\epsilon},$$

both with a polynomial dependence on s. The line of integration can be pulled back to $\mathfrak{Re}\, v = 0$, and it is known that

$$\zeta(1+it)^{-1} \ll (1+|t|)^{\epsilon}.$$

As the sum over t_j converges absolutely, and the decay in the imaginary part of s is super exponential, it follows from Lemma 12.5 that

$$I_{1,3}(X,y) \ll y^{\epsilon}\ell^{-1/2}.$$

12.3. **The case $f = g$.** Recall that when $f = g$,

$$\begin{aligned}
S(Q) &= \mathrm{Res}_{s=0}\, L_Q(s+1, f\otimes f)\Gamma(s)(x/2)^s \\
&+2\sum_{d|Q}\frac{\mu(d)\,\mathrm{Res}_{w=0}\,\Gamma(w)x^w Z_Q(1-k/2, w+1/2; F_d, F_d))}{d} + \mathcal{O}\left(Q^{\theta-1/2}\right).
\end{aligned} \qquad (12.39)$$

In this case, for any $x > 1$,

$$S(X,y) = S_1(X,y) + 2S_2(X,y) + \mathcal{O}\left(\frac{X^{1/2}}{y}\right),$$

where

$$S_1(X,y) = -i\int_{(2)} \frac{e^{\pi v^2/y^2}X^v}{y}\sum_{Q\geq 1}\frac{\mathrm{Res}_{s=0}\, L_Q(s+1,f\otimes f)\Gamma(s)(x/2)^s}{Q^v} dv \qquad (12.40)$$

and

$$
\begin{aligned}
S_2(X,y) = -i \int_{(2)} \frac{e^{\pi v^2/y^2} X^v}{y} \\
\times \sum_{Q\geq 1} \frac{\sum_{d|Q} \frac{\mu(d)}{d} \operatorname{Res}_{w=0} \Gamma(w) x^w Z_{Q/d}(1-k/2, w+1/2; F_d, F_d)}{Q^v} dv.
\end{aligned}
\tag{12.41}
$$

As before, the simplest of the two is $S_1(X,y)$. The sum over Q in (12.40) can be analyzed just as in the case $f \neq g$, the only difference being that there will be a main term which is a multiple of $\log x$, owing to the presence of a double pole at $s = 0$. The result is

$$
\begin{aligned}
&\sum_{Q\geq 1} \frac{\operatorname{Res}_{s=0} L_Q(s+1, f\otimes f)\Gamma(s)(x/2)^s}{Q^v} \\
&\qquad = \frac{\zeta(v)L(1,f,\vee^2)}{L(1+v,f)L(1+v,\bar{g})} E(v,f,f)\log x + \zeta(v)E_1(v,f,f),
\end{aligned}
$$

where $E(v,f,f), E_1(v,f,f)$ are analytic for $\mathfrak{Re}\, v > -1$.

Substituting in (12.40) gives

$$
\begin{aligned}
S_1(X,y) &= \frac{L(1,f,\vee^2)e^{\pi/y^2}X}{L(1,f)L(1,\bar{g})y} E(1,f,f)\left(\log x + c(f)\right) \\
&\quad - i\int_{(2)} \frac{e^{\pi v^2/y^2}X^v}{y} \frac{\zeta(v)L(1,f,\vee^2)}{L(1+v,f)L(1+v,\bar{g})} E(v,f,f)dv \\
&= \frac{L(1,f,\vee^2)e^{\pi/y^2}X}{L(1,f)L(1,\bar{g})y} E(1,f,f)\left(\log x + c(f)\right) + \mathcal{O}(X^\epsilon).
\end{aligned}
\tag{12.42}
$$

As before, the integral was estimated as $\mathcal{O}(X^\epsilon)$ by changing variables $v \to yv$ and applying stationary phase to save a power of $y^{1/2}$ from the ratio of gamma factors relating $\zeta(yv)$ to $\zeta(1-yv)$.

Turning to $S_2(X,y)$,

$$S_2(X,y) = -i\int_{(2)} \frac{e^{\pi v^2/y^2}X^v}{y} \times \sum_{Q\geq 1} \frac{\sum_{d|Q}\frac{\mu(d)}{d}\operatorname{Res}_{w=0}\Gamma(w)x^w Z_{Q/d}(1-k/2,w+1/2;F_d,G_d)}{Q^v}dv$$

$$= -i\int_{(2)} \frac{e^{\pi v^2/y^2}X^v}{y} \times \sum_{d\geq 1}\frac{\mu(d)}{d^{1+v}}\sum_{Q'\geq 1}\frac{\operatorname{Res}_{w=0}\Gamma(w)x^w Z_{Q'}(1-k/2,w+1/2;F_d,G_d)}{(Q')^v}dv$$

$$= -i\int_{(2)} \frac{e^{\pi v^2/y^2}X^v}{y} \times \sum_{d\geq 1}\frac{\mu(d)}{d^{1+v}}\operatorname{Res}_{w=0}\Gamma(w)x^w M(1-k/2,w+1/2,v;F_d,G_d)dv. \quad (12.43)$$

Now move the v line of integration to $\mathfrak{Re}\, v=\epsilon$. The shifted integral is estimated in an identical way to the case $f\neq g$ and is $\mathcal{O}(X^\epsilon)$. A pole of $M(1-k/2,w+1/2,v;F_d,G_d)$ is passed over at $2s'+v=2$, i.e at $v=1-2w$. We are left with

$$S_2(X,y) = 2\pi\operatorname{Res}_{w=0}\frac{e^{\pi(1-2w)^2/y^2}X^{1-2w}}{y} \times\sum_{d\geq 1}\frac{\mu(d)}{d^{2-2w}}\Gamma(w)x^w\operatorname{Res}_{v=1-2w}M(1-k/2,w+1/2,v;F_d,G_d)+\mathcal{O}(X^\epsilon). \quad (12.44)$$

The function $\operatorname{Res}_{v=1-2w}M(1-k/2,w+1/2,v;F_d,G_d)$ has a simple pole at $w=0$. When multiplied by $\Gamma(w)$ the resulting function has a double pole at $w=0$ and we finally get

$$S_2(X,y) = 2\pi\frac{e^{\pi/y^2}X}{y}\kappa_f\left(\log(x/X^2)+c(f)\right)+\mathcal{O}(X^\epsilon),$$

where

$$\kappa_f = \sum_{d\geq 1}\frac{\mu(d)}{d^2}\operatorname{Res}_{w=0}\operatorname{Res}_{v=1-2w}M(1-k/2,w+1/2,v;F_d,G_d).$$

The coefficient of $\log x$ appearing in the approximation of $S_1(X,y)$ is

$$\frac{L(1,f,\vee^2)e^{\pi/y^2}X}{L(1,f)L(1,\bar{g})y}E(1,f,f)$$

As the sum $S_1(X,y)+2S_2(X,y)$ is independent of x, it must be the case that

$$\frac{L(1,f,\vee^2)e^{\pi/y^2}X}{L(1,f)L(1,\bar{g})y}E(1,f,f)+4\pi\frac{e^{\pi/y^2}X}{y}\kappa_f=0,$$

which implies

$$\kappa_f=-\frac{L(1,f,\vee^2)}{4\pi L(1,f)L(1,\bar{g})E(1,f,f)}.$$

Substituting, finally, in

$$S(X,y)=S_1(X,y)+2S_2(X,y)+\mathcal{O}\left(\frac{X^{1/2}}{y}\right),$$

we have

$$\begin{aligned} S(X,y)&=-4\pi\frac{e^{\pi/y^2}X\log X}{y}\kappa_f+\frac{C(f)X}{y}+\mathcal{O}\left(\frac{X^{1/2}}{y}\right)+\mathcal{O}(X^\epsilon)\\ &=\frac{L(1,f,\vee^2)e^{\pi/y^2}X\log X}{L(1,f)L(1,\bar{g})E(1,f,f)y}+\frac{C(f)X}{y}+\mathcal{O}\left(\frac{X^{1/2}}{y}\right)+\mathcal{O}(X^\epsilon), \end{aligned} \tag{12.45}$$

as claimed in the Theorem.

References

[1] J. Bernstein and A. Reznikov. Estimates of Automorphic Functions, Moscow Math. J. vol 4, number 1]ages 19–37.

[2] V. Blomer. Shifted convolution sums and subconvexity bounds for automorphic L-functionsIMRN, 73:3905–3924 (2004)

[3] V. Blomer and F. Brumley. On the Ramanujan conjecture over number fields, Annals of Math. 174 (2011), 581-605

[4] V. Blomer and G. Harcos. Hybrid bounds for twisted L-functions J. Reine Angew. Math. Volume 2008, Issue 621 pp 53–79

[5] V. Blomer, and G. Harcos. Twisted L-functions over number fields and Hilbert's eleventh problem, GAFA 20 (2010), 1-52

[6] B. Conrey, and H. Iwaniec. The cubic moment of central values of automorphic L-functions, Annals of Mathematics, 151 (2000), 11751216

[7] M. Deshouillers and H. Iwaniec. Kloosterman sums and Fourier coefficients of cusp forms, Invent. Math. 70 (1982), no. 2, 219288.

[8] W. Duke, J. Friedlander, IH. Iwaniec. Bounds for automorphic L-functions II, Invent. Math. 115 (1994) 219239.

[9] W. Duke, J. Friedlander, IH. Iwaniec. Bounds for automorphic L-functions III, Invent. Math. 143 (2001) 221248.

[10] W. Duke, J. Friedlander, IH. Iwaniec. The subconvexity problem for Artin L-functions, Invent. Math. 149 (2002) 489577.

[11] The second moment of Dirichlet twists of Hecke L-functions Peng Gao, Rizwanur Khan, and Guillaume Ricotta, arXiv:0812.2606v2 [math.NT] 26 Mar 2009

[12] D. Goldfeld. Analytic and Arithmetic Theory of Poincaré Series. Astérisque 61 (1979 95–107

[13] D. Goldfeld, J. Hoffstein, D. Lieman. Appendix to Coefficients of Maass forms and the Siegel Zero, Annals of Math, Vol 140 No.1

[14] I.S. Gradshteyn and I.M. Ryzhik. Table of Integrals, Series and Products, fifth edition, Academic Press Inc.

[15] D. Hansen. Test vectors in global trilinear forms and triple product L-functions, preprint

[16] T. Hulse. Thesis (in preparation)

[17] D. R. Heath-Brown. The fourth power mean of Dirichlets L-functions. Analysis 1 (1981), no. 1 2532.

[18] A. Ichino. Trilinear forms and the central values of triple product L-functions, Duke Math. J. 145:281-307 (2008)

[19] H. Iwaniec,and E. Kowalski. Analytic Number Theory, AMS Colloquium Publications, volume 53

[20] R. Khan. Simultaneous non-vanishing of $GL(3) \times GL(2)$ and GL(2) L-functions, Arxiv, Number Theory 1103.5937, Theorem 1.1

[21] H. H. Kim. Functoriality for the exterior square of $GL(4)$ and the symmetric fourth of $GL(2)$, J. Amer. Math. Soc. 16 (2003), no. 1, 139183, with Appendix 1 by D. Ramakrishnan and Appendix 2 by H. Kim and P. Sarnak.

[22] H. Kim, F. Shahidi. Cuspidality of symmmetric powers with applications, Duke Math. J. 112 (2002), 177197.

[23] Y-K Lau, J. Liu and Y. Ye. Shifted convolution sums of Fourier coefcients of cusp forms, Number theory, 108-135, Ser. Number Theory Appl., 2 World Sci. Publ., Hackensack, NJ 2007

[24] Y-K Lau, J. Liu and Y. Ye. Subconvexity bounds for RankinSelberg L-functions for congruence subgroups, Journal of Number Theory 121 (2006) 204223

[25] R.A. Rankin. Contributions to the theory of Ramanujan's function $\tau(n)$ and similar arithmetic functions, Proc. Cambridge Philos. Soc. 35 (1939) 357-372

[26] P. Sarnak. Estimates for Rankin-Selberg L-functions and quantum unique ergodicity, Journal of Functional Analysis 184, 419–453 (2001)

[27] A. Selberg. Bemerkungenüber eine Dirichletsche Reihe, die mit der Theorie der Modulformen nahe Verbunden ist, Arch. Math. Naturrid, 43 (1940), 44–50

[28] A. Selberg. On the Estimation of Fourier Coefficients of Modular Forms, Proc. Symp. Pure Math, AMS, Vol III, Theory of Numbers (1965) 1–15

[29] K. Soundararajan. The fourth moment of Dirichlet L-functions, Analytic number theory, Clay Math. Proc., vol. 7, Amer. Math. Soc., Providence, RI, 2007, pp. 239246.

[30] K. Soundararajan, M. Young.The second moment of quadratic twists of modular L-functions. J. Eur. Math. Soc. (JEMS) 12 (2010), no. 5, 10971116.

[31] T Stefanicki. Non-vanishing of L-functions attached to automorphic representations of GL(2) over Q, J. Reine Angew. Math. 474 (1996), 124.

[32] T.C. Watson. Rankin triple products and quantum chaos, arXiv:0810.0425v3 [math.NT]

[33] M. Young. The fourth moment of Dirichlet L-functions Pages 1-50 from Annals of Math. Volume 173 (2011), Issue 1 1–50

E-mail address: `jhoff@math.brown.edu`

Department of Mathematics, Brown University, Providence, RI, 02912